

**ARIZONA GAME AND FISH DEPARTMENT  
HERITAGE DATA MANAGEMENT SYSTEM**

**Animal Abstract**

**Element Code:** AFCJB37050

**Data Sensitivity:** No

**CLASSIFICATION, NOMENCLATURE, DESCRIPTION, RANGE**

**NAME:** *Rhinichthys osculus*

**COMMON NAME:** Speckled Dace

**SYNONYMS:** *Rhinichthys nubilus*, *Apocope oscula oscula*, *Agosia couesii*, *Argyreus osculus*

**FAMILY:** Cyprinidae

**AUTHOR, PLACE OF PUBLICATION:** Girard, 1856, Proc. Acad. Nat. Sci. Philadelphia, 8:165-213.

**TYPE LOCALITY:** Species was described from Babocomari Creek, flood tributary of San Pedro River, just north of Fort Huachuca, Cochise County, Arizona (Lowe 1964).

**TYPE SPECIMEN:**

**TAXONOMIC UNIQUENESS:** Eight species of this genus in United States. *R. osculus* only species of *Rhinichthys* found in Arizona. Two subspecies of *R. osculus* found in Arizona; *R.o. osculus* inhabits the southern part of the Gila River System, and *R.o. yarrowi* in larger rivers and creeks to the north.

**DESCRIPTION:** Small minnow, rarely exceeding 7.6 cm (3.0 in.) in length. Body chunky, rounded, somewhat flattened ventrally; body depth usually about equal to length of head. Upper jaw protractile, or if a frenum is present, it is very small. Barbels usually present at sides of upper lips. Fins generally rounded, dorsal fin origin above or just behind pelvic-fin insertion. Moderately small scales, 60 to 90 along lateral line. Dorsal fin with 6 to 9 rays, usually 8. Anal fin with 7 rays (rarely 6). Pharyngeal teeth in two rows, 1, 4-4, or 2, 4-4, 2, or a combination there of.

Coloration highly variable, drab olivaceous with patterns ranging from large black blotches on body, through a single or double lateral band, to almost unicolored. Breeding males vivid red on bases of paired fins and on body near those fins, on and near anal fin base, the lower caudal lobe, the mouth, and near the upper part of gill cleft (Minckley 1973).

**AIDS TO IDENTIFICATION:** Small eye; a broad, blackish mid dorsal stripe; a conspicuous black spot at base of the caudal fin. *Rhinichthys osculus* differs from *Agosia chrysogaster*, longfin dace, in that *A. chrysogaster* has nearly terminal mouth, white spots at front and rear of dorsal fin, white bar on caudal peduncle, no scales on breast, belly and part of back; lacks barbel, groove on snout. Differs from *R. cataractae*, longnose dace, in that *R. cataractae* has a long fleshy snout in front of mouth, no groove separating snout from upper lip (Page and Burr 1991).

**ILLUSTRATIONS:** B&W photo (Minckley 1973:129)  
Color line drawing (Page and Burr 1991:100).  
Photo (USDA, Forest Service Rocky Mtn Forest and Range Experiment Station 1991:17)  
B&W photo (Wildlife Habitat Management Staff Group 1975:147).

**TOTAL RANGE:** Native to all major western drainages from the Columbia and Colorado rivers south to Sonora, Mexico (Lee et al. 1980).

**RANGE WITHIN ARIZONA:** In Arizona, found in Colorado, Bill Williams, and Gila River drainages, except slower and warmer portions of Colorado River mainstream. Minckley (AGFD Native Fish Diversity Review. 1995. Tempe, Arizona) states that there are 4 populations in the southern part of its range.

## **SPECIES BIOLOGY AND POPULATION TRENDS**

**BIOLOGY:** Where this species has been more thoroughly studied, as in the Great Basin, high levels of morphological and genetic variability have been found. In many cases, individual springs and isolated basins have been found to have unique species or subspecies (Simons 1987). In Arizona there are at least two major body forms, a small, highly speckled or blotched, chubby-bodied form found in the southern part of the Gila River system, and a larger, banded or unicolored, more streamlined form found north of the Mogollon Rim. Can persist for long periods in intermittent pool even though they are crowded, diseased and starving. Relatively low tolerance for water with high temperature or low oxygen content.

**REPRODUCTION:** Two breeding periods, one in spring and the other in late summer. Spawn over coarse substrate using broadcast spawning method. Congregate in large groups and release many eggs.

**FOOD HABITS:** Omnivorous feeder, subsists on algae and other plant material, small crustaceans, insect larvae, and small snails (McClane 1974).

**HABITAT:** A bottom dweller, found in rocky riffles, runs, and pools of headwaters, creeks, and small to medium rivers: rarely in lakes (Page and Burr 1991). Reside in water less than 0.5 m (1.6 ft.) deep, with current averaging about 0.4m/sec (1.3ft/sec). Often congregate below riffles and eddies. Breeding adults prefer swift water.

**ELEVATION:** Peak abundance found 2,000 to 3,000 m (6562 - 9843 ft.), rarely below 1,500 m (4921 ft.). Based on records in the Heritage Data Management System (HDMS), which does not include the entire range in Arizona, elevation ranges from 1,550 - 8,920 ft. (473 - 2721 m) (AGFD, unpublished data accessed 2001).

**PLANT COMMUNITY:**

**POPULATION TRENDS:** Stable

**SPECIES PROTECTION AND CONSERVATION**

|                                       |   |
|---------------------------------------|---|
| <b>ENDANGERED SPECIES ACT STATUS:</b> | None (USDI, FWS 1996)<br>[C2 USDI, FWS 1994]  |
| <b>STATE STATUS:</b>                  | None  |
| <b>OTHER STATUS:</b>                  | No Forest Service Status (USDA, FS Region<br>3 1999)<br>[Forest Service Sensitive, USDA, FS Region<br>3 1988]<br>Bureau of Land Management Sensitive<br>(USDI, BLM AZ 2000, 2005)<br>No NNDFW Status (NNDFW, NESL 2000)<br>[Group 4, NNDFW, NESL 1994]<br>Listed Endangered (Secretaría de Medio<br>Ambiente 2000)<br>[Listed Endangered Secretaría de Desarrollo<br>Social 1994] |

**MANAGEMENT FACTORS:** Speckled dace are widespread, abundant, and not in danger of extinction. Little management needed except to promote land use practices which maintain natural aquatic habitats. This species does not fare well in the presence of non-native predatory fish; introductions of these fishes should be prevented (Simons 1987).

**PROTECTIVE MEASURES TAKEN:****SUGGESTED PROJECTS:**

**LAND MANAGEMENT/OWNERSHIP:** BIA - Hualapai and San Carlos Reservations, and Navajo Nation; BLM - Arizona Strip, Kingman, Phoenix and Safford Field Offices; NPS - Grand Canyon National Park and Montezuma Castle National Monument; USFS - Apache-Sitgreaves, Coconino, Coronado, Prescott and Tonto National Forests; State Land Department; AGFD Black River Lands; Sonoita Creek State Natural Area; TNC - Aravaipa Canyon and Muleshoe Ranch Preserves, and Patagonia-Sonoita Creek; Private.

**SOURCES OF FURTHER INFORMATION****REFERENCES:**

- AGFD Native Fish Diversity Review. 1995. Tempe, Arizona.
- Eddy, S. and J. Underhill. 1978. How to know the freshwater fishes. W.C. Brown Company Publishers. p. 65.
- Lee, D.S., C.R. Gilbert, C.H. Hocutt, R.E. Jenkins, D.E. McAllister and J.R. Stauffer, Jr. 1980. Atlas of North America freshwater fishes. North Carolina State Museum of Natural History, Raleigh, North Carolina.

- Lowe, C.H. Editor. 1964. The vertebrates of Arizona. University of Arizona Press, Tucson, Arizona. p. 141.
- McClane, A. 1974. Field guide to freshwater fishes of North America. Holt, Rinehart and Winston. p. 77.
- Minckley, W.C. 1973. Fishes of Arizona. Arizona Game and Fish Department, Phoenix. pp. 129-132.
- Navajo Nation Department of Fish and Wildlife. 1994. Navajo Endangered Species List. p. 4.
- Navajo Nation Department of Fish and Wildlife. 2000. Navajo Endangered Species List.
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America, North of Mexico. Houghton Mifflin Co., Boston. pp. 100-101.
- Rinne, J.N. and W.L. Minckley. 1991. Native fishes of arid lands: a dwindling resource of the desert southwest. U.S. Department of Agriculture, Forest Service, General Technical Report RM-206. Fort Collins, Colorado. pp.16-17.
- Secretaría de Desarrollo Social. 1994. Diario Oficial de la Federacion. p. 53.
- Secretaría de Medio Ambiente. 2000. Diario Oficial de la Federacion, PROY-NOM-059-ECOL-2000. p. 46.
- Simons, L.H. 1987. Speckled Dace. Arizona Game and Fish Department, Nongame field notes. pp.1-3.
- USDA, Forest Service Region 3. 1988. Regional Forester's Sensitive Species List.
- USDA, Forest Service Region 3. 1999. Regional Forester's Sensitive Species List.
- USDI, Bureau of Land Management. 2000. Arizona BLM Sensitive Species List. Instruction Memorandum No. AZ-2000-018.
- USDI, Bureau of Land Management. 2005. Arizona BLM Sensitive Species List.
- USDI, Fish and Wildlife Service. 1994. Endangered and Threatened Wildlife and Plants; Animal Candidate Review for Listing as Endangered or Threatened Species; Proposed Rule. Federal Register 59(219):59000.
- USDI, Fish and Wildlife Service. 1996. Endangered and Threatened Wildlife and Plants: Review of Plant and Animal Taxa that are Candidates for Listing as Endangered or Threatened Species. Federal Register 61(40):7596-7613.
- Wildlife Habitat Management Staff Group. 1975. Endangered and unique fish and wildlife of the southwestern national forests. U.S. Department of Agriculture, Forest Service, Southwestern Region. p. 147.

**MAJOR KNOWLEDGEABLE INDIVIDUALS:****ADDITIONAL INFORMATION:**

**Revised:** 1994-07-19 (LOC)  
 1994-07-25 (SMS)  
 1995-01-29 (KLY)  
 1997-03-05 (SMS)  
 2001-10-18 (SMS)  
 2002-12-04 (RHB)

To the user of this abstract: you may use this entire abstract or any part of it. We do request, however, that if you make use of this abstract in plans, reports, publications, etc. that you credit the Arizona Game and Fish Department. Please use the following citation:

Arizona Game and Fish Department. 20XX (= **year of last revision as indicated at end of abstract**). X...X (= **taxon of animal or plant**). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. X pp.