

Three-Year Summary of Fish Salvaged at the TFCF

These data represent the period of April 1, 2000 through March 31, 2003. For this period, average pumping at the Tracy Pumping Plant was 2,480,000 acre-ft/year and total fish salvage was 6,520,000 fish/year. Daily salvage estimates were based on 10-minute counts performed at the TFCF and reported to the California Department of Fish and Game. Data are rounded to three significant figures.

Common Name	Genus and Species	Family	Fish Salvaged/ Year, 3- Year Average	3-Year Percent of Fish Salvaged
Threadfin shad	<i>Dorosoma petenense</i>	Clupeidae	4,550,000	69.8%
Striped bass	<i>Morone saxatilis</i>	Moronidae	771,000	11.8%
American shad	<i>Alosa sapidissima</i>	Clupeidae	541,000	8.30%
Yellowfin goby	<i>Acanthogobius flavimanus</i>	Gobiidae	270,000	4.14%
White catfish	<i>Ameiurus catus</i>	Ictaluridae	119,000	1.83%
Bluegill	<i>Lepomis macrochirus</i>	Centrarchidae	79,100	1.21%
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	Salmonidae	31,900	0.488%
Inland silverside	<i>Menidia beryllina</i>	Atherinidae	30,300	0.465%
Largemouth bass	<i>Micropterus salmoides</i>	Centrarchidae	24,300	0.373%
Splittail	<i>Pogonichthys macrolepidotus</i>	Cyprinidae	23,200	0.356%
Delta smelt	<i>Hypomesus transpacificus</i>	Osmeridae	19,300	0.296%
Channel catfish	<i>Ictalurus punctatus</i>	Ictaluridae	17,200	0.264%
Longfin smelt	<i>Spirinchus thaleichthys</i>	Osmeridae	16,100	0.246%
Golden shiner	<i>Notemigonus crysoleucas</i>	Cyprinidae	7,410	0.114%
Prickly sculpin	<i>Cottus asper</i>	Cottidae	4,770	0.0732%
Steelhead trout	<i>Oncorhynchus mykiss</i>	Salmonidae	4,350	0.0668%
Black crappie	<i>Pomoxis nigromaculatus</i>	Centrarchidae	1,420	0.0217%
Redear sunfish	<i>Lepomis microlophus</i>	Centrarchidae	1,220	0.0187%
Lampreys, all species			1,210	0.0186%
Shimofuri goby	<i>Tridentiger bifasciatus</i>	Gobiidae	1,180	0.0180%
Bigscale logperch	<i>Percina macrolepida</i>	Percidae	1,050	0.0160%
Warmouth	<i>Lepomis gulosus</i>	Centrarchidae	932	0.0143%
Tule perch	<i>Hysterocarpus traski</i>	Embiotocidae	460	<0.010%
Threespine stickleback	<i>Gasterosteus aculeatus</i>	Gasterosteidae	292	< 0.010%
Brown bullhead	<i>Ameiurus nebulosus</i>	Ictaluridae	280	<0.010%
Sacramento sucker	<i>Catostomus occidentalis</i>	Catostomidae	192	< 0.010%
Mosquitofish	<i>Gambusia affinis</i>	Poeciliidae	188	< 0.010%

Common Name	Genus and Species	Family	Fish Salvaged/ Year, 3- Year Average	3-Year Percent of Fish Salvaged
Common carp	<i>Cyprinus carpio</i>	Cyprinidae	173	< 0.010%
Wakasagi	<i>Hypomesus nipponensis</i>	Osmeridae	148	< 0.010%
Pacific staghorn sculpin	<i>Leptocottus armatus</i>	Cottidae	76	< 0.010%
Fathead minnow	<i>Pimephales promelas</i>	Cyprinidae	72	< 0.010%
White crappie	<i>Pomoxis annularis</i>	Centrarchidae	64	< 0.010%
Striped mullet	<i>Mugil cephalus</i>	Mugilidae	44	< 0.001%
Riffle sculpin	<i>Cottus gulosus</i>	Cottidae	40	< 0.001%
Starry Flounder	<i>Platichthys stellatus</i>	Pleuronectidae	36	< 0.001%
Red shiner	<i>Cyprinella lutrensis</i>	Cyprinidae	28	< 0.001%
Goldfish	<i>Carassius auratus</i>	Cyprinidae	24	< 0.001%
Sacramento blackfish	<i>Orthodon microlepidotus</i>	Cyprinidae	24	< 0.001%
Blue catfish	<i>Ictalurus furcatus</i>	Ictaluridae	16	< 0.001%
Sacramento pikeminnow	<i>Ptychocheilus grandis</i>	Cyprinidae	12	< 0.0001%
Black bullhead	<i>Ameiurus melas</i>	Ictaluridae	9	< 0.001%
Chameleon goby	<i>Tridentiger trigonocephalus</i>	Gobiidae	8	< 0.001%
Green sturgeon	<i>Acipenser medirostris</i>	Acipenseridae	8	< 0.001%
Hardhead	<i>Mylopharodon conocephalus</i>	Cyprinidae	4	< 0.001%
Hitch	<i>Lavinia exilicauda</i>	Cyprinidae	4	< 0.001%
Miscellaneous Species			4	< 0.001%