



# Weekly Assessment of CVP and SWP Delta Operations on ESA-listed Species

January 16, 2024

## Executive Summary

### Operational Conditions

See Weekly Fish and Water Operation Outlook document for January 16 – 22.

### Winter-run Chinook Salmon

Loss of natural winter-run Chinook Salmon (by length at date, LAD) has occurred in the past week at the State or Federal fish salvage facilities. Loss of natural winter-run Chinook Salmon at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities is likely to occur over the next week. 35-45% of juvenile natural winter-run Chinook Salmon from brood year (BY) 2023 are estimated to be present in the Delta.

### Spring-run Chinook Salmon

No loss of natural YOY spring-run Chinook Salmon (by length at date, LAD) has occurred in the past week at the State or Federal fish salvage facilities. 15-30% of juvenile natural spring-run Chinook Salmon was estimated in the Delta. CV spring-run Chinook Salmon adults have completed spawning and eggs are in gravel and are emerging. Fry are beginning to migrate downstream. Loss of genetically confirmed yearling spring-run Chinook Salmon occurred on 1/11/24.

### Central Valley Steelhead

Loss of natural California CV (CCV) steelhead has occurred in the past week at the Federal or State fish salvage facilities. Loss of CCV steelhead at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities is likely to occur over the next week. 5-10% of CCV steelhead were estimated in the Delta.

### DCC gates recommendation

The DCC gates were closed for the OMR Season on 11/27/2023.

## Delta Smelt

Based on distribution patterns over the past decade and low detections in this water year, Delta Smelt are unlikely to be prevalent in the Central and South Delta. Limited detection data from the past month supports Delta Smelt presence in the lower Sacramento River. Based on historical data, the centroid of distribution may have shifted downstream with shifting X2 (Sommer et al. 2011). The last Delta Smelt observation was on 1/16/2024 in the lower Sacramento River. The likelihood of Delta Smelt entrainment is low due to seasonal timing. OMR Management based on Salmonid presence has begun as of 1/1/2024. The Integrated Early Winter Pulse Protection (IEWPP) period began on 12/1/2023. “First Flush” conditions that would trigger IEWPP regulations are not anticipated but will be monitored this week.

## Monitoring Teams summary

There were no non-consensus issues to report from the Salmon Monitoring Team or Smelt Monitoring Team.

## Operational and Regulatory Conditions

See current Weekly Fish and Water Operation Outlook document.

## Biology, Distribution, and Evaluation Winter-run Chinook salmon, Spring-run Chinook salmon, Central Valley Steelhead

### Population Status

#### *Winter-run Chinook Salmon*

- Delta Life Stages:
  - Juveniles, Adults
- Brood Year 2023 Productivity:
  - Natural winter-run Chinook salmon: The Final WR JPE for BY2023 is 234,896.
  - Mean cumulative weekly passage of winter-run Chinook salmon through December 31st at Red Bluff Diversion Dam (RBDD) for the last 21 years of passage data is 96.7% (one SD of 3.5%). By 12/31/2023, 1,064,812 winter-run Chinook salmon were estimated to have passed RBDD compared to the total cumulative passage for WY 2023 of 240,059 winter-run Chinook salmon.
  - Hatchery winter-run Chinook salmon: One release of Livingston Stone National Fish Hatchery CWT tagged winter-run Chinook salmon occurred on 12/28/23.

### ***Spring-run Chinook Salmon***

- Delta Life Stages:
  - Young-of-year (YOY) and Yearlings
- Brood Year 2023 Productivity:
  - Natural spring-run Chinook salmon: No JPE has been established for spring-run Chinook salmon.
  - Hatchery spring-run Chinook salmon surrogates associated with the Proposed Action (PA 4.10.5.10.2 Additional Real-Time OMR Restrictions and Performance Objectives):
    - Approximately 60,764 brood year 2023 late-fall Chinook salmon were released on December 22, 2023 into Battle Creek at the Coleman NFH. This group is 100% marked with an adipose-fin clip and coded-wire tagged.
    - Approximately 71,049 Coleman NFH brood year 2023 late-fall Chinook Salmon were released on December 29, 2023 into Battle Creek at Coleman NFH. This group is 100% marked (with an adipose-fin clip and CWT) and has an overall estimated average fork length of 145 mm.
    - Approximately 67,018 Coleman NFH brood year 2023 late-fall Chinook Salmon were released on January 11, 2023 into Battle Creek at Coleman NFH. This group is 100% marked (with an adipose-fin clip and CWT) and has an overall estimated average fork length of 145 mm.

### ***Central Valley Steelhead***

- Delta Life Stages:
  - Spawning Adults, Kelts, Juveniles
- Brood Year 2023 Productivity:
  - Spawner abundance: There is limited information about the adult steelhead population. It is estimated to be small, contributing to the limited productivity of the population.
  - Natural steelhead: No JPE has been established for steelhead. Data are limited.
  - Hatchery steelhead: Reclamation's Proposed Action has no hatchery steelhead triggers.

### **Distribution**

#### ***Winter-run Chinook Salmon***

### **Current Distribution:**

- For winter-run Chinook Salmon observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.
- As of 1/16/2024, no genetic natural winter-run Chinook Salmon have been identified at the CVP or SWP facilities.

### **Historical Trends**

- For historical winter-run Chinook salmon trends in salvage, see Table 3.
- Loss of natural winter-run Chinook salmon at the CVP and SWP fish collection facilities is likely to occur over the next week based on hydrology. Based on historic trends in salvage winter-run Chinook salmon loss may increase over the next week.

### **Forecasted Distribution within Central Valley and Delta regions**

- The STARS model projects route-specific proportion of entrainment, survival, and travel times (Table 5). This model does not estimate entrainment into the lower Sacramento River sloughs (i.e., Three-Mile Slough).

### ***Spring-run Chinook salmon***

#### **Current Distribution**

- For spring-run Chinook salmon observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.

#### **Historical Trends**

- For historical spring-run Chinook salmon trends in salvage, see Table 3. Based on historic trends in salvage YOY spring-run Chinook salmon loss may increase over the next week.

#### **Forecasted Distribution within Central Valley and Delta regions**

- YOY spring-run Chinook are migrating into the Delta.

### ***Central Valley Steelhead***

#### **Current Distribution**

- For CCV Steelhead observations reported to SaMT since previous meeting, see Table 1.
- For SaMT distribution estimates, see Table 2.

- As of 1/16/2024, 7 unclipped CCV steelhead have been identified at the CVP or SWP facilities.

### Historical Trends

- For historical CCV steelhead trends in salvage, see Table 2. Based on historic trends in salvage, juvenile CCV steelhead loss may occur over the next week.

### Forecasted Distribution within Central Valley and Delta regions

- The entrainment tool estimates of CCV steelhead loss to be low (Table 6, Fig. 1).

Table 1. Fish observation reported since the previous SaMT meeting. NAs represent no data reported. See Operations Outlook for notes on interruptions in any surveys.

Locations	Reporting Period	SR Chinook	WR Chinook	LFR Chinook	Steelhead (Wild)	Green Sturgeon
Butte Creek RST	12/11-1/1	1685	0	0	0	0
Tisdale RST	1/2-1/7	73	72	0	1	0
Knights Landing RST	1/7-1/13	5	16	0	1	0
Lower Sacramento RST	12/29-1/7	189	54	2	3	0
Beach Seines	12/31-1/6	5	9	1	0	0
Sac. Trawl	12/31-1/6	0	0	1	0	0
Chipps Island Midwater Trawl	12/31-1/6	0	0	0	0	0
Mossdale Kodiak Trawl	12/31-1/6	0	0	0	0	0
EDSM	12/31-1/6	0	1	1	2	0
Feather River Herring RST	12/31-1/7	0	0	0	0	0
Feather River Eye Side RST	12/31-1/7	0	0	0	0	0
Lower Feather River	1/8-1/12	0	0	0	0	0

Table 2. Salmonid distribution estimates

Location	Yet to Enter Delta (%)	In the Delta (%)	Exited Delta past Chipps Island (%)
Young-of-year (YOY) winter-run Chinook salmon	Current: 55-65% Last Week: 60-70%	Current: 35-45% Last Week: 30-40%	Current: 0% Last Week: 0%
YOY spring-run Chinook salmon	Current: 70-85% Last Week: 75-85%	Current: 15-30% Last Week: 15-25%	Current: 0% Last Week: 0%
YOY hatchery winter-run Chinook salmon	Current: 80-95% Last Week: 100%	Current: 5-20% Last Week: 0%	Current: 0% Last Week: 0%

Location	Yet to Enter Delta (%)	In the Delta (%)	Exited Delta past Chipps Island (%)
Natural origin steelhead	Current: 90-95% Last Week: 95-99 %	Current: 5-10% Last Week: 1-5%	Current: 0% Last Week: 0%

Table 3. Historic migration and salvage patterns. Last updated 01/16/2023.

Species	Red Bluff Diversion Dam	Tisdale Rst	Knights Landing Rst	Sac Trawl Sherwood Catch Index	Chipps Island Trawl Catch Index	Salvage
Chinook, Winter-run, Unclipped	97.8%(96.3%,99.3%) BY: 2014 - 2022	80.5%(68.0%,93.1%) BY: 2014 - 2022	78.9%(58.4%,99.3%) BY: 2014 - 2022	39.8%(13.3%,66.4%) BY: 2014 - 2022	4.9%(-2.8%,12.6%) BY: 2014 - 2022	22.8%(4.8%,40.8%) WY: 2014 - 2023
Chinook, Spring-run, Unclipped	14.1%(3.9%,24.3%) BY: 2014 - 2022	20.6%(4.8%,36.3%) BY: 2014 - 2022	29.8%(7.7%,51.9%) BY: 2014 - 2022	5.2%(-2.8%,13.2%) BY: 2014 - 2022	0.0%(0.0%,0.0%) BY: 2014 - 2022	0.0%(0.0%,0.0%) WY: 2014 - 2023
Steelhead, Unclipped (January-December)	0.7%(-0.5%,2.0%) BY: 2014 - 2023	15.4%(0.4%,30.4%) BY: 2014 - 2023	15.5%(-2.4%,33.5%) BY: 2014 - 2023	2.5%(-3.4%,8.4%) BY: 2014 - 2023	3.4%(-1.7%,8.5%) BY: 2014 - 2023	N/A
Steelhead, Unclipped (December-March)	N/A	N/A	N/A	N/A	N/A	10.2%(-1.0%,21.4%) WY: 2014 - 2023
Steelhead, Unclipped (April-June)	N/A	N/A	N/A	N/A	N/A	N/A

Table 4. Mean daily flow and percent change (Wilkins Slough, Deer Creek, Mill Creek; cfs from CDEC) and temperature and percent change (Knights Landing; °F from RST).

Date	Mill Creek (MLM): mean daily flow (cfs)	Mill Creek (MLM): flow percent change	Mill Creek (MLM): Alert	Deer Creek (DCV): mean daily flow (cfs)	Deer Creek (DCV): flow percent change	Deer Creek (DCV): Alert	Wilkins Slough (WLK): mean daily flow (cfs)	Knights Landing RST: water temperature (f)	Alert Triggered
1/15/2024	298.5	-47.5%	Flow>95cfs	320.1	-17.0%	Flow>95cfs	17,231.5	N/A	N/A

Date	Mill Creek (MLM): mean daily flow (cfs)	Mill Creek (MLM): flow percent change	Mill Creek (MLM): Alert	Deer Creek (DCV): mean daily flow (cfs)	Deer Creek (DCV): flow percent change	Deer Creek (DCV): Alert	Wilkins Slough (WLK): mean daily flow (cfs)	Knights Landing RST: water temperature (f)	Alert Triggered
1/14/2024	568.2	123.8%	Flow>95cfs Change>50%	385.6	127.7%	Flow>95cfs Change>50%	9,106.2	N/A	N/A
1/13/2024	253.9	62.2%	Flow>95cfs Change>50%	169.4	8.1%	Flow>95cfs	9,023.7	N/A	N/A
1/12/2024	156.6	-14.3%	Flow>95cfs	156.6	-12.9%	Flow>95cfs	8,951.1	N/A	N/A
1/11/2024	182.7	11.6%	Flow>95cfs	179.9	8.7%	Flow>95cfs	8,856.1	N/A	N/A
1/10/2024	163.8	8.1%	Flow>95cfs	165.5	10.1%	Flow>95cfs	8,897.7	N/A	N/A
1/9/2024	151.4	0.2%	Flow>95cfs	150.3	-0.3%	Flow>95cfs	9,012.2	N/A	N/A

Table 5. STARS model simulations for route-specific entrainment, travel times, and survival. Travel time is calculated in days.

Stock	Date	Route	Median Travel Time	Survival	Routing Probability
Winter Chinook	2024-01-14	Overall	6.08	0.41	N/A
Winter Chinook	2024-01-14	Sacramento River	5.68	0.45	0.61
Winter Chinook	2024-01-14	Yolo Bypass	9.97	0.50	0.00
Winter Chinook	2024-01-14	Sutter Slough	5.92	0.37	0.13
Winter Chinook	2024-01-14	Steamboat Slough	5.31	0.49	0.13
Winter Chinook	2024-01-14	Interior Delta	9.03	0.17	0.13
Late-fall Chinook	2024-01-14	Overall	9.46	0.48	N/A
Late-fall Chinook	2024-01-14	Delta Cross Channel	N/A	N/A	0.00
Late-fall Chinook	2024-01-14	Georgiana Slough	13.57	0.24	0.23
Late-fall Chinook	2024-01-14	Sacramento River	8.18	0.60	0.48
Late-fall Chinook	2024-01-14	Sutter and Steamboat Slough	8.79	0.50	0.29

The entrainment tool estimates a median and maximum loss of winter-run Chinook Salmon and juvenile CCV Steelhead each week (Table 6a).

Table 6a-b. WY 2024 loss and salvage predictor data: Environmental details, current and forecast. Model results from 1/16/2024.

a) WY 2024 loss and salvage predictor data: Predicted weekly loss of winter-run Chinook salmon and steelhead at CVP and SWP facilities.

Parameter	Modeled Current Week	Modeled Next Week
Predicted Steelhead, Median %	0	0
Predicted Steelhead, High %	40	17
Predicted Chinook Winter Run, Median %	0	0
Predicted Chinook Winter Run, High %	7	9

b) Environmental details, current and forecast.

Parameter	Data	Forecast
Temperature (Mallard Island, C)	10.5	10.5
Precipitation (5-d running sum, inches)	0.19	0.19
Old and Middle River Flows (cfs)	-4874	-4874
Sacramento River Flow (Freeport, cfs)	17323	17323
DCC Gates	closed	closed
San Joaquin River Flow (Vernalis, cfs)	1824	1824
Export	6005	6005







Figure 1. Predicted weekly loss of steelhead and winter-run Chinook salmon at the CVP and SWP facilities

Figure 1 is two-line graphs of the predicted weekly loss of steelhead and winter-run Chinook salmon for water year 2024 beginning on December 1, 2023. The first line graph shows the cumulative loss of winter-run Chinook salmon comparing the predicted loss of over 50% and the observed loss of 3%. The second line graph shows the cumulative loss

of Steelhead salmon comparing the predicted loss of about 55% and the observed loss of about 9%.





Figure 2. Cumulative natural winter-run Chinook salmon loss for the year (blue) and 2009 – 2018 historic cumulative loss (gray, different symbols). Historic daily mean plotted in black circles.

Figure 2 is two line graphs showing the natural winter-run Chinook salmon cumulative loss for water year 2024 beginning on October 1, 2023, ending on June 30, 2024. Both

graphs combine observed Chinook salmon loss with historical loss calculated for water year 2009 through 2018.

The first line graph shows water year 2024 cumulative loss combined with the historical minimum and maximum weekly sum and water year 2024 cumulative loss combined with the historical daily mean. Water year 2024 cumulative loss combined with the historical daily minimum and maximum weekly sum reaches the 50 percent single-year threshold on January 1, 2024. Water year 2024 cumulative loss combined with the historical daily mean reaches the 50 percent single-year threshold on February 20, 2024.

The second line graph shows water year 2024 cumulative loss combined with historical daily water years from 2009 – 2018. Water year 2024 cumulative loss combined with historical daily water year loss exceeds the 50 percent single-year threshold for water years 2009, 2010, 2011, 2012, 2013, and 2014. Water year 2024 cumulative loss combined with historical daily water year loss does not exceed the 50 percent single-year threshold for water years 2015, 2016, 2017, and 2018.




Figure 3. Cumulative natural steelhead loss for the year (blue) and 2009 – 2018 historic cumulative loss (gray). Historic daily mean plotted in black circles.

Figure 3 is a line graph that shows cumulative natural steelhead loss for WY2024 compared with 2009-2018 historic cumulative natural steelhead loss beginning on December 1, 2023 and finishing in June 15, 2023. The line graph shows the WY2024 cumulative loss and historical daily mean begin to approach the 50% single-year threshold at the end of January 2023, falling just short of the 50% single-year threshold at the end of March, 2023. The WY2024 cumulative loss and historical daily mean begins to approach the 50% single-year threshold at the start of April, 2023 and reaches just above 500 on June 15, 2023.

## Evaluation

1. After January 1, are more than 5% of juveniles from one or more salmonid species present in the Delta?

Greater than 5% of all salmonid species are estimated to be present in the Delta.

2. Does the operational outlook’s ranges impact fish movement and change the potential distribution of fish?

Potential effects within the 7 days (near-term) in the operations outlook.

OMR flow is expected to remain at or more positive than -5,000 cfs this upcoming week. OMR flows more positive than -5,000 cfs are hypothesized to have minimal impact on movement and distribution of salmonids in the South Delta.

Potential effects longer than the 7 days (longer-term) in the operations outlook.

Not applicable, see response above to (2) (i).

3. What is the likelihood of increased loss exceeding the next annual loss threshold (50%, 75% or 90% of threshold) resulting in OMR management actions based on population distribution, abundance, and behavior of fish in the Delta?

#### Winter-run Chinook salmon

Total juvenile natural winter-run Chinook salmon (LAD) loss is 23.08 fish (as of 01/16/2024). Loss of juvenile winter-run Chinook salmon has occurred in the past week at the CVP and SWP fish salvage facilities. Final JPE calculations have been established for brood year (BY) 2023 winter-run Chinook salmon.

#### Central Valley Steelhead

Total natural juvenile steelhead loss is 28.88 fish (as of 01/16/2024). Loss of natural juvenile steelhead has occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believe that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold.

4. If an annual loss threshold has been exceeded, do continued OMR restrictions benefit fish movement and survival based on real-time information?

#### Winter-run Chinook salmon

The annual loss threshold for winter-run Chinook salmon has not been calculated in WY 2024.

#### Central Valley Steelhead

The annual loss threshold for steelhead (December 1 – March 31) has not been exceeded in WY 2024.

5. If OMR is more negative than -5,000 cfs, are there changes in spawning, rearing, foraging, sheltering, or migration behavior beyond those anticipated to occur under OMR management at -5,000 cfs?

Expected OMR flows are -4,000 to -5,100 cfs for the next week. Under OMR flows more negative than -5,000 cfs the SaMT expects impacts to rearing, foraging, sheltering, or migration of salmonids present in the south Delta. Salmonid presence in the south Delta is difficult to assess because of limited observations and there is uncertainty in how much of the population might be impacted.

## Biology Distribution and Evaluation of Green Sturgeon

### Population Status

- Delta Life Stages:
  - Adults and Juveniles

### Distribution

#### *Current Distribution*

- Adults: Most abundant during spring spawning migration period of March through May, and post spawning out-migration periods May through June; October through January depending on first winter storm event resulting in significant Sacramento River flow increases. Adult presence year-round to a lesser extent mainly in San Pablo Bay.
- Juveniles: Age-1 through Age-3 juveniles present year-round and widely distributed. Juveniles tagged with acoustic tags in the main channel Sacramento River near Sherman Island detected in the Sacramento River as far upstream as the Cache Slough complex, in the San Joaquin River at the Antioch Bridge, in Threemile, Horseshoe Bend, and Montezuma Sloughs. Seasonal abundance at the primary sampling site (near Sherman Island) appears to be highest during summer in based on capture and telemetry data. Residence time at the primary sampling site for individual fish ranges from one day to over one year but telemetry data show outmigration from the primary sampling site to the Pacific Ocean ranges from 27 to 552 days. Recent capture data shows diurnal depth preference in the main channel of the Sacramento River. No recent documentation of shallow water habitat presence or foraging.

#### *Historical Trends*

- Juvenile and adult green sturgeon are historically present in the San Joaquin and Sacramento rivers and Delta.

#### *Forecasted Distribution within Central Valley and Delta regions*

- Juvenile and adult green sturgeon are likely present in the San Joaquin and Sacramento rivers and Delta during the next week.

## Evaluation

1. Is there likely to be salvage that may exceed the annual loss limit?

Green sturgeon salvage is 0 fish (as of 1/16/2024). The agencies in the SaMT assessed the likelihood of salvage occurring in the next week is unlikely to occur.

## Biology, Distribution, and Evaluation of Delta Smelt

### Population Status

- Delta Smelt Life Stages:
  - Subadults, Adults
- Brood Year 2023:
- Abundance estimate:
  - The most recent non-zero abundance estimate for Delta Smelt is from December 8, 2023, and was 1,321 (95% CI: 183 to 4,775).
- Biological Conditions:
  - Adult and subadult Delta Smelt are expected to be present in the Low Salinity Zone and Sacramento Deep Water Shipping Channel and have been most recently detected in the lower Sacramento River. The Smelt Monitoring Team discussed the most recent monitoring data (TABLE 7) and considered published literature and professional judgement on the historical trends in regional distribution.

### Distribution

#### *Current Distribution*

- Real time detection data are currently limited to EDSM, Chipps Island Trawl and SLS. Bay Study provides data as available.
- Three adult and two juvenile Delta Smelt have been detected by surveys in the lower Sacramento River between 10/5/2023-1/16/2024. Two of these fish have been from experimental releases.
- No Delta Smelt have been detected in salvage at the SWP and CVP this water year.
- Larval sampling at the Skinner Fish Facility (SFF) and the Tracy Fish Collection Facility (TFCF) has not yet been initiated this year.
- COA 8.5.2: Spawning has not yet begun.



Table 7. Summary of newly reported detections of Delta Smelt by Region and Salvage Facilities since the last assessment. Regions are those defined by EDSM sampling. Delta Smelt >58mm FL are considered adults. Subadult fish are considered by the SMT to be fish from the previous year’s cohort based on size and timing of collection. Young of year are considered juveniles and larvae. Salvage values reflect pre-expansion salvage.

Life Stage	North	South	West	Far West	Salvage
Adult	0	0	1	0	0
Subadult	0	0	0	0	0
Larvae/Juvenile	0	0	0	0	0

Table 8. Summary of recent Delta Smelt detections reported since last assessment and the total detections for the current water year. Notes reflect latest information on reported detections or completion of survey for the water year and include both larval and adult detections. Total Fish counts do not distinguish between hatchery origin and wild Delta Smelt. Table indicates new detections and previously reported detections that have undergone preliminary ID, QA/QC, and genetic confirmation. Numbers are updated as QA/QC and genetic confirmation become available. Tracy and Skinner facility counts reflect pre-expansion salvage.

Sampling Method	Frequency	New Detections	Preliminary Detections	QA/QC Detections	Genetically Confirmed to Date	Total WY2024	Notes
EDSM	Weekly	1	1	4	N/A	5	Phase 1 began 12/4/2023
SKT	Monthly	0	N/A	N/A	N/A	0	not occurring this year
SLS	Biweekly	0	N/A	N/A	N/A	0	Began 12/11/2023
20-mm	Biweekly	0	N/A	N/A	N/A	0	Begins: 3/18/2024
Summer Towntnet	Biweekly	0	N/A	N/A	N/A	0	Begins:
Bay Study	Monthly	0	N/A	N/A	N/A	0	Ongoing
FMWT	Monthly	0	N/A	N/A	N/A	0	Complete
Chippis Island Trawl	Weekly	0	N/A	N/A	N/A	0	Ongoing

Sampling Method	Frequency	New Detections	Preliminary Detections	QA/QC Detections	Genetically Confirmed to Date	Total WY2024	Notes
FCCL Brood Stock Collections	Weekly	0	N/A	N/A	N/A	0	Begins: 11/27/2023
LEPS	As available	0	N/A	N/A	N/A	0	Began 1/2/2024
FRP	Daily	0	N/A	N/A	N/A	0	Ongoing
Tracy Fish Collection Facility (CVP)	Daily	0	N/A	N/A	N/A	0	Ongoing
Skinner Fish Facility (SWP)	Daily	0	N/A	N/A	N/A	0	Ongoing
Total	N/A	N/A	N/A	N/A	N/A	5	Sum of all Delta Smelt observed during the water year

### ***Cultured Delta Smelt Experimental Releases***

- Experimental releases completed in Water Year 2024 include:
  - Release 1: 14,104 fish released at Sacramento River at Rio Vista (truck release)
  - Release 2: 13,089 fish released at Sacramento River at Rio Vista (hard and soft releases)
  - Release 3: 12,691 fish released at Sacramento River at Rio Vista (hard carboy and trailer release)
  - Release 4: 25,649 fish released at Sacramento River at Rio Vista (truck release)
- Other experimental releases for Water Year 2024 are planned for:
  - Release 5: 1/24-1/25/2024
  - Release 6: 1/31-2/1/2024
- See additional details at: [Central Valley Prediction & Assessment of Salmon](#)

Table 9. Weekly summary of the origin of Delta Smelt. These identifications are considered tentative and additional genetic testing will confirm the identity of individuals. Individuals with no tags are provided alive to the FCCL as potential additions to the FCCL Broodstock.

Date	Survey	Stratum/Station	Total Caught	Ad. Clipped	VIE	No Tag
1/16/2024	EDSM	Lower Sacramento	1	X	N/A	N/A

### *Historical Trends*

- Upstream migration for Delta Smelt occurs between September and December and in response to “first flush” conditions (Sommer et al. 2011, Grimaldo et al. 2009). Migration typically ranges one to four weeks after flow and turbidity increases, based on salvage data (Sommer et al. 2011).
- Historically, detections of ripe Delta Smelt began in January and peaked in February and March and the majority of Delta Smelt spawning occurs within a temperature range of 9-18°C (Damon et al. 2016).
- Based on historical monitoring data from the past few years (<https://github.com/Delta-Stewardship-Council/deltafish>), first detection of larvae in the Central and South Delta has typically occurred by mid to late March. ([https://www.cbr.washington.edu/sacramento/tmp/hrtsalvage\\_1676407207\\_694.html](https://www.cbr.washington.edu/sacramento/tmp/hrtsalvage_1676407207_694.html)).
- Salvage data as presented on SacPas indicates that adult Delta Smelt salvage in recent years has reached the 50th percentile at the end of February – beginning of March.
- Historically, the highest peak in salvage is in May and the second highest is in June (Grimaldo et al 2009).

### *Forecasted Distribution within Central Valley and Delta regions*

- Predicting the distribution of Delta Smelt is currently difficult because detection data is limited to a few wild individuals and historic patterns may not be representative of the low population levels.
- The SMT uses turbidity as a surrogate for Delta Smelt presence and in making assessments of the likelihood of entrainment for larval Delta Smelt after spawning begins.
- The potential of experimentally released Delta Smelt to distribute from their release site is unknown at this time and SMT cannot predict their distribution beyond the original release site and subsequent recaptures. There is a high degree of uncertainty regarding the response of cultured fish to environmental cues typically applied to wild Delta Smelt.

## Abiotic Conditions

### *Turbidity*

- Moderate precipitation (between a quarter and half of an inch) expected on Tuesday and Wednesday, with rain likely for the remainder of the week. In Stockton, ESE winds 6-9 mph. In Antioch, E winds around 6 mph.
- Turbidity is below 12 FNU at OBI and at other stations in the central and south Delta. Turbidity may increase with expected precipitation this week.

Table 10. Relevant Environmental Factors to the current management actions for Delta Smelt.

Date Reported	FPT 3-day Running Average Flow (cfs)	FPT 3-day Running Average Turbidity (FNU)
1/15/2024	16938.81	15.59

### *X2 Conditions*

- As of 1/15/2024, X2 was estimated to be around 75 km

### *Other Environmental Conditions*

- The Fish and Water Operation Outlook OMR Index values are expected to range between -4,000 to -5,100 cfs this week.
- QWEST was +1500 cfs as of 1/15/2024 and is estimated to range between -600 to +5000 cfs this week, depending on precipitation totals. Real time tracking of environmental conditions, relevant thresholds and Delta Smelt catch data are updated daily at: [http://www.cbr.washington.edu/sacramento/workgroups/delta\\_smelt.html](http://www.cbr.washington.edu/sacramento/workgroups/delta_smelt.html).

## Evaluation

### USBR and DWR Proposed Operations:

- Monthly Delta Outflow for January greater than 6,000 cfs; E/I ratio not to exceed 0.65; OMR flow is less negative than or equal to -5,000 cfs. SWP is targeting a proportional share of combined exports that would result in an OMRI of -3500 on a 7-day average per the recommendation regarding the ITP COA 8.4.2.

1. Between December 1 and January 31, has any first flush condition been exceeded?

No first flush condition has been exceeded. The predicted amount of precipitation for this week is unlikely to create “First Flush” conditions and trigger IEWPP regulations this week. However, conditions will be monitored throughout the rest of the week.

2. Do DSM have a high risk of migration and dispersal into areas at high risk of future entrainment? (December 1- January 31)

Based on distribution patterns over the past decade and recent detection data, Delta Smelt are unlikely to be prevalent in the South Delta. No live Delta Smelt have been detected in the South Delta. A fourth experimental release of hatchery Delta Smelt occurred on 1/10/2024 at Rio Vista, which is outside of the South Delta. Information regarding their behavior post-release is limited and catch will be monitored.

3. Has a spent female been collected?

This question is not applicable until Turbidity Bridge Avoidance begins.

4. If OMR of -2000 cfs does not reduce OBI turbidity below 12NTU/FNU, what OMR target is deemed protective between -2000 and -5000 cfs?

This question is not applicable until Turbidity Bridge Avoidance begins.

5. If OBI is 12 NTU/FNU, what do other station locations show?

This question is not applicable until Turbidity Bridge Avoidance begins.

6. If OBI is 12 NTU/FNU, is a turbidity bridge avoidance action not warranted? What is the supporting information?

This question is not applicable until Turbidity Bridge Avoidance begins.

7. After March 15 and if QWEST is negative, are larval or juvenile DSM within the entrainment zone of the CVP and SWP pumps based on surveys?

This question is not applicable until after March 15.

8. Based on real-time spatial distribution of Delta Smelt and currently available turbidity information, should OMR be managed to no more negative than -3,500?

This question is not applicable until March 15.

9. What do hydrodynamic models, informed by EDSM or other relevant data, suggest the estimated percentage of larval and juvenile DSM that could be entrained may be?

This question is not applicable until March 15.

## Delta Smelt References

- Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. *California Fish and Game* 102(4):188–210.
- Hobbs, J. A., Lewis, L. S., Willmes, M., Denney, C., & Bush, E. (2019). Complex life histories discovered in a critically endangered fish. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-52273-8>
- Grimaldo, L. F., T. Sommer, N. Van Ark, G. Jones, E. Holland, P. B. Moyle, B. Herbold & P. Smith (2009) Factors Affecting Fish Entrainment into Massive Water Diversions in a Tidal Freshwater Estuary: Can Fish Losses be Managed? *North American Journal of Fisheries Management*, 29:5, 1253-1270, DOI: 10.1577/M08-062.1
- Gross, E. S. (2021). Modeling Delta Smelt Distribution for Hypothesized Swimming Behaviors. *San Francisco Estuary and Watershed Science*, 19(1).
- Kimmerer, W. J. (2008). Losses of Sacramento River Chinook Salmon and Delta Smelt to Entrainment in Water Diversions in the Sacramento-San Joaquin Delta. *San Francisco Estuary and Watershed Science*, 6(2).
- Polansky, L., Newman, K.B., Nobriga, M.L. et al. Spatiotemporal Models of an Estuarine Fish Species to Identify Patterns and Factors Impacting Their Distribution and Abundance. *Estuaries and Coasts* 41, 572–581 (2018). <https://doi.org/10.1007/s12237-017-0277-3>
- Smith, W. E., Polansky, L., and M. L Nobriga. 2021. Disentangling risks to an endangered fish: using a state-space life cycle model to separate natural mortality from anthropogenic losses. *Canadian Journal of Fisheries and Aquatic Sciences*, 78: 1008-1029.
- Sommer, T., F. Mejia, M. Nobriga, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science* 9(2).