



# Weekly Fish and Water Operations Outlook

2/27/2024 – 3/5/2024

## Water Project Operational Intent for Week

- Effective 2/7 operations will be limited to a 7-day average OMRI flow no more negative than -3,500 cfs per the 2023 IOP/ITP COA 8.5.2. until the average measured Secchi depth, as defined in the IOP and ITP, is greater than 1 meter; X2 at Chipps for 28 days in February; E/I <= 0.35.
- Proposed Action- Water project operations are currently limited to a 14-day average OMRI of no more negative than -2,500 cfs based on the 75% loss threshold for Steelhead being exceeded.
- Reclamation will be in close coordination with other agencies to resolve the operational plan in response to the 100% exceedance for Steelhead

## Forecasted Weather

- A weak system today brings some showers and snow into the Central Valley. A more active pattern begins late in the week through this weekend. Precipitation, snow, gusty winds and abnormally cold temperatures near freezing with morning frost are forecasted to return with this system.

## Tables

Table 1: Anticipated weekly operational ranges by tributary. Environmental and fish conditions updated by respective watershed groups at varying intervals that may not coincide with the weekly range of Water Operations

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Clear Creek	<ul style="list-style-type: none"> <li>• Current Release: 200 cfs</li> <li>• Anticipated Weekly Range of Releases: 200 – 700 cfs</li> </ul>	<ul style="list-style-type: none"> <li>• Spring-run Chinook salmon fry rearing.</li> <li>• Fall-run Chinook salmon eggs are incubating and hatching. Juveniles are emerging and rearing.</li> <li>• Late fall-run Chinook Salmon adults are spawning and eggs are incubating.</li> <li>• Adult O. mykiss are migrating and spawning. Their eggs are incubating and juveniles are rearing.</li> </ul> <p><i>(Updated 2/5/2024)</i></p>

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Sacramento River	<ul style="list-style-type: none"> <li>• Shasta Storage: 3.897 MAF</li> <li>• Current Release: 35,000 cfs</li> <li>• Anticipated Weekly Range of Releases: 20,000 cfs to 35,000 cfs for flood management.</li> </ul>	<ul style="list-style-type: none"> <li>• Late fall-run Chinook adults are continuing their spawning but we are past the peak period.</li> <li>• Late-fall run Chinook eggs/alevins are currently incubating/residing in the gravel.</li> <li>• Fall-run redds are mostly emerged. Fry are very actively migrating downstream.</li> <li>• Adult winter-run are arriving in the upper river and holding.</li> </ul> <p><i>(Updated 2/12/2024)</i></p>
Feather River	<ul style="list-style-type: none"> <li>• Oroville Storage: 2.973 MAF</li> <li>• Current Release: 12,000 cfs</li> <li>• Anticipated Weekly Range of Releases: 8,000 cfs to 20,000 cfs for flood management.</li> </ul>	<ul style="list-style-type: none"> <li>• Fall-run Chinook salmon adult spawning has ended. Eggs are incubating in gravel. Fry are beginning to emerge and migrate downstream.</li> <li>• O. mykiss juveniles are rearing. Adults are spawning and eggs are incubating in gravel.</li> <li>• Adult green sturgeon are still holding in the LFC near Fish Barrier Dam.</li> <li>• Spring-run Chinook salmon adult spawning has ended. Fry are emerging and migrating downstream.</li> </ul> <p><i>(Updated 2/27/2024)</i></p>
American River	<ul style="list-style-type: none"> <li>• Folsom Storage: 617 TAF</li> <li>• Current Release: 6,000 cfs</li> <li>• Anticipated Weekly Range of Releases: 6,000 cfs to 8,000 cfs for flood management</li> </ul>	<ul style="list-style-type: none"> <li>• O. mykiss juveniles are rearing.</li> <li>• Adult fall-run Chinook salmon spawning has ended.</li> <li>• Eggs are incubating in gravel.</li> <li>• Fry are beginning to emerge and migrate downstream.</li> <li>• O. mykiss – Adults present</li> </ul> <p><i>(Updated 1/30/2024)</i></p>
Stanislaus River	<ul style="list-style-type: none"> <li>• New Melones Storage: 1.981 MAF</li> <li>• Current Release: 1,000 cfs</li> <li>• Anticipated Weekly Range of Releases: 1,000 – 3,000 cfs for flood management</li> </ul>	<ul style="list-style-type: none"> <li>• O. mykiss - Adult and juveniles present</li> <li>• Fall-run Chinook salmon spawning has ended.</li> <li>• Eggs are incubating in gravel. Fry are beginning to emerge and migrate downstream.</li> </ul> <p><i>(Updated 1/29/2024)</i></p>

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Delta	<ul style="list-style-type: none"> <li>• Freeport: 50,000 to 70,000 cfs</li> <li>• Vernalis: 6,000 to 10,000 cfs</li> <li>• Delta Outflow index: 70,000 to 150,000 cfs</li> <li>• Combined Exports: 4,600 to 7,700 cfs</li> <li>• JPP: Current 4,200 cfs, Range 3,600 cfs to 4,200 cfs</li> <li>• CCF: Current 2,400 cfs, Range 1,000 cfs to 3,500 cfs</li> <li>• Expected Daily OMR Index Values: -2,000 cfs to -2,500 cfs</li> <li>• DCC Gates: Closed on 11/27 for season</li> <li>• X2 &lt; 56 km</li> <li>• Tides: Transitioning from Spring to Neap; Last Quarter Moon on 3/3</li> </ul>	<ul style="list-style-type: none"> <li>• Juvenile and adult <i>O. mykiss</i> present.</li> <li>• Juvenile Chinook Salmon present.</li> <li>• Adult winter-run Chinook Salmon are present.</li> <li>• Adult and juvenile Green Sturgeon present</li> <li>• Delta Smelt adults have been detected in the lower Sacramento River, Suisun Marsh, Western Delta, lower San Joaquin River, Liberty Island, and the South Delta since 2/6. DS population scale migration is likely completed, and water temperatures are suitable for spawning.</li> <li>• Longfin Smelt sub-adults and adults have been detected in Suisun Marsh and Bay, Honker and Grizzly bays, San Pablo Bay, Chipps Island, the Confluence, and the Lower Sacramento River. Longfin Smelt larvae have been detected in the Napa River, Carquinez Strait, San Pablo Bay, Suisun Bay and Marsh, the Confluence, lower Sacramento and San Joaquin rivers, and the Central and South Delta. LFS population scale migration and spawning are ongoing.</li> </ul> <p style="text-align: right;"><i>(Updated 2/27/2024)</i></p>

Table 2a-b: WY 2024 relevant Fish and Environmental Criteria and Status in 2019 Reclamation LTO Action Cumulative loss for the duration of 2019 Biological Opinion began upon signature of ROD, 2/19/2020.

Table 2a: WY 2024 Salmonid Current Loss and Delta Smelt Abiotic Conditions. Additional Real-Time OMR Restrictions and Performance Objectives (4.10.5.10.2, 4.10.5.10.3) and Onset of OMR Management (4.10.5.10.1). Genetic identification of salmon is not used in calculating loss, but results are included in the Assessment as they become available. The Final WR JPE for BY 2023 is 234,896. The ITL and performance thresholds are TBD.

Species/run	Threshold	Current Status	Weekly Trend	Updated
Green sturgeon	WY 2024 salvage = 74	WY 2024 salvage = 0 (0%)	No change expected	2/26/2024

Species/run	Threshold	Current Status	Weekly Trend	Updated
Natural winter-run Chinook Salmon	WY 2024 loss = 2061 (75% of 1.17% of JPE)	WY 2024 loss = 1443.47 (70.03%) 50% threshold exceeded 2/25/2024	Increasing	2/26/2024
Natural Steelhead	Dec 1 – Mar 31 = 1414; (100% of 1,414) Apr 1 – June 15 = 776 (50% of 1,552)	WY 2024 loss = 1,778 Dec 1 – Mar 31 = 1,778 75% threshold exceeded 2/22/24 and 100% threshold loss exceeded on 2/23/24 50% threshold exceeded 2/11/2024 Apr 1 – June 15 = 0 (0% of the 50% threshold)	Increasing	2/27/2024
Sacramento River Hatchery winter-run Chinook salmon	WY 2024 loss = 140.93 (50% of 0.12% of JPE)	WY 2024 loss = 0 (0%)	Increasing	2/27/2024
Battle Creek Hatchery winter-run Chinook salmon	WY 2024 loss = 234.90 (1% of JPE)	WY 2024 loss = 0 (0%)	No change expected	2/27/2024
Proposed Action Hatchery yearling spring-run Chinook salmon surrogates	> 0.5% of each release group 1) 12/22/2023 group 1: 60,764 = 303.82 2) 12/29/2023 group 2: 71,049 = 355.25 3) 1/11/2024 group 3: 67,018 = 335.09	WY 2024 loss = 1) 36.84 (12.12%) 2) 31.75 (8.94%) 3) 75.69 (22.6%)	Increasing	2/27/2024
Delta Smelt	After Dec. 1: Running 3-day avg. flows at Freeport >25,000 cfs AND Running 3-day avg. turbidity at Freeport =>50 FNU	Offramped. Implemented 1/23/2024-2/5/2024.  Flow = N/A; Turbidity = N/A	Not relevant	2/12/2024

Species/run	Threshold	Current Status	Weekly Trend	Updated
Delta Smelt	Daily avg. Turbidity at OBI=>12 FNU	In effect. Not triggered.  OBI Daily Average = 6.0 FNU	Remaining stable or increasing	2/20/2024
Delta Smelt	Daily avg. Temperature at CCF > 25°C for three consecutive days	Not in effect.  CCF (CLC CDEC station) daily avg. Temperature = N/A	Not relevant	2/12/2024

Table 2b. 10-Year Salmonid Cumulative Loss

Species/run	Threshold	Current Status	Updated
Natural winter-run Chinook salmon	Loss = 8,738	Cumulative loss = 1813.72 (20.76%)	2/26/2024
Hatchery winter-run Chinook salmon	Loss = 5,356	Cumulative loss = 6.71 (0.13%)	2/26/2024
Natural steelhead	Loss = 6,038 (Dec 1 – Mar 31) Loss = 5,826 (Apr 1 – June 15)	Cumulative loss = 3355.11 (55.6%, Dec 1 – Mar 31) 1012.50 (17.4%, Apr 1 – June 15)	2/26/2024

Table 3a-c: Relevant Water Year 2024 Fish Criteria and Status for Listed Fish under the SWP Long-Term Incidental Take Permit.

Table 3a: Chinook Salmon

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
OMR Mgmt. triggered (8.3.2)	Jan. 1 - Jun. 30 (when ≥ 5% of spring-run or winter-run in Delta)	In effect	≥ 5% of the Winter-run or Spring-run population in Delta	N/A	N/A	2/26/2024	In effect as of January 1.

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Winter-run yearly loss (8.6.1)	Nov. 1 - Jun. 30	In effect	2,748.28 (1.17% of Natural LAD WR of Final JPE) 232.30 (0.12% of LSNFH WR of Final JPE)	Loss of total LAD Winter-run = 1374.27 (50.00% of Natural LAD WR threshold)	Salvage is likely to occur in the upcoming week	2/26/24	Natural-origin LAD winter-run Chinook salmon (WR) were observed in salvage the previous week.
Winter-run discrete daily loss (8.6.2)	Nov. 1 - Dec. 31	Not in effect	26 older juvenile/day	Max Older Juvenile discrete daily loss observed last week = N/A	N/A	1/22/2024	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Mid and late season Winter-run daily loss threshold (8.6.3)	Jan 1 – May 31	In effect	2/1-2/29: loss of 5.43 fish /day of natural-origin LAD older juvenile (Winter-run, Yearling fall-run, and Late fall-run). Updated with genetic results as they become available. If genetics confirms that the older juvenile is NOT a WR then that fish will not count towards the threshold. Upcoming: 3/1-3/31: loss of 8.74 fish /day of natural-origin LAD older juvenile	Max daily Loss of older juvenile Chinook occurred on 2/23 with loss of 316.59.	Salvage of older juveniles is likely in the upcoming week.	2/26/2024	4 genetically confirmed WR were observed in salvage for the season.
Spring-run surrogate protection (8.6.4)	Feb. 1 - Jun. 30	Not in effect	TBD	N/A	N/A	9/29/23	Will be updated when in effect

Table 3b: Delta Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Integrated Early Winter Pulse Protection ('First Flush') (8.3.1)	Dec. 1 - Jan. 31	Off-ramped	- three-day Freeport daily flow running avg $\geq$ 25,000 AND  [three-day Freeport turbidity running avg $\geq$ 50 FNU OR Smelt Monitoring Team recommendation]	Not relevant	Not relevant	2/12/24	N/A
Turbidity Bridge Avoidance (8.5.1)	Dec. 15 - Apr. 1	In effect; not triggered	Occurs after the Integrated Early Winter Pulse protection or February 1 whichever comes first until April 1 -avg. OBI turbidity $>$ 12 FNU	OBI daily average = 5.4 FNU	Remaining stable or increasing	2/25/24	N/A



Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Larval and/Juvenile Delta smelt Protection (8.5.2)	Nov. 1 – Jun. 30	Triggered 2/5/2024, also 2/21/2024	- If 5-day cum. salvage of juv.DS >= 1 [average 3-yr FMWT index + 1], then –5000 OMR - If DS in SLS/20mm or 3-d temp at Jersey Point >= 12C, and SLS/20mm Secchi for 12 south delta stations <= 1m, then – 3500 OMR	Current 5-day salvage = 0 3-day SJJ temp = 12.5 °C Average Secchi Depth = 83cm (as of 2/21)	N/A	2/25/24	N/A

Table 3c: Longfin Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Early Adult Protection (8.3.3)	Dec. 1 - Feb. 28	Off-ramped	-Cum. salvage > [most recent FMWT/10] = 46 fish (Sept.-Dec. Index) OR -Smelt Monitoring Team determines high likelihood of LFS movement into high-risk areas	Cumulative salvage = 0	N/A	12/26/23	N/A

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
OMR Mgt. for Adults (8.4.1)	Onset of OMR mgmt - Feb. 28	Off-ramped	-Smelt Monitoring Team recommendation	N/A	N/A	12/19/23	N/A
Larval and Juvenile Longfin Smelt Entrainment Protection (8.4.2)	Jan 1 – Jun 30	Not triggered by SLS 4	-LFS larvae or juveniles in >=4 SLS or 20 mm stations in central and south Delta, OR -LFS catch/tow >5 larvae or juveniles in >=2stations	SLS 4 detected 2 larvae at 809 on 2/20/24.	N/A	2/13/24	N/A
High Flow OMR Off-Ramp for Longfin Smelt (8.4.3)	Based on the status of 8.3.3, 8.4.1, & 8.4.2	Triggered 2/6/2024	-Sac. R. at Rio Vista >55,000, OR SJR at Vernalis >8,000	Rio Vista = 70,000 – 130,000 – cfs SJ = 6,000 – 10,000 cfs	Flows are decreasing	2/26/24	Off-ramps 8.4.2 until Sac flows at Rio Vista <40,000 cfs

Table 4: Fish monitoring gear efficiency and disruptions. Status Categories: [1] Active (ongoing sampling), [2] Partial Interruption (some sampling interruptions), [3] Interrupted (sampling fully suspended), [4] Not Active (sampling not scheduled)

Monitoring survey	Region	Notes (as of 2/26/2024)	Status
SWP regular counts, CWT reading	Delta	Active	1
SWP larval sampling	Delta	Not Active (Will begin 3/11)	4
CVP regular counts, CWT reading	Delta	Active	1
CVP larval sampling	Delta	Active	1
Smelt Larval Survey	Delta	Active	1
LEPS	Delta	Active	1
20mm Survey	Delta	Not Active	4
Fall Mid-water Trawl	Delta	Not Active	4
Summer Townet Survey	Delta	Not Active	4

<b>Monitoring survey</b>	<b>Region</b>	<b>Notes (as of 2/26/2024)</b>	<b>Status</b>
Bay Study	Delta	Active	1
DJFMP- Chipps and Sacramento Trawls	Delta	Active	1
DJFMP- Seines	Delta	Active	1
EDSM	Delta	Active	1
Environmental Monitoring Program (EMP)	Delta	Active	1
Mossdale Trawl	Delta	Active	1
USGS Flow monitoring	Delta	Active	1
Red Bluff Diversion Dam Rotary Screw Trap (RST)	Sacramento River	Active	1
Knights Landing RST	Sacramento River	Active	1
Tisdale RST	Sacramento River	Active	1
Yuba River (Hallwood) RST	Yuba River	Active	1
Redd dewatering and stranding surveys	Sacramento River	Active	1
Sacramento Carcass and Redd Surveys (Late fall-run Chinook salmon)	Sacramento River	Active	1
Lower Sacramento RST	Sacramento River	Active	1
Feather River (upper DWR) RST	Feather River	Active	1
Feather River (lower CDFW) RST	Feather River	Active	1
Lower American River at Watt Ave RST	American River	Not Active (as of 2/18 due to heavy debris)	4
SJRRP CDFW Field Monitoring	San Joaquin River	Active	1
SJRRP USBR Field Monitoring	San Joaquin River	Active	1
Stanislaus Fish Weir	Stanislaus River	Active	1
American River Carcass/Redd Surveys (Fall-run Chinook salmon)	American River	Not Active	4
Stanislaus Carcass Survey (Steelhead)	Stanislaus River	Active	1
Caswell RST	Stanislaus River	Active	1
Wallace Weir	Cache Slough	Active	1
Butte Creek RST/Diversion Trap	Butte Creek	Active	1