

Weekly Fish and Water Operations Outlook 10/11/2022 – 10/17/2022

FORECASTED WEATHER: Warm and dry, but cooler than last week.

Table 1: Anticipated weekly operational ranges by tributary. Environmental and fish conditions updated by respective watershed groups at varying intervals that may not coincide with the weekly range of Water Operations

cical cicci	Current Release: 200 cfsAnticipated Weekly Range of	Spring-run Chinook salmon spawning is
	Releases: 200 cfs	 winding down. Fall-run Chinook Salmon are entering the Creek and spawning is ramping up. (Updated 10/3/22)
	 Shasta Storage: 1.484 MAF Current Release: 4,100 cfs Anticipated Weekly Range of Releases: 4,100 cfs to 3,500 cfs 	 Winter-run Chinook salmon are the dominant run passing at RBDD currently in relatively low and variable numbers. All other species either absent or caught in sporadic and low numbers. Spring-run Chinook salmon adults present and spawning is underway. Spawning surveys began in early September for adult spring-run Chinook salmon. Winter-run Chinook juvenile salmon are present in upper river and are both emerging from redds and migrating downstream. Fall-run Chinook salmon are beginning to spawn. Carcass surveys for fall-run are underway. (Updated 10/3/22)

Tributary/Division	Anticipated Weekly Ranges	Related Environmental and Fish Conditions
Feather River	 Oroville Storage: 1.202 MAF Current Release: 2,500 cfs Anticipated Weekly Range of Releases: 2,500 cfs to 2,300 cfs Daily temperature maximum: 51 F at Fish Hatchery 	 Adult Fall-run Chinook Salmon are just beginning to spawn, some still migrating upstream and some holding Adult Spring-run Chinook Salmon are spawning, eggs in gravel. Approximately 15-20 green sturgeon present form Sunset Pumps to Boyd's Pump. Adult and juvenile steelhead present. (Updated 10/11/22)
American River	 Folsom Storage: 333 TAF Current Release: 1,500 cfs Anticipated Weekly Range of Releases: 1500 cfs to 1,300 cfs 	 Adult fall-run Chinook Salmon are present and holding. Juvenile and adult O. mykiss are present.
Stanislaus River	 New Melones Storage: 614 TAF Current Release: 200 cfs Anticipated Range of Weekly Releases: 200 cfs to 1,300 cfs. (Fall pulse begins 10/12) 	 Juvenile and adult O. mykiss are present. Adult fall-run Chinook salmon are present and holding. (Updated 10/4/22)
Delta	 Freeport: 6,500 to 7,500 cfs Vernalis: 300 to 1,000 cfs Delta Outflow index: 3,500 to 5,000 cfs Combined Exports: 1,200 to 2,300 cfs JPP: 900 cfs to 1,800 cfs CCF: 300 cfs to 500 cfs Expected Daily OMR Index Values: - 1,200 to -2,000 cfs DCC Gates: Closed as of 10/10. Expected operations: Closed during the week and open on weekends, but may need flexibility to change if necessary. 	 Adult spring-run Chinook Salmon entering Delta and migrating upstream towards spawning grounds Adult O. mykiss present Adult and juvenile Green Sturgeon present Delta Smelt sub-adults are present in the north delta arc and downstream of the confluence. Longfin Smelt sub-adults and adults are downstream of the confluence. (Updated 10/11/22)

Table 2a-b: WY 2023 relevant Fish and Environmental Criteria and Status in 2019 Reclamation LTO Action Cumulative loss for the duration of 2019 Biological Opinion began upon signature of ROD, 2/19/2020.

Table 2a: WY 2023 Salmonid Current Loss and Delta Smelt Abiotic Conditions. Additional Real-Time OMR Restrictions and Performance Objectives (4.10.5.10.2) and Onset of OMR Management (4.10.5.10.1). * TBD – no draft JPE produced, ITL and performance thresholds are TBD currently

Species/run	Threshold	Current Status	Weekly Trend	Updated
Green sturgeon	WY 2023 salvage = 74	WY 2023 salvage = 0 (0%)	No change expected	10/11/2022
Natural winter-run Chinook Salmon	WY 2023 loss = TBD * (50% of 1.17% of JPE)	WY 2023 loss = 0	No change expected	10/11/2022

Species/run	Threshold	Current Status	Weekly Trend	Updated
Natural Steelhead	Dec 1 – Mar 31 = 707 (50% of 1,414) Apr 1 – June 15 = 776 (50% of 1,552)	WY 2023 loss = 0 Dec 1 – Mar 31 = 0 (0%) Apr 1 – June 15 = 0 (0%)	No change expected	10/11/2022
Sacramento River Hatchery winter-run Chinook salmon	WY 2023 loss = TBD* (50% of 0.12% of JPE)	WY 2023 loss = 0 (0%)	No change expected	10/11/2022
Battle Creek Hatchery winter-run Chinook salmon	WY 2023 loss = TBD * (50% of 0.12% of JPE)	WY 2023 loss = 0 (0%)	No change expected	10/11/2022
Proposed Action Hatchery yearling spring-run Chinook salmon surrogates	> 0.5% of each release group	WY 2023 loss = 0 (0%)	No change expected	10/11/2022
Delta Smelt	Daily avg. Turbidity at OBI=>12 FNU	OBI daily Avg Turbidity = Not relevant	Not relevant	10/11/2022
Delta Smelt	Daily avg. Temperature at CCF > 25°C for three consecutive days	CCF daily avg. Temperature = Not relevant	Not relevant	10/11/2022

Table 2b. 10-Year Salmonid Cumulative Loss

Species/run	Threshold	Current Status	Updated
Natural winter-run Chinook salmon	Loss = 8,738	Cumulative loss = 264.2 (3.0%)	10/11/2022
Hatchery winter-run Chinook salmon	Loss = 5,356	Cumulative loss = 6.71 (0.13%)	10/11/2022
Natural steelhead	Loss = 6,038 (Dec 1 – Mar 31) Loss = 5,826 (Apr 1 – June 15)	Cumulative loss = 530.2 (8.8%, Dec 1 – Mar 31) 474.5 (8.1%, Apr 1 – June 15)	10/11/2022

Table 3a-d: Relevant Water Year 2023 Fish Criteria and Status for Listed Fish under the SWP Long-Term Incidental Take Permit.

Table 3a: Chinook Salmon

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
OMR Mgmt. triggered (8.3.2)	Jan. 1 - Jun. 30 (when ≥ 5% of spring-run or winter- run in Delta)	Not in effect	-5% of the Winter-run or Spring-run population in Delta	N/A	N/A	10/10/22	N/A
Winter-run yearly loss (8.6.1)	Nov. 1 - Jun. 30	Not in effect (based on JPE)	TBD (based on JPE)	N/A	N/A	10/10/22	N/A
Winter-run discrete daily loss (8.6.2)	Nov. 1 - Dec. 31	Not in effect	11/1-11/30: loss of 6/day unclipped older juv. Winter-run 12/1-12/31: loss of 26/day unclipped older juv. Winter-run	N/A	N/A	10/10/22	N/A
Mid and late season Winter-run daily loss threshold (8.6.3)	Jan 1 – May 31	Not in effect	May 1-May 31 (0.0077% of JPE) = 9.63	N/A	N/A	10/10/22	N/A
Spring-run surrogate protection (8.6.4)	Feb. 1 - Jun. 30	Not in effect	TBD (based on the number of fish released)	N/A	N/A	10/10/22	N/A

Table 3b: Delta Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Integrated Early Winter Pulse Protection ('First Flush') (8.3.1)	Dec. 1 - Jan. 31	Dec 17, last	 three-day Freeport daily flow running avg> = 25,000 AND [three-day Freeport turbidity running avg > = 50 NTU OR Smelt Monitoring Team recommendation] 	N/A	N/A	10/3/22	N/A
Turbidity Bridge Avoidance (8.5.1)	Dec. 15 - Apr. 1	Offramped	Occurs after the Integrated Early Winter Pulse protection or February 1 (whichever until April 1) comes first -avg. OBI turbidity>12 NTU	N/A	N/A	10/3/22	N/A
Larval and/Juvenile Delta smelt Protection (8.5.2)	ongoing	In effect, not triggered	 - 5-day cum. salvage of juv.DS > = 1[average 3- yrFMWT index + 1] OR, - 3-day cum. Salvage of juv. DS >11 	N/A	N/A	10/3/22	N/A

Table 3c: Longfin Smelt

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
Early Adult Protection (8.3.3)	Dec. 1 - Feb. 28	Offramped	-Cum. salvage > [most recent FMWT/10] =1 fish (SeptOct. Index) OR -Smelt Monitoring Team determines high likelihood of LFS movement into high-risk areas	N/A	N/A	10/3/22	N/A
OMR Mgt. for Adults (8.4.1)	Dec. 1 -Feb. 28	Offramped	-Smelt Monitoring Team recommendation	N/A	N/A	10/3/22	N/A
Larval and Juvenile Longfin Smelt Entrainment Protection (8.4.2)	Jan 1 – Jun 30	In effect, not triggered	-LFS larvae or juveniles in >=4 SLS or 20 mm stations in central and south Delta, OR -LFS catch/tow >5 larvae or juveniles in >=2stations	N/A	N/A	10/3/22	N/A
High Flow OMR Off- Ramp for Longfin Smelt (8.4.3)	Based on the status of 8.3.3, 8.4.1, & 8.4.2	In effect, not triggered	-Sac. R. at Rio Vista>55,000, OR SJR at Vernalis >8,000	Rio Vista = 2,300 to 5,000 cfs SJ = 300 to 1,000 cfs	N/A	10/10/22	N/A

Table 3d: OMR

Action	Timeframe	Current Action Status	Threshold(s)	Current Relevant Data	Weekly Trend	Last Updated	Comments
OMR Mgmt. Offramp (8.8)	Jun. 1 – Jun. 30	Not in effect	 >95% of the Winter-run and Spring run populations have migrated past Chipps Island AND -Current daily average water temperature at Mossdale and Prisoners Point. Days exceeded: Criteria met as of 6/16/2022 	N/A	N/A	10/10/22	N/A

Table 4: Fish monitoring gear efficiency and disruptions. Status Categories: [1] Active (ongoing sampling), [2] Partial Interruption (some sampling interruptions), [3] Interrupted (sampling fully suspended), [4] Not Active (sampling not scheduled)

Monitoring survey	Region	Notes (as of 10/11/2022)	Status
SWP regular counts, CWT reading	Delta	Active	1
SWP larval sampling	Delta	Not Active	4
CVP regular counts, CWT reading	Delta	Active	1
CVP larval sampling	Delta	Not Active	4
Smelt Larval Survey	Delta	Not Active	4
LEPS	Delta	Not Active	4
20mm Survey	Delta	Not Active	4
Spring Kodiak Trawl	Delta	Not Active	4
Fall Mid-water Trawl	Delta	Active	1
Summer Townet Survey	Delta	Not Active	4
Bay Study	Delta	Active	1
DJFMP- Chipps and Sacramento Trawls	Delta	Active (sampling four days a week starting in February)	1
DJFMP- Seines	Delta	Active	1
EDSM	Delta	Active	1
EMP	Delta	Active	1
Mossdale	Delta	Active	1
USGS Flow monitoring	Delta	Active	1
Red Bluff Diversion Dam screw trap	Sacramento River	Active	1

Monitoring survey	Region	Notes (as of 10/11/2022)	Status
Knights Landing screw trap	Sacramento River	Not Active	4
Tisdale screw trap	Sacramento River	Active	1
GCID screw trap	Sacramento River	Not Active	4
Yuba River (Hallwood) screw trap	Yuba River	Not Active	4
Redd dewatering and stranding surveys	Sacramento River	Not Active	4
Sacramento Carcass and Redd Surveys	Sacramento River	Active	1
Lower Sacramento Rotary Screw Trap	Sacramento River	Active	1
Feather River (upper DWR)	Sacramento River	Not Active	4
Feather River (lower CDFW)	Sacramento River	Not Active	4
SJRRP CDFW Field Monitoring	San Joaquin River	Active	1
SJRRP USFWS and USBR Field Monitoring	San Joaquin River	Not Active	4
Stanislaus Fish Weir	San Joaquin River	Active	1