Weekly Fish and Water Operations Outlook 2/15/2022 – 2/21/2022

Quick moving system on Monday brings showers in northern mountains Monday afternoon and drops temperatures down about 10-20 degrees from Sunday by Tuesday. Locally breezy conditions on Tuesday and Wednesday may elevate some fire concerns. Gradual rebound of temperatures with dry conditions for remainder of week. Change in weather pattern for the start of the extended forecast is possible with a few light showers in the mountains late Sunday into Monday.

| Tributary/ Division | Anticipated Weekly Ranges | Related Environmental and Fish Conditions* |
|---------------------|---|---|
| Clear Creek | Current Release: 200 cfs Anticipated Weekly Range of Releases: 200 cfs | Spring-run Chinook Salmon juveniles are rearing in Clear Creek. Fall-run Chinook salmon continue their emergence and many fry are rearing in Clear Creek. O. mykiss/steelhead adults are present, migrating, and spawning. Their spawning will continue through March. A few fry are starting to emerge. Late-fall run Chinook Salmon are spawning and it will continue through February. (updated 2/14/22) |

| Tributary/ Division | Anticipated Weekly Ranges | Related Environmental and Fish Conditions* |
|---------------------|---|--|
| Sacramento River | Shasta Storage: 1.660 MAF Current Release: 3,250 cfs Anticipated Weekly Range of Releases: 3,250 cfs | Winter-run Chinook Salmon juveniles present in very low numbers above RBDD and distributed throughout the Sacramento River and to the Delta. Hatchery releases of juvenile winter-run Chinook salmon from Livingston Stone Hatchery are in progress, some of these fish have been released on 2/9/2022 at boat ramp in Redding area. Other releases are going to occur in near future as fish grow to release size. Winter-run Chinook salmon adults present in increasing numbers and are holding from RBDD to Keswick Dam. Spring-run Chinook Salmon fry present and migrating downstream. Spring-run juvenile passage has slowed and fall-run Chinook salmon are the dominant run passing at RBDD currently. Some fall-run Chinook Salmon alevin remain in gravel, fry emergence and passage at RBDD is ongoing daily. Late fall-run Chinook Salmon spawning occurring, eggs and alevins in the gravel. Green Sturgeon adults and juveniles present in lower river with Delta entry likely. Adult steelhead present; juveniles absent from upper river catch. (updated 2/14/22) |
| Feather River | Oroville Storage: 1.650 MAF Current Release: 3,500 cfs Anticipated Weekly Range of Releases: 3,000 to 4,000 cfs Daily temperature targets: 55°F at Fish Hatchery | Fall-run Chinook Salmon fry are emerging and emigrating Adult steelhead are present and their spawning has commenced and will continue into April. Spring-run Chinook Salmon fry are present and emigrating (updated 2/1/22) |

| Tributary/ Division | Anticipated Weekly Ranges | Related Environmental and Fish Conditions* |
|---------------------|--|---|
| American River | Folsom Storage: 531 TAF Current Release: 2,000 cfs Anticipated Weekly Range of Releases: 1,750 to 2,000 cfs | Fall-run Chinook Salmon have finished spawning. Egg are in gravel. Fry are emerging and present in RST samples. Juvenile and adult O. mykiss are present and holding. Spawning has begun, and eggs are in gravel. (updated 1/24/22) |
| Stanislaus River | New Melones Storage: 984 TAF Current Release: 900 cfs Anticipated Range of Weekly Releases: 600 - 900 cfs for D- 1641 flow requirement at Vernalis | Juvenile and adult O. mykiss are present. O. mykiss are spawning. Fall-run Chinook Salmon eggs are in gravel and fry are emerging. (updated 2/14/22) |

| Tributary/ Division | Anticipated Weekly Ranges | Related Environmental and Fish Conditions* |
|---------------------|--|--|
| Delta | Freeport: 12,000 to 13,500 cfs Vernalis: 900 to 1,400 cfs Delta Outflow index: 11,200 to 12,000 cfs Combined Exports: 800 to 2,800 cfs JPP: 800 cfs to 1,800 cfs CCF: 0 cfs to 1,000 cfs Expected OMR Index Values: -500 to -2,000 cfs DCC Gates: Closed as of 11/30/2021 | Juvenile winter-run Chinook Salmon 10-25% yet to enter Delta, 74-85% in Delta, 1-5% exited Delta past Chipps Island YOY spring-run Chinook Salmon: 22-41% yet to enter Delta, 59-78% in Delta, 0% exited past Chipps Island Juvenile Steelhead: 43-62% yet to enter Delta, 28-42% in Delta, 10-15% exited Delta past Chipps Island Adult fall-run Chinook Salmon entering Delta and migrating upstream towards spawning grounds Adult steelhead present Adult and juvenile Green Sturgeon present Adult Delta Smelt expected to be present in the Sacramento Deep Water Ship Channel (DWSC), lower Sacramento River, Suisun Marsh, South Delta and lower San Joaquin River. Delta Smelt with a migratory life history are expected to have moved upstream and water temperatures are within the range for spawning. Longfin Smelt >60mm have been detected at Chipps, lower Sacramento River, Suisun Bay/Marsh, San Pablo, and Central Bay. Longfin Smelt larvae have been detected in the confluence, lower Sacramento River, station 716, the central and south delta, and lower San Joaquin River. (updated 2/15/22) |

* Environmental and fish conditions updated by respective watershed groups at varying intervals that may not coincide with the weekly range of Water Operations

Table 2a-b: WY 2022 relevant Fish and Environmental Criteria and Status in 2019 Reclamation LTO Action Cumulative loss for the duration of 2019 Biological Opinion began upon signature of ROD, 2/19/2020.

Table 2a: WY 2022 Salmonid Current Loss and Delta Smelt Abiotic Conditions. Additional Real-Time OMR Restrictions and Performance Objectives (4.10.5.10.2) and Onset of OMR Management (4.10.5.10.1).

| Species/run | Threshold | Current Status | Weekly Trend | Updated |
|---|---|---|------------------------------|-----------|
| Green sturgeon | WY 2022 salvage = 74 | WY 2022 salvage = 0 (0%) | No change expected | 2/13/2022 |
| Natural winter-run Chinook Salmon | WY 2022 loss = 731.47 (50% of 1.17% of JPE) | WY 2022 loss = 19.99 (2.73%) | Possible additional salvage | 2/13/2022 |
| Natural Steelhead * | Dec 1 – Mar 31 = 707 (50% of 1,414) Apr 1 – June 15 = 776 (50% of 1,552) | Dec 1 – Mar 31 = 59.80 (8.5%) Apr 1 – June 15 = 0 (0%) | Possible additional salvage | 2/13/2022 |
| Sacramento River Hatchery winter-run Chinook salmon | WY 2022 loss = 90.93 (50% of 0.12% of JPE) | WY 2022 loss = 0 (0%) | Unlikely salvage | 2/13/2022 |
| Hatchery yearling spring- run Chinook salmon surrogates | > 0.5% of each release group: 1) 12/15/2021: 84,343 = 421.7 2) 12/22/2021: 82,626 = 413.1 3) 1/6/2022: 77,325 = 386.6 | 1) 48.18 (11.4%) 2) 17.61 (4.3%) 3) 16.16 (4.2%) | Likely additional salvage | 2/13/2022 |
| Delta Smelt | Daily avg. Turbidity at OBI=>12 FNU | OBI daily Avg Turbidity = 3.83 | No Change Expected | 2/15/2022 |

* Steelhead observed on 10/30/2021 and 11/19/2021 are included in the December 1 through March 31 period

Table 2b. 10-Year Salmonid Cumulative Loss

| Species/run | Threshold | Current Status | Updated | |
|-----------------------------------|--------------|----------------------------------|-----------|--|
| Natural winter-run Chinook salmon | Loss = 8,738 | Cumulative loss = 224.88 (2.57%) | 2/13/2022 | |

| Species/run | Threshold | Current Status | Updated |
|------------------------------------|---|--|-----------|
| Hatchery winter-run Chinook salmon | Loss = 5,356 | Cumulative loss = 0 (0%) | 2/13/2022 |
| Natural steelhead | Loss = 6,038 (Dec 1 – Mar 31) Loss = 5,826 (Apr 1 – June 15) | Cumulative loss = 503.04 (8.33%, Dec 1 – Mar 31) 374.8 (6.4%, Apr 1 – June 15) | 2/13/2022 |

Table 3a-d: Relevant Water Year 2022 Fish Criteria and Status for Listed Fish under the SWP Long-Term Incidental Take Permit.

Table 3a: Chinook Salmon

| Action | Timeframe | Current Action Status | Threshold(s) | Current Relevant Data | Weekly Trend | Last Updated | Comments |
|--------------------------------------|--|--|--|---|---|--------------|---------------------------------------|
| OMR Mgmt. triggered (8.3.2) | Jan. 1 - Jun. 30 (when ≥ 5% of spring-run or winter- run in Delta) | In effect | - 5% of the Winter-run or Spring-run population in Delta | Winter-run = 74- 85% estimated in the Delta; Spring- run = 58-75% estimated in the Delta | Potential increase in presence of winter and spring-run | 2/14/22 | Based on 2/8/22 SaMT discussion |
| Winter-run yearly loss (8.6.1) | Nov. 1 - Jun. 30 | In effect WY 2022 loss = 1462.94 | 731.47 (50% of 1.17% of JPE) | Current yearly WR loss (natural) = 19.99; Current yearly WR loss (hatchery) = 0 | Possible additional salvage of natural Winter-run | 2/14/22 | Based on salvage data from 2/13/22 |

| Action | Timeframe | Current Action Status | Threshold(s) | Current Relevant Data | Weekly Trend | Last Updated | Comments |
|---|------------------|---|--|---|--------------|--------------|---------------------------------------|
| Winter-run discrete daily loss (8.6.2) | Nov. 1 - Dec. 31 | Not in effect | 12/1-12/31: loss 26/day unclipped older juv. Winter-run | max single daily loss from previous week = 3.52 fish | N/A | 1/3/22 | N/A |
| Mid and late season Winter-run daily loss threshold (8.6.3) | | In effect | Feb 1-Feb 28 (0.0000991% of JPE) = 12.39 | max single daily loss of Winter-run from previous week = 2.88 fish (observed at CVP on 1/28/22) | N/A | 2/14/22 | Based on salvage data from 2/13/22 |
| Spring-run surrogate protection (8.6.4) | Feb. 1 - Jun. 30 | Not in effect (Fish has not been released) | TBD (based on the number of fish released) | N/A | N/A | 2/7/22 | N/A |

Table 3b: Delta Smelt

| Action | Timeframe | Current Action Status | Threshold(s) | Current Relevant Data | Weekly Trend | Last Updated | Comments |
|--|---------------------|---|---|------------------------------|-----------------------|--------------|--|
| Integrated EarlyWinter Pulse Protection ('First Flush') (8.3.1) | Dec. 1 - Jan. 31 | Not in effect, triggered on Dec 17, last day of action was 1/2/22 | - three-day Freeport daily flow running avg > = 25,000 <u>AND</u> [three-day Freeport turbidity running avg > = 50 NTU <u>OR</u> Smelt Monitoring Team recommendation] | N/A | N/A | N/A | N/A |
| Turbidity Bridge Avoidance (8.5.1) | Dec. 15 - Apr. 1 | In effect, not triggered | Occurs after the Integrated Early Winter Pulse protection or February 1 (whichever until April 1) comes first - avg. OBI turbidity > 12 NTU | OBI 3.78 FNU | No change expected | 2/14/22 | Data from 2/13/22 |
| Larval and/Juvenile Delta smelt Protection (8.5.2) | ongoing | In effect, not triggered | - 5-day cum. salvage of juv. DS >= 1 [average 3-yr FMWT index + 1] <u>OR</u>, 3-day cum. salvage of juv. DS > 11 | current 5-day salvage = 0 | no change expected | 2/14/22 | Based on salvage data from 2/13/22 |

Table 3c: Longfin Smelt

| Action Early Adult Protection (8.3.3) | Timeframe Dec. 1 - Feb. 28 | Current Action Status Offramped | Threshold(s) - Cum. salvage > [most recent FMWT/10] = 1 fish (Sept Oct. Index) <u>OR</u> - Smelt Monitoring Team determines high likelihood of LFS movement into high-risk | Current Relevant Data N/A | Weekly Trend N/A | Last Updated N/A | Comments N/A |
|--|--|--|---|--|--|------------------------|-------------------------------------|
| OMR Mgt. for Adults (8.4.1) | Dec. 1 -Feb. 28 | Offramped | areas - Smelt Monitoring Team recommendation | N/A | N/A | N/A | N/A |
| Larval and Juvenile Longfin Smelt Entrainment Protection (8.4.2) | Jan 1 – Jun 30 | ln effect, Triggered | LFS larvae or juveniles in >=4 SLS or 20 mm stations in central and south Delta, OR LFS catch/tow >5 larvae or juveniles in >=2 stations | SLS 2 detected larvae at 4 stations in the south and central delta | Hatching started and will increase | 2/13/22 | SLS 2 sampled week of 1/24/22 |
| High Flow OMR Off-Ramp for Longfin Smelt (8.4.3) | Based on the status of 8.3.3, 8.4.1, & 8.4.2 | In effect, not triggered | - Sac. R. at Rio Vista >55,000, <u>OR</u> SJR at Vernalis >8,000 | Rio Vista = 9,000 to 11,000 cfs SJ = 9,00 to | N/A | 2/14/22 | N/A |
| | | | | 1,400 cfs | | | |

Table 3d: OMR

| Action | Timeframe | Current Action Status | Threshold(s) | Current Relevant Data | Weekly Trend | Last Updated | Comments |
|------------------------------------|---------------------|-----------------------------|--|-----------------------------|--------------|-----------------|----------|
| OMR Mgmt. Offramp (8.3.2) | Jun. 1 – Jun. 30 | Not in effect | 95% of the Winter-run and Spring run populations have migrated past Chipps Island <u>AND</u> Current daily average water temperature at Mossdale exceeds22.2°C for 7 nonconsecutive days in June <u>AND</u> Current daily average water temperature at Prisoners Point exceeds 22.2°C for 7 nonconsecutive days in June. Current daily average water temperature at CCF is greater than 25°C for three consecutive days | N/A | N/A | 10/4/21 | N/A |

| Monitoring Survey | Notes (as of 2/15/2022) | Status * |
|--|-------------------------|----------|
| Delta | N/A | N/A |
| SWP regular counts, CWT reading, and larval sampling | Active | 1 |
| CVP regular counts, CWT reading, and larval sampling | Active | 1 |
| Smelt Larval Survey | Active | 1 |
| LEPS | Active | 1 |
| 20mm Survey | Not Active | 4 |
| Spring Kodiak Trawl | Active | 1 |
| Fall Mid-water Trawl | Not Active | 4 |
| Summer Townet Survey | Not Active | 4 |
| Bay Study | Active | 1 |
| DJFMP- Chipps and Sacramento Trawls | Active | 1 |
| DJFMP- Seines | Active | 1 |
| EDSM | Active | 1 |
| EMP | Active | 1 |
| Mossdale | Active | 1 |
| USGS Flow monitoring | Active | 1 |
| Sacramento River | N/A | N/A |

Table 4. Fish monitoring gear efficiency and disruptions.

| Monitoring Survey | Notes (as of 2/15/2022) | Status * |
|--|--|----------|
| Red Bluff Diversion Dam screw trap | Active | 1 |
| Knights Landing screw trap | Active | 1 |
| Tisdale screw trap | Active | 1 |
| GCID screw trap | Active (not active since 2/7) | 2 |
| Redd dewatering and stranding surveys | Not Active | 4 |
| Sacramento Carcass and Redd Surveys | Active | 1 |
| Lower Sacramento Rotary Screw Trap | Active (cones raised on 2/6, trapping resumed 2/7) | 2 |
| Feather River | N/A | N/A |
| Feather River screw trap | Active (weekdays only) | 1 |
| San Joaquin River | N/A | N/A |
| SJRRP CDFW Field Monitoring | Active | 1 |
| SJRRP USFWS and USBR Field Monitoring | Active | 1 |
| Stanislaus Fish Weir | Active | 1 |

* Status: Weekly categories include:

[1] Active (ongoing sampling)

[2] Partial Interruption (some sampling interruptions)

[3] Interrupted (sampling fully suspended)[4] Not Active (sampling not scheduled)