

Weekly Assessment of CVP and SWP Delta Operations on ESA-listed Species

Executive Summary

a) Operations anticipated during the week

See Weekly Fish and Water Operation Outlook document for February 1 – February 7

b) Winter-run Chinook Salmon summary

Loss of natural winter-run Chinook Salmon (by length at date, LAD) has occurred in the past week at the Federal fish salvage facility (WY 2022 total loss = 19.99 fish, as of 1/30/2022). Loss of natural winter-run Chinook Salmon at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities could occur over the next week based on hydrology and recent observations. 74-85% of juvenile natural winter-run Chinook Salmon from brood year (BY) 21 are estimated to be present in the Delta. The Delta Cross Channel (DCC) gates closure for the season reduces exposure of winter-run Chinook Salmon juveniles that are present in the Sacramento River near the DCC gates into the interior Delta.

c) Spring-run Chinook salmon summary

Juvenile natural spring-run Chinook Salmon from BY 21 are migrating past the DCC gates. The exposure and effects of DCC closure are unlikely for natural spring-run Chinook Salmon. Length-at-date spring run were observed in the Delta within the previous seven days. Larger, older juveniles were observed that may be yearling spring run. 57-72% young of year spring-run Chinook Salmon are estimated to be in the Delta.

d) Central Valley Steelhead summary

Loss of natural Central Valley California (CCV) steelhead has not occurred in the past week at the State and Federal fish salvage facilities (WY 2022 total loss = 54.36 fish, as of 1/30/2022). Loss of Central Valley steelhead at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities could possibly occur over the next week. 27-39% of juvenile CCV Steelhead are estimated to be present in the Delta. Closure of the DCC gates for the season will reduce exposure to Central Valley steelhead juveniles that are potentially present in the Sacramento River near the DCC gates.

e) Green Sturgeon summary

Loss of green sturgeon has not occurred in the past week at the State and Federal fish salvage facilities. Loss of green sturgeon is unlikely to occur over the next week due to their rare presence in the South Delta.

f) Delta Smelt summary

Based on distribution patterns over the past decade and rare detections, Delta Smelt are unlikely to be prevalent in the South Delta, but a 54mm, adipose clipped, Delta Smelt was salvaged at the Tracy Fish Collection Facility on 1/16/2022. Limited detection data support Delta Smelt being present in the Sacramento Deep Water Ship Channel, Lower San Joaquin River, South Delta, Suisun Marsh, and the Lower Sacramento River. Two marked individuals have been collected since 1/25/2022. High winds may elevate turbidity at OBI between 2/1-3/2022 but a turbidity bridge avoidance action will not be possible since the OMR Index will be less negative than -2,000 cfs. The likelihood of Delta Smelt adult entrainment may elevate if turbidity increases in the OMR corridor. The expected less negative OMR Index should decrease the overall likelihood of entraining adults. Water temperatures are within the range for Delta Smelt spawning.

g) Monitoring Teams summary

There were no non-consensus issues to report from the Salmon Monitoring Team.

There were no non-consensus issues to report from the Smelt Monitoring Team.

Operational and Regulatory Conditions

See current Weekly Fish and Water Operation Outlook document.

Biology, Distribution, and Evaluation Winter-run Chinook salmon, Spring-run Chinook salmon, Central Valley Steelhead

POPULATION STATUS

Winter-run Chinook salmon

Delta Life Stages:

Juveniles, Adults

Brood Year 2021 Productivity:

- Natural winter-run Chinook salmon: A draft juvenile production estimate (JPE) calculation has been established for brood year (BY) 2021 winter-run Chinook salmon. The final winter-run Chinook salmon JPE is 125,038 fish. Among other topics, the group discussed the thiamine vitamin deficiency that is being observed again in broodstock at the Livingston Stone NFH similar to last year's observation. Last year the thiamine vitamin deficiency appeared to negatively affect survival of juvenile fish as they migrate downstream towards the Delta. Mean cumulative weekly passage of winter-run Chinook salmon through 1/28/2022 for the last 19 years of passage data is 98.4% (one SD of 1.9%). By 1/14/2022, 571,662 winter-run Chinook salmon were estimated to have passed RBDD compared to the cumulative passage last year of 3,813,580 winter-run Chinook salmon.
- Hatchery winter-run Chinook salmon:
 - No hatchery winter-run Chinook salmon have been released in WY 2022.

Spring-run Chinook salmon

Delta Life Stages:

Young-of-year (YOY) and Yearlings

Brood Year 2021 Productivity:

- Natural spring-run Chinook salmon: No JPE has been established for spring-run Chinook salmon. Approximately 16.2% juvenile spring-run sized Chinook salmon have been observed passing Red Bluff Diversion dam as of 1/30 based on historical data.
- Hatchery spring-run Chinook salmon surrogates:

- Approximately 84,343 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 12/15/2021. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 145mm. This is the first spring-run Chinook salmon surrogates release group. There has been loss this water year of fish associated with the first surrogate release group.
- Approximately 82,626 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 12/22/2021. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 125mm. This is the second spring-run Chinook salmon surrogates release group. There has been loss this water year of fish associated with the second release group.
- Approximately 77,325 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 1/6/2022. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 145mm. This is the third spring-run Chinook salmon surrogates release group. There has been loss this water year of fish associated with the third release group.
- The agencies in the SaMT discussed the thiamine vitamin deficiency that was observed in winter run Chinook salmon broodstock at the Livingston Stone NFH in BY 2021. Last year the thiamine deficiency appeared to negatively affect survival of juvenile fish as they migrate downstream towards the Delta. The thiamine deficiency issue is also likely impacting spring-run Chinook salmon. Although the egg take goals have been met at the Feather River Fish Hatchery, they are still experiencing fertility issues that are impacting production.

Central Valley Steelhead

Delta Life Stages:

- Spawning Adults, Kelts, Juveniles

Brood Year 2021 Productivity:

- Spawner abundance: There is limited information about the adult steelhead population. It is estimated to be small, contributing to the limited productivity of the population
- Natural steelhead: No JPE has been established for steelhead. Data are limited.
- Hatchery steelhead: Reclamation's Proposed Action has no hatchery steelhead triggers.
 - Approximately 610,911 steelhead from Coleman NFH were released at Battle Creek from 12/12/2021 to 12/13/2021. This group is 100%

marked only (with an adipose-fin clip) and has an estimated average fork length of 195mm.

- Approximately 57,056 steelhead from Mokelumne River Fish Hatchery will be released into the Mokelumne River at Feist Ranch from 1/31/2022 to 2/1/2022. This group is 100% marked (CWT and adipose fin clip) and has an estimated average fork length of approximately 210mm.
- Approximately 424,000 steelhead from Nimbus Fish Hatchery will be released into the Lower American River at the Sunrise Boat Ramp from 1/31/2022 through 2/4/2022. This group is 100% marked (adipose fin clip) and has an average fork length of approximately 240mm.

DISTRIBUTION

Winter-run Chinook Salmon

Current Distribution:

- On 2/1/2022, SaMT estimated 74-85% of juvenile winter-run Chinook salmon were present in the Delta and 1-5% were estimated to have exited the Delta (Table 1). The SaMT estimate is largely based on historical trends since few winter-run Chinook salmon have been seen in recent monitoring. The SaMT expressed some concern that these numbers seem low especially given the precipitation events that occurred early in the Water Year (October 2021).
- Combined total natural winter-run Chinook salmon loss equals 19.99 fish (as of 1/30/2022).
- Fish have been steadily arriving since the beginning of October 2021.
- Winter-run Chinook Salmon have not been observed in RST monitoring locations over the past week (Knights Landing, Tisdale, Sac Trawls, Beach Seines) however the fish appear to no longer be holding in the middle reaches of the Sacramento River and are migrating downstream (Table 2).
- Movement of winter-run Chinook Salmon juveniles into the lower reaches of the Sacramento River and upper Delta is continuing. Mill and Deer creeks daily flows were recorded higher than 95 cfs during the past week.

Historic Trends

- Based on historical trends in salvage, 24.5% of winter-run Chinook salmon should have been observed in salvage by this time of the water year (Table 3). Loss of natural winter-run Chinook salmon at the CVP and SWP fish collection facilities could occur over the next week based on hydrology and

recent observations. If historic trends in salvage were to continue, winter-run Chinook salmon loss is expected to remain the same over the next week.

- Based on historic loss patterns and historic cumulative loss, it is possible that in WY 2022 50% of the wild winter-run Chinook salmon Delta Performance single-year loss threshold (731.47 wild winter-run Chinook salmon, as set by the JPE estimate) could be exceeded (Figure 1, Operations Outlook Table 2a). However, members of the SaMT agree the likelihood of exceeding the next annual loss threshold and loss occurring in the next week is unlikely to lead to exceedance of the threshold.

Forecasted Distribution within Central Valley and Delta regions

- Movement of winter-run Chinook salmon juveniles into the lower reaches of the Sacramento River and upper Delta are likely to occur with precipitation events and increasing river flows and turbidity. The agencies in the SaMT don't believe precipitation events will occur over the next week (see Ops Outlook). The STARS model projects route-specific proportion of entrainment, survival, and travel times (Table 4). This model does not estimate entrainment into the lower Sacramento River sloughs (i.e., Three-Mile Slough). The DCC gates were closed 11/30/21 and are expected to remain closed through mid-May 2022. If little precipitation is forecasted there may be a need to open the DCC gates to meet D-1641 water quality standards
- The entrainment tool estimates a median loss of 0 fish and a maximum loss of 12 fish during this week (SacPAS last updated on 1/31/2022, Table 5, Figure 2).
- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.
- For preliminary results of acoustically tagged salmonids currently in the system see Table 6.

Spring-run Chinook salmon

Current Distribution

- On 2/1/2022, SaMT estimated 57-72% of young of year CV spring-run Chinook salmon were present in the Delta (Table 1).
- Spring-run chinook are being observed at upstream monitoring sites including Tisdale, Knights Landing, on the Feather River and Butte Creek, and in Beach Seines.

Historical Trends

- Based on historical trends in salvage, 0% of YOY spring-run Chinook salmon were observed in salvage by this time of the water year (Table 3). If historic trends in salvage were to continue YOY spring-run Chinook salmon loss is unlikely to increase over the next week.

Forecasted Distribution within Central Valley and Delta regions

- On 2/1/2022 SaMT noted that some juvenile CV YOY spring-run Chinook Salmon have yet to emerge. Yearling CV spring run Chinook Salmon are moving out from natal tributaries. Mill and Deer creek flows continue to exceed 95 cfs indicating that yearling spring-run Chinook Salmon may begin to move and migrate into the mainstem Sacramento River (Table 4).
- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.
- For preliminary results of acoustically tagged salmonids in the system see Table 6.

Central Valley Steelhead

Current Distribution

- On 2/1/2022 SaMT estimated 27-39% of juvenile CCV steelhead were present in the Delta (Table 1).
- Combined total loss of hatchery steelhead equals 130.3 fish as of 1/30/2022.
- Combined total loss of natural steelhead between December 1 and March 31 equals 54.36 fish as of 1/30/2022.

Historical Trends

- Based on historical trends in salvage, 11.9% (December through March) of juvenile CCV steelhead should have been observed in salvage by this time of the water year. If historic trends in salvage were to continue, juvenile CCV steelhead loss is likely to not increase over the next week.

Forecasted Distribution within Central Valley and Delta regions

- Natural steelhead were not observed in key monitoring locations this past week (Tisdale and Knight's landing).
- SaMT estimated that 27-39% of the population of CCV steelhead may be present in the Delta at this time. Closure of the DCC gates for the season will

reduce exposure and possible entrainment of juvenile CCV steelhead into the interior Delta via the DCC gates.

- The entrainment tool estimates a median loss of 0 fish and a maximum loss of 17 fish during this week (SacPAS last updated on 1/31/2022, Table 5, Figure 2).
- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.

Table 1. Salmonid distribution estimates

Location	Yet to Enter Delta	In the Delta	Exited Delta past Chipps Island
Young-of-year (YOY) winter-run Chinook salmon	10-25%	74-85%	1-5%
YOY spring-run Chinook salmon	28-43%	57-72%	0%
YOY hatchery winter-run Chinook salmon*	NA	NA	NA
Natural origin steelhead	60-70%	27-39%	1-3%

Table 2. Historic migration and salvage patterns.

Date (1/30)	Red Bluff Diversion Dam	Tisdale RST	Knights Landing RST	Sac Trawl (Sherwood) Catch Index	Chipps Island Trawl Catch Index	Salvage
Chinook, Winter-run, Unclipped	98.0% (96.2%,99.8%) BY: 2012 - 2020	77.8% (53.0%,102.7%) BY: 2012 - 2020	73.8% (46.0%,101.5%) BY: 2013 - 2020	39.8% (9.3%,70.3%) BY: 2012 - 2020	3.9% (-0.5%,8.2%) BY: 2012 - 2020	24.5% (3.8%,45.2%)
Chinook, Spring-run, Unclipped	16.2% (4.0%,28.4%) BY: 2012 - 2020	25.4% (-1.3%,52.1%) BY: 2012 - 2020	23.9% (-2.8%,50.5%) BY: 2013 - 2020	5.2% (-3.5%,13.8%) BY: 2012 - 2020	0.0% (0.0%,0.0%) BY: 2012 - 2020	0.0% (0.0%,0.0%)
Steelhead, Unclipped (December - March)	N/A	N/A	N/A	N/A	N/A	11.9% (-2.5%,26.4%)

Table 3. STARS model output.

Date (1/30/2022)	DCC	Georgiana Slough	Sacramento River	Sutter and Steamboat
Proportion of Entrainment	0	0.27	0.46	0.27
Survival	N/A	0.19	0.54	0.42
Travel Time	N/A	16.7d	10.3d	10.7d

Table 4. Mean daily flow and percent change (Wilkins Slough, Deer Creek, Mill Creek; cfs from CDEC) and temperature and percent change (Knights Landing; °F from RST).

Date	Mill Creek flow (MLM)	MLM Change	MLM Alert	Deer Creek flow (DCV)	DCV Change	DCV Alert	Wilkins Slough flow (WLK)	Knights Landing temperature (°F)	Alert Triggered
1/30/2022	170.7	-0.3%	Flow> 95cfs	155.5	-1.8%	Flow> 95cfs	5924.1	N/A	N/A
1/29/2022	171.3	-1.4%	Flow> 95cfs	158.3	-2.2%	Flow> 95cfs	6045.0	N/A	N/A
1/28/2022	173.7	-2.0%	Flow> 95cfs	161.9	-3.1%	Flow> 95cfs	6193.5	N/A	N/A
1/27/2022	177.3	-3.3%	Flow> 95cfs	167.2	-3.2%	Flow> 95cfs	6311.1	41.4	N/A
1/26/2022	183.4	-3.6%	Flow> 95cfs	172.8	-2.5%	Flow> 95cfs	6426.2	41.7	N/A
1/25/2022	190.2	-0.2%	Flow> 95cfs	177.2	-1.0%	Flow> 95cfs	6550.0	41.7	N/A
1/24/2022	190.5	-0.1%	Flow> 95cfs	178.9	-1.1%	Flow> 95cfs	6667.7	41.6	N/A

Table 5. a) WY 2022 loss and salvage predictor data: Predicted weekly loss of winter-run Chinook salmon and steelhead at CVP and SWP facilities. b) Environmental details, current and forecast.

Week	17	18
a)	Model	N/A
Steelhead median	0	0
Steelhead high	17	26
Winter-run Chinook median	0	0
Winter-run Chinook high	12	18

Week	17	18
b)	Data	Forecast
Temperature (Mallard Island, C)	10.1	10.0
Precipitation (5-d running sum, inches)	0.00	0.00
Old + Middle river flows (cfs)	-5,140	-5,140
Sacramento River flow (Freeport, cfs)	12,542	12,542
DCC Gates	Open	Closed
San Joaquin River flow (Vernalis, cfs)	898	898
Export	5,666	5,666

Table 6. Preliminary survival (Φ) results for acoustically tagged salmonids.

Project	Group (n)	Release Date(s)	Release Location	Min. Φ to Tower Bridge	Min. Φ to Benicia Bridge (East Span)	Min. through-Delta Φ (Sac to Benicia)	Notes
Hatchery-origin Chinook salmon: Seasonal Survival Study	Week 1 (200)	12/15/2021 – 12/16/2021	RBDD	75% (SE = 3.1)	16.6% (SE = 2.7)	22.2% (SE = 3.4)	Fish detected at Old River receiver

Table updated 1/31/2022. Additional information for all projects can be found online at <https://oceanview.pfeg.noaa.gov/CalFishTrack/>




FIGURE 1. WY 2022 cumulative natural winter-run Chinook salmon loss (blue) and 2009 – 2018 historic cumulative loss (gray, different symbols). Historic daily mean plotted in black circles.




FIGURE 2. Predicted weekly loss of steelhead and winter-run Chinook salmon at the CVP and SWP facilities

EVALUATION

1. **After January 1, are more than 5% of juveniles from one or more salmonid species present in the Delta?**

Greater than 5% of juvenile winter-run Chinook salmon, CCV steelhead, and young of year spring-run Chinook salmon are present in the Delta.

2. **Does the operational outlook's ranges impact fish movement and change the potential distribution of fish?**

Potential effects within the 7 days (near-term) in the operations outlook.

OMR flow is expected to remain at or more positive than -5,000 cfs this upcoming week.

Potential effects longer than the 7 days (longer-term) in the operations outlook.

See response above to (2) (i).

3. **What is the likelihood of increased loss exceeding the next annual loss threshold (50%, 75% or 90% of threshold) resulting in OMR management actions based on population distribution, abundance, and behavior of fish in the Delta?**

Winter-run Chinook salmon

Total juvenile natural winter-run Chinook salmon (LAD) loss is 19.99 fish (as of 1/20/2022). Loss of juvenile winter-run Chinook salmon has occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believe that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold.

Spring-run Chinook salmon

Total juvenile spring-run Chinook salmon (LAD) loss is 0 fish (as of 1/30/2022). Loss of juvenile spring-run Chinook salmon has not occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believes that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold.

Central Valley Steelhead

Total natural juvenile steelhead loss (December 1 through March 31) is 54.36 fish (as of 1/30/2022). Loss of natural juvenile has not occurred in the past week at the CVP and SWP fish salvage facilities. Total clipped steelhead loss is 130.3 fish (as of 1/30/2022). The agencies in the SaMT assessed the likelihood of exceeding the next

annual loss threshold and believe that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold.

4. If an annual loss threshold has been exceeded, do continued OMR restrictions benefit fish movement and survival based on real-time information?

Winter-run Chinook salmon

The annual loss threshold for winter-run Chinook salmon has not been exceeded in WY 2022.

Spring-run Chinook salmon

The annual loss threshold for spring-run Chinook salmon has not been exceeded in WY 2022.

Central Valley Steelhead

The annual loss threshold for steelhead (December 1 – March 31) has not been exceeded in WY 2022.

5. If OMR is more negative than -5,000 cfs are there changes in spawning, rearing, foraging, sheltering, or migration behavior beyond those anticipated to occur under OMR management at -5,000 cfs?

OMR flows are not expected to be more negative than -5,000 cfs during the next week.

Biology Distribution and Evaluation of Green Sturgeon POPULATION STATUS

Delta Life Stages:

- Adults and Juveniles

Juvenile Abundance:

- In 2021, 1037 larval green sturgeon and 5 juvenile green sturgeon were observed at the Red Bluff Diversion Dam during continuous fish monitoring using RSTs in the upper Sacramento River. 14 juveniles were captured and implanted with micro-acoustic tags during the month of October, and all were presumed to leave the upper Sacramento River during the unprecedented storm /run-off event that occurred on October 24, 2021 that resulted in flows at Bend Bridge of ~37,000 cfs.

DISTRIBUTION

Current Distribution

- Adults: Most abundant during spring spawning migration period of March through May, and post spawning out-migration periods May through June; October through January depending on first winter storm event resulting in

significant Sacramento River flow increases. Adult presence year-round to a lesser extent mainly in San Pablo Bay. Recent sporadic occurrences of adult green sturgeon in the San Joaquin River system but spawning has not been documented. Unknown if spawning occurred historically in the San Joaquin River system.

- Juveniles: Age-1 through Age-3 juveniles present year-round and widely distributed. Juveniles tagged with acoustic tags in the main channel Sacramento River near Sherman Island detected in the Sacramento River as far upstream as the Cache Slough complex, in the San Joaquin River at the Antioch Bridge, in Threemile, Horseshoe Bend, and Montezuma Sloughs. Seasonal abundance at the primary sampling site (near Sherman Island) appears to be highest during summer in based on capture and telemetry data. Residence time at the primary sampling site for individual fish ranges from one day to over one year but telemetry data show outmigration from the primary sampling site to the Pacific Ocean ranges from 27 to 552 days. Recent capture data shows diurnal depth preference in the main channel of the Sacramento River. No recent documentation of shallow water habitat presence or foraging but likely. The remains of a juvenile/young adult green sturgeon were collected at CVP on 1/6/22; however, this does not count towards green sturgeon salvage totals.

Historical Trends

- Juvenile and adult green sturgeon are historically present in the San Joaquin and Sacramento rivers and Delta

Forecasted Distribution within Central Valley and Delta regions

- Juvenile and adult green sturgeon are present in the San Joaquin and Sacramento rivers and Delta during the next week.

EVALUATION

1. Is there likely to be salvage that may exceed the annual loss limit?

Green sturgeon salvage is 0 fish (as of 1/30/2022). The agencies in the SaMT assessed the likelihood of salvage occurring in the next week is unlikely to occur.

Biology, Distribution, and Evaluation of Delta Smelt POPULATION STATUS

Delta Smelt Life Stages:

Subadults and Adults

Brood Year 2021:

Abundance estimate:

The All Strata abundance estimate for Delta Smelt was 1,301 for data collected between 1/24-28/2022. The most recent detection of Delta Smelt was on 1/31/2022 (2 individuals, marked, EDSM) caught in the Sacramento Deep Water Shipping Channel stratum.

Biological Conditions:

Adult Delta Smelt are expected to be present in the Sacramento Deep Water Ship Channel, the Lower Sacramento, Suisun Bay, South Delta, and Lower San Joaquin River based on the survey detections. Delta Smelt are expected to have migrated in response to increases in turbidity and flow from “first flush” conditions (Sommer et al 2011). Delta water temperatures have reached the range in which Delta Smelt spawning may be occurring (Damon et al 2016). The Smelt Monitoring Team discussed the most recent monitoring data (Table 4) and considered professional opinion on the historical trends in regional distribution.

DISTRIBUTION

Current Distribution

- Real time detection data is currently limited to EDSM sampling, Chipps Island, Bay Study, LEPS, and SLS. Since there are only a few recent detections of Delta Smelt, the Smelt Monitoring Team’s capacity to estimate where they are within the Delta is limited.
- The last Delta Smelt detection was on 1/31/2022 in the Sacramento Deep Water Shipping Channel stratum.
- Larval sampling at the Skinner Fish Facility (SFF) and the Tracy Fish Collection Facility (TFCF) will begin on 2/7/2021.

Table 6. Summary of recently reported detections of Delta Smelt by Region and Salvage Facilities between 1/25/2022 and 1/31/2022. Start and End dates reflect period of time between updates to SMT. Regional categories are determined from EDSM sampling. Delta Smelt >58mm FL are considered adults. Subadult fish are considered by the SMT to be fish from the previous year’s cohort based on size and timing of collection. Young of year are considered juveniles and larvae.

Life Stage	North	South	West	Far West	Salvage
Adult	2	0	0	0	0

Life Stage	North	South	West	Far West	Salvage
Subadult	0	0	0	0	0
Larvae/Juvenile	0	0	0	0	0

Table 7. Summary of recent Delta Smelt detections reported since last assessment and the total detections for the current water year. Notes reflect latest information on reported detections or completion of survey for the water year and include both larval and adult detections. Total Fish counts do not distinguish between hatchery origin and wild Delta Smelt. Table may include preliminary data that may not have received full QA/QC, but any corrections will be made the following week. * This individual was collected by FCCL broodstock collection on 12/22/2021 and genetically identified as a hybrid.

Sampling Method	Frequency	New Detections	WY2022	Notes
EDSM	Weekly	2	21	Phase 1 began 11/29/2021
SKT	Monthly	0	0	Ongoing
SLS	Biweekly	0	0	Survey 2: Processing
20-mm	Biweekly	0	0	Begins: 3/21/2022
Summer Trawnet	Biweekly	0	0	Complete
Bay Study	Monthly	0	0	Ongoing
FMWT	Monthly	0	0	Complete
Chipps Island Trawl	Weekly	0	1	Ongoing
FCCL Brood Stock Collections	Weekly	0	1*	Ongoing
LEPS	Weekly	0	0	Ongoing
TFCF	Daily	0	1	Ongoing
Total	N/A	N/A	23	Sum of all Delta Smelt observed during the OMR Management Season

Cultured Delta Smelt Experimental Releases

- On 12/14/2021-12/15/2021, the Experimental Release Technical Team trucked approximately ~12800 brood year 2021 Delta Smelt from the FCCL to Sacramento River at Rio Vista for release into the Delta. This release will include 11392 Adipose fin clipped individuals and 1408 Visible Implant Elastomer (VIE) tagged individuals.
- On 01/11/2022-01/12/2022, the Experimental Release Technical Team will truck approximately ~12800 brood year 2021 Delta Smelt from the FCCL to Sacramento River at Rio Vista for release into the Delta. This release will include 12416 Adipose fin clipped individuals and 384 Visible Implant Elastomer tagged individuals.
- Genetic analysis was completed on the two potentially wild Delta Smelt and they were determined to be of hatchery origin with poor quality fin clips. A third smelt collect by the FCCL was identified as a hybrid and additional analysis is being run to determine parentage.

Table 8. Weekly summary of the origin of Delta Smelt. These identifications are considered tentative and additional genetic testing will confirm the identity of individuals. Individuals with no tags are provided alive to the FCCL as potential additions to the FCCL Broodstock.

Date	Survey	Stratum	Total Caught	Ad. Clipped	VIE	No Tag
1/31/2022	EDSM	Sacramento Deep Water Shipping Channel	2	2	0	0

Historical Trends

- Wild Delta Smelt detections in the Sacramento Deep Water Ship Channel indicate presence upstream of the confluence, but the fish may be freshwater residents and not representative of the migratory life history patterns in Delta Smelt (Hobbs 2019).
- Historically, the highest peak in salvage is in May and the second highest is in June (Grimaldo et al 2009; figure 5).

Forecasted Distribution within Central Valley and Delta regions

- Predicting the distribution of adult wild Delta Smelt is currently difficult because detection data is limited to a few individuals and historic patterns may not be representative of the low population levels.

- The SMT uses turbidity as a surrogate for Delta Smelt presence and in making assessments of the likelihood of entrainment for larval Delta Smelt after spawning begins.
- Given water temperatures are conducive to spawning, increasing turbidity and spring tides may influence the distribution of adults.
- The potential of experimentally released Delta Smelt to distribute from their release site is unknown at this time and SMT cannot predict their distribution beyond the original release site and subsequent recaptures.

ABIOTIC CONDITIONS

Turbidity

- As of 1/31/2022 turbidity continues to be less than 12 FNU at Old River at Bacon Island (OBI) (See attachment B).
- Turbidity in the Delta may further increase as a consequence of high winds. North to Northwest winds as high as 40 mph are predicted for 2/1-2/2022.
- Integrated Early Winter Pulse Protection ended on 1/2/2021.
- Precipitation is not expected in the next seven days.
- Three Station Daily Average water temperature on 1/31/2021 was 10.46° C. This temperature is within the range of Delta Smelt spawning; 9-18 ° C (Damon et al 2016).

Table 9. Relevant Environmental Factors to the current management actions for Delta Smelt.

Date Reported	OBI Daily Turbidity (FNU)
2/1/2022	4.77

X2 Conditions

- X2 is estimated to be at 85 km.
- When X2 is above 81 km, the SMT uses the X2_EC_Graph.xlsx tool to estimate the position of X2 for both the Sacramento and San Joaquin Rivers and assumes the average of the two is representative of an approximate X2 position.

Other Environmental Conditions

- The Fish and Water Operation Outlook OMR Index values are expected to range between -500 to -1,500 cfs from 2/1/2022 to 2/7/2022.
- Real time tracking of environmental conditions, relevant thresholds and Delta Smelt catch data are updated daily at:
http://www.cbr.washington.edu/sacramento/workgroups/delta_smelt.html

EVALUATION

1. Between December 1 and January 31, has any first flush condition been exceeded?

The running 3-day average flows and running 3-day average turbidity at Freeport exceeded the triggers for Integrated Early Winter Pulse Protection on 12/17/2021 (running 3-day averages: 27,152 cfs and 67 FNU). The CVP and SWP reduced exports from 12/20/2021 until 1/2/2022.

2. Do DSM have a high risk of migration and dispersal into areas at high risk of future entrainment? (December 1- January 31)

Based on distribution patterns over the past decade and recent detection data, Delta Smelt are unlikely to be prevalent in the South Delta. Delta Smelt are expected to have made their migration in response to first flush and be holding in anticipation for temperatures conducive to spawning. Turbidity may become elevated due to high winds. The likelihood of entraining adult delta smelt may increase if turbidity increases in the OMR corridor. However, the OMR index will already be more positive than possible under a turbidity bridge avoidance action.

3. Has a spent female been collected?

No spent female has been caught by SKT as of 2/1/2022

4. If OMR of -2000 does not reduce OBI turbidity below 12NTU/FNU, what OMR target is deemed protective between -2000 and -5000?

OBI turbidity is currently below 12 FNU and a turbidity bridge avoidance action is not possible even if turbidity increases due to wind in the next seven days since the OMR index is already less negative than -2,000.

5. If OBI is 12 NTU/FNU, what do other station locations show?

OBI turbidity is currently below 12 FNU. The daily average turbidities on 1/31/2022 at Prisoners Point (6.91 NTU), Holland Cut (station offline) and Victoria Canal (3.77 NTU) are likely to increase after high winds and expected to decrease after windspeeds reduce in the next seven days.

6. If OBI is 12 NTU/FNU, is a turbidity bridge avoidance action not warranted? What is the supporting information?

The turbidity at OBI is below 12 FNU, and a turbidity bridge avoidance action is not possible in the next 7 days due to the OMR index being less negative than -2,000 cfs.

7. After March 15 and if QWEST is negative, are larval or juvenile DSM within the entrainment zone of the CVP and SWP pumps based on surveys?

This question is not applicable until March 15th.

8. Based on real-time spatial distribution of Delta Smelt and currently available turbidity information, should OMR be managed to no more negative than -3,500?

This question is not applicable until March 15th.

9. What do hydrodynamic models, informed by EDSM or other relevant data, suggest the estimated percentage of larval and juvenile DSM that could be entrained may be?

This question is not applicable until March 15th

DELTA SMELT REFERENCES

- Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. *California Fish and Game* 102(4):188–210.
- Hobbs, J. A., Lewis, L. S., Willmes, M., Denney, C., & Bush, E. (2019). Complex life histories discovered in a critically endangered fish. *Scientific Reports*, 9(1).
<https://doi.org/10.1038/s41598-019-52273-8>
- Lenny F. Grimaldo, Ted Sommer, Nick Van Ark, Gardner Jones, Erika Holland, Peter B. Moyle, Bruce Herbold & Pete Smith (2009) Factors Affecting Fish Entrainment into Massive Water Diversions in a Tidal Freshwater Estuary: Can Fish Losses be Managed? *North American Journal of Fisheries Management*, 29:5, 1253-1270, DOI: 10.1577/M08-062.1
- Polansky, L., Newman, K.B., Nobriga, M.L. et al. Spatiotemporal Models of an Estuarine Fish Species to Identify Patterns and Factors Impacting Their Distribution and Abundance. *Estuaries and Coasts* 41, 572–581 (2018).
<https://doi.org/10.1007/s12237-017-0277-3>
- Sommer, T., F. Mejia, M. Nobriga, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science* 9(2).

Attachment A

Objective

Background

DSM2 model results: summary tables

DSM2 model results: figures

DSM2 model interpretation entrainment into Delta strata regions

DSM2 channel locations information

Objective

Weekly modeling efforts are conducted to examine the effects of varying OMR conditions on the behavior of salmonids present in the Delta in a one-week “look ahead” or outlook. Members of the Salmon Monitoring Team (SaMT) use DSM2 modeling results to help answer how changing pumping regimes translates to differences in flows and velocities modeled at various channel locations within the Delta and what impact modeled environmental parameters have on rearing, foraging, migrating, and holding salmonids.

Each series of runs consists of three OMR conditions: minimum and maximum scenarios bounded by expected OMR index values for that week (Ops Outlook, Table 1) and a baseline which represents an anticipated operational value. Assumptions are made to best estimate future hydrologic characteristics. These inputs are more confident for the future one, two, and three-day timeframes; days four through six have lower confidence. Model scenarios hold hydrology inputs between runs constant and adjust Delta export pumping rates to compare between scenario OMR index values (unless otherwise noted). Although hydrologic ensembles could be used, a single value or deterministic projection is used for efficiency.

SaMT members use weekly DSM2 model results from a range of scenarios as part of a suite of tools to help assess distribution and changes to behavior of salmonids. At each channel location over a six-day action period, environmental parameters are examined: modeled flow and velocity general statistics (e.g., magnitude, range, percent positive), differences in modeled flow and velocity values compared with the baseline scenario, etc. That information, in conjunction with channel location (e.g., close to the Delta pumping facilities, closer to areas with higher tidal influence, etc.) and other environmental considerations (e.g., tidal cycle, upcoming storms, etc.), is then interpreted from a biological perspective. SaMT explores the possible effects to salmonids of changing OMR index scenarios, assuming each of those potential operations could be that week’s controlling factor.

Background

Category	Notes
Process	Notes
Weekly process	<ul style="list-style-type: none"> • DSM2 model runs use a historic and forecasted hydrological input dataset with no assumptions provided by DWR Thursdays and updated by Reclamation Mondays. • Input File updated Monday after initial distribution from DWR for removal of forecasted in lieu of historic input. • Reclamation provides scenarios based on expected OMR index values for the upcoming week. • DSM2 model runs produced Monday and distributed to SaMT members.
Hydraulic footprint information	Notes
Updated	1/31/2022 (baseline and scenarios)
DSM2 modeling results range	2/1/2022 – 2/7/2022
OMR index value scenarios	Notes
Baseline	-950 cfs
Scenario 1	-500 cfs (difference of 450 cfs)
Scenario 2	-1,500 cfs (difference of 550 cfs)
Changes between scenarios	Notes
Hydrology	No (see special considerations section below)
Delta Exports	Yes
Common assumptions	Notes
DSM2 run results based on the following assumptions	<ul style="list-style-type: none"> • CCFB Gates are operating to Priority 2 through January, then to Priority 1 starting 2/1/2022. • The Delta Cross Channel gates were closed on November 30 and will remain closed through the end of the forecast period. • Suisun Marsh salinity control flashboards are in, and 2 of the Suisun Marsh salinity control gates are in open position as of December 21, and the remaining 1 gate is closed for maintenance. • See Figure A1a for Sacramento River flow at Freeport during the forecast period. • See Figure A1b for San Joaquin River flow at Vernalis during the forecast period. • Clifton Court Forebay and Tracy Pumping Plant pumping is shown for model runs in Figure A2.

Category	Notes
Additional information	Notes
Considerations for current DSM2 model run	<ul style="list-style-type: none"> The expected range in OMR is due to an anticipated range in Delta inflow hydrology, not export variability as is depicted herein. There are limitations in the DSM2 "set-up protocol" that does not coincide with reality.
Caveats	<ul style="list-style-type: none"> Time-step: DSM2 generates results at 15-minute time-steps. Visualizations of DSM2 model run results are aggregated over daily timesteps. Operations function on a more granular scale than daily time-step. <ul style="list-style-type: none"> Salmonid behavior: DSM2 provides flow fields which salmonids may encounter but salmonids are not neutrally buoyant particles. Models which incorporate behavior from acoustic tagged salmonids are being developed for South Delta (ePTM, ecoPTM).

DSM2 model results: figures




FIGURE A1. Daily forecasted Freeport and Vernalis flows (cfs).






FIGURE A2a-c. Daily forecasted pumping: Jones and Banks (cfs) for (a) Baseline, (b) Scenario 1, and (c) Scenario 2.

DSM2 model results: summary tables

Table A1. Summary of minimum, maximum, mean, and percent positive flows (cfs) and velocities (ft/s) by DSM2 channel for OMR scenarios over a 6 day time period. For scenario values refer to “Hydraulic Footprint Information” in the conditions / assumptions section above.

Scenario (cfs)	DSM2 Channel	Flow Min.	Flow Max.	Flow Mean	Flow % Positive	Velocity Min.	Velocity Max.	Velocity Mean	Velocity % Positive
Baseline-950	6	-736.8	2663.1	1856.3	98	-0.3	1.4	0.9	98
OMR-1500	6	-683.8	2663.4	1856.4	98	-0.3	1.4	0.9	98
OMR-500	6	-736.8	2671.2	1856.3	98	-0.3	1.4	0.9	98
Baseline-950	21	-8218.7	7416.4	740.1	55	-0.5	0.5	0.1	55
OMR-1500	21	-8260.2	7385.8	725.4	55	-0.5	0.5	0.1	55
OMR-500	21	-8218.7	7416.4	755.6	55	-0.5	0.5	0.1	55
Baseline-950	49	-167726.5	152433.2	1826.7	52	-2.0	2.0	0.1	52
OMR-1500	49	-169763.1	151871.3	1339.6	52	-2.1	2.0	0.0	52
OMR-500	49	-167726.5	152433.2	2173.7	52	-2.0	2.0	0.1	52
Baseline-950	81	-4675.7	2172.6	-1336.9	41	-1.2	0.6	-0.4	41
OMR-1500	81	-4984.0	2359.9	-1618.1	36	-1.2	0.7	-0.4	36
OMR-500	81	-4403.3	3143.2	-886.3	47	-1.2	0.9	-0.2	47
Baseline-950	94	-13302.6	11145.0	-258.6	52	-1.7	1.6	0.0	52
OMR-1500	94	-13547.5	10749.2	-585.7	51	-1.7	1.5	-0.1	51
OMR-500	94	-13302.6	11145.0	-28.6	52	-1.7	1.6	0.0	52
Baseline-950	107	-6216.5	4586.7	-323.1	52	-1.6	1.3	-0.1	52

Scenario (cfs)	DSM2 Channel	Flow Min.	Flow Max.	Flow Mean	Flow % Positive	Velocity Min.	Velocity Max.	Velocity Mean	Velocity % Positive
OMR-1500	107	-6246.4	4494.3	-411.8	51	-1.6	1.3	-0.1	51
OMR-500	107	-6216.5	4586.7	-261.3	52	-1.6	1.3	-0.1	52
Baseline-950	124	-19481.4	13228.6	-2526.0	45	-0.6	0.4	-0.1	45
OMR-1500	124	-19501.3	13076.0	-2694.1	44	-0.6	0.4	-0.1	44
OMR-500	124	-19481.4	13228.6	-2412.5	45	-0.6	0.4	-0.1	45
Baseline-950	148	-8206.4	6605.0	-295.9	51	-0.8	0.7	0.0	51
OMR-1500	148	-8341.0	6398.0	-449.4	51	-0.9	0.7	0.0	51
OMR-500	148	-8206.4	6605.0	-188.3	52	-0.8	0.7	0.0	52
Baseline-950	160	-5201.0	3864.9	-41.6	53	-0.5	0.5	0.0	53
OMR-1500	160	-5263.4	3778.5	-107.4	53	-0.5	0.5	0.0	53
OMR-500	160	-5201.0	3864.9	9.0	53	-0.5	0.5	0.0	53
Baseline-950	434	-178565.1	170195.6	7042.2	53	-1.9	2.0	0.1	53
OMR-1500	434	-178750.4	170031.9	6918.2	53	-1.9	2.0	0.1	53
OMR-500	434	-178565.1	170195.6	7129.1	53	-1.9	2.0	0.1	53




FIGURE A3: Spatial representation of DSM2 modeled flow by channel.

DSM2 model interpretation entrainment in Delta strata regions

North Delta into Interior and Central Delta

Changes in flow and velocity related to the modeled flow conditions would not likely be detected in the main-stem Sacramento River but could be detected in other locations in the North Delta. It's unlikely that listed salmonids would experience changes to rearing, foraging, and /or sheltering; however, it is likely that migrating fish may experience changes. Higher flows should increase travel rate (and decrease travel time) and aid in out-migration.

San Joaquin River and Central Delta into South Delta

Changes in flow and velocity related to the modeled flow conditions would not likely be detected and it's unlikely that listed salmonids would experience changes to rearing, foraging, and /or sheltering. Towards the end of the forecast period, less negative flow may increase travel rate (and decrease travel time) to more northern sites (e.g., Frank's Tract).

South Delta into facilities

Changes in flow and velocity related to the modeled flow conditions would likely be detected and it's likely that listed salmonids would experience changes to rearing, foraging, and /or sheltering. Decrease in flow may result in a decreased travel rate (and longer travel times) towards the pumps (with influence increasing closer towards the pumps)

DSM2 channel locations information




FIGURE A4. Highlighted DSM2 channels by Delta Strata.

TABLE A2. Description of channel location, by Delta Strata region. Not all listed channels have model results presented in every weekly Proposed Action Assessment.

DSM2 Channel	Description
North Delta into Interior and Central Delta	N/A
CHAN049	San Joaquin River at Sherman Island
CHAN310	Three-Mile Slough
CHAN421	Sacramento River at Delta Cross Channel
CHAN422	Sacramento River at Delta Cross Channel
CHAN423	Sacramento River at Delta Cross Channel
CHAN434	Sacramento River at Sherman Island
San Joaquin River and Central Delta into South Delta	N/A
CHAN006	San Joaquin River at Head of Old River (HOR)
CHAN021	San Joaquin River downstream from confluence with Calaveras River
CHAN024	San Joaquin River upstream of Turner Cut
CHAN054	Old River at confluence with San Joaquin River (HOR)
CHAN107	Old River north of Rock Slough
CHAN117	Old River south of Franks Tract
CHAN124	Old River between Franks Tract and San Joaquin River
CHAN160	Columbia Cut
CHAN173	Turner Cut
South Delta into Facilities	N/A
CHAN148	Middle River
CHAN227	Victoria Canal
CHAN081	Grant Line Canal
CHAN094	Old River

Attachment B: Delta Turbidity Report

Department of Water Resources
Division of Operations and Maintenance
SWP Water Operations Office

Delta Turbidity Conditions Report

For conditions through: January 31, 2022

General Conditions:

Inflows:


Freeport	12020 cfs
Yolo Bypass	55 cfs
Vernalis	994 cfs
Cosumnes	298 cfs
Mokelumne	334 cfs
Calaveras	50 cfs

Exports:

Clifton Court	2500 cfs
Jones	3431 cfs

Other:

OMR (Index)	-5075 cfs
QWEST	-2729 cfs
NDOI	5811 cfs



OBI station Turbidity values are CDEC daily data.
All other stations Turbidity values are daily average calculated from CDEC event data.

Figure 2. Delta Daily Turbidity Trends Through 01/31/2022