

Weekly Assessment of CVP and SWP Delta Operations on ESA-listed Species

Executive Summary

a) Operations anticipated during the week

See Weekly Fish and Water Operation Outlook document for May 24 – May 30

b) Winter-run Chinook Salmon summary

Loss of natural winter-run Chinook Salmon (by length at date, LAD) has not occurred in the past week at the fish salvage facilities (WY 2022 total loss = 73.04 fish, as of 5/22/2022). Loss of natural winter-run Chinook Salmon at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities could occur over the next week based on hydrology and recent observations. Loss of hatchery winter-run Chinook salmon has not occurred in the past week at the fish salvage facilities (WY 2022 total loss = 6.71 fish, as of 5/22/2022). 1-5% of juvenile natural winter-run Chinook Salmon from brood year (BY) 2021 are estimated to be present in the Delta. The Delta Cross Channel (DCC) gates closure for the season reduces exposure of winter-run Chinook Salmon juveniles that are present in the Sacramento River near the DCC gates into the interior Delta.

c) Spring-run Chinook salmon summary

Juvenile natural spring-run Chinook Salmon from BY 21 are migrating past the DCC gates. The exposure and effects of DCC closure are unlikely for natural spring-run Chinook Salmon. Length-at-date spring run were observed in the Delta within the previous seven days. Larger, older juveniles were observed that may be yearling spring run. Natural spring-run Chinook salmon have been observed at the fish collection facilities in the past week. 7-19% young of year spring-run Chinook Salmon are estimated to be in the Delta.

d) Central Valley Steelhead summary

Loss of natural Central Valley California (CCV) steelhead has occurred in the past week at the State and Federal fish salvage facilities (WY 2022 total loss December through March = 84.28 fish and WY 2022 total loss April through June = 151.66 fish, as of 5/23/2022). Loss of Central Valley steelhead at the Central Valley Project (CVP) and State Water Project (SWP) fish collection facilities could possibly occur over the next week. 5-20% of juvenile CCV Steelhead are estimated to be present in the Delta. Closure of the DCC gates for the season will reduce exposure to Central Valley steelhead juveniles that are potentially present in the Sacramento River near the DCC gates.

e) Green Sturgeon summary

Loss of green sturgeon has not occurred in the past week at the State and Federal fish salvage facilities (WY 2022 total loss = 0 fish, as of 5/23/2022). Loss of green sturgeon is unlikely to occur over the next week due to their rare presence in the South Delta.

f) Delta Smelt summary

One larval Delta Smelt was detected over a month ago in the OMR Corridor and any fish in this area are at an increased likelihood of entrainment. Six juvenile Delta Smelt from EDSM were collected on 4/22, 5/9, 5/10 in the Sacramento Deep Water Ship Channel. Larval and juvenile Delta Smelt have been detected in the Lower San Joaquin River, the Lower Sacramento River, Old River, the confluence, and the North Delta. Delta Smelt adults and sub-adults are less likely to move into the south and central delta since turbidity remains low. Delta Smelt spawning is likely winding down due to increased water temperatures and timing.

g) Monitoring Teams summary

There were no non-consensus issues to report from the Salmon Monitoring Team.

There were no non-consensus issues to report from the Smelt Monitoring Team.

Operational and Regulatory Conditions

See current Weekly Fish and Water Operation Outlook document.

Biology, Distribution, and Evaluation Winter-run Chinook salmon, Spring-run Chinook salmon, Central Valley Steelhead

POPULATION STATUS

Winter-run Chinook salmon

Delta Life Stages:

Juveniles, Adults

Brood Year 2021 Productivity:

- Natural winter-run Chinook salmon: A draft juvenile production estimate (JPE) calculation has been established for brood year (BY) 2021 winter-run Chinook salmon. The final winter-run Chinook salmon JPE is 125,038 fish. Among other topics, the group discussed the thiamine vitamin deficiency that is being observed again in broodstock at the Livingston Stone NFH similar to last year's observation. Last year the thiamine vitamin deficiency appeared to negatively affect survival of juvenile fish as they migrate downstream towards the Delta. Mean cumulative weekly passage of winter-run Chinook salmon through 5/23/2022 for the last 19 years of passage data is 100.0% (one SD of 0.0%). By 4/22/2022, 572,535 winter-run Chinook salmon were estimated to have passed RBDD compared to the cumulative passage last year of 3,813,580 winter-run Chinook salmon.
- Hatchery winter-run Chinook salmon:
 - Approximately 123,975 Livingston Stone NFH brood year 2021 winter Chinook salmon were released at dusk on 2/9/2022 into the Sacramento River at Caldwell Park in Redding, CA. This is the first release of LSNFH brood year 2021 winter Chinook salmon comprising of approximately 24% of the total hatchery production for the Sacramento River supplementation program. The release group is 100% marked (adipose-fin clip and CWT) with an overall estimated average fork length of 85mm. There has been loss this water year with this release group. Some fish from this group were acoustically tagged and are being tracked (Table 6).
 - Approximately 396,310 LSNFH brood year 2021 winter Chinook salmon were released at dusk on 3/2/2022 into the Sacramento River at Caldwell Park in Redding, CA. This is the second release of LSNFH brood year 2021 winter Chinook salmon. The release group is 100% marked (adipose-fin clip and CWT) with an overall estimated average

fork length of 85mm. There has not been loss this water year with this release group. Some fish from both groups were acoustically tagged and are being tracked (Table 6).

- Approximately 136,958 Coleman NFH Complex brood year 2021 winter Chinook salmon were released 3/16/2022 and 3/17/2022 on the North Fork Battle Creek. The release group is 100% marked (adipose fin-clip and a left pelvic-fin clip and CWT) with an overall estimated average fork length of 85mm.

Spring-run Chinook salmon

Delta Life Stages:

Young-of-year (YOY) and Yearlings

Brood Year 2021 Productivity:

- Natural spring-run Chinook salmon: No JPE has been established for spring-run Chinook salmon. Approximately 99.7% juvenile spring-run sized Chinook salmon have been observed passing Red Bluff Diversion dam as of 5/22 based on historical data.
- Hatchery spring-run Chinook salmon surrogates associated with the Proposed Action (PA 4.10.5.10.2 Additional Real-Time OMR Restrictions and Performance Objectives):
 - Approximately 84,343 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 12/15/2021. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 145mm. This is the first spring-run Chinook salmon surrogates release group associated with the Proposed Action. There has been loss this water year of fish associated with the first surrogate release group.
 - Approximately 82,626 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 12/22/2021. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 125mm. This is the second spring-run Chinook salmon surrogates release group associated with the Proposed Action. There has been loss this water year of fish associated with the second release group.
 - Approximately 77,325 late-fall Chinook salmon from Coleman NFH were released at Battle Creek on 1/6/2022. This group is 100% marked with adipose-fin clip and CWT and have an estimated average fork length of 145mm. This is the third spring-run Chinook salmon surrogates release group associated with the Proposed Action. There

has been loss this water year of fish associated with the third release group.

- An additional Coleman National Fish Hatchery (CNFH) hatchery surrogate group was released on 4/5/2022 and 4/7/2022 which totaled 339,094 fall-run (FR) of which 339,094 FR were marked with an adipose fin clip and CWT.
- Hatchery spring-run Chinook salmon surrogates associated with the ITP (Condition Of Approval 8.6.4):
 - Approximately 2.8 million (2,865,384) fall-run Chinook salmon from Coleman NFH were released into Battle Creek on 3/15/2022. This group is 25% marked (adipose-fin clip and CWT) and 75% unmarked and has an estimated fork length of 75mm. This is the first release from Coleman NFH associated with ITP COA 8.6.4.
 - Approximately 3.0 million (2,982,466) fall-run Chinook salmon from Coleman NFH were released into Battle Creek on 3/18/2022. This group is 25% marked (adipose-fin clip and CWT) and 75% unmarked and has an estimated fork length of 75mm. This is the second release from Coleman NFH associated with ITP COA 8.6.4.
 - Approximately 1,458,758 brood year 2021 spring-run Chinook salmon from Feather River Fish Hatchery will be released into the Feather River at Gridley and Boyd's pump on 3/30/2022. This group is 50% tagged (adipose fin clip and CWT) and has an estimated fork length of 68 mm. This is the first release group from Feather River Fish Hatchery associated with ITP COA 8.6.4.
 - Approximately 4.22 million brood year 2021 fall-run Chinook salmon from Coleman NFH were released between 3/31/2022 and 4/1/2022 at Battle Creek. This group is 25% marked (adipose fin clip and CWT) and 75% unmarked and has an estimated fork length of 75mm. This is the third release group from Coleman NFH associated with ITP COA 8.6.4. There has been loss this water year associated with the third release group.
 - An additional Coleman National Fish Hatchery (CNFH) hatchery surrogate group was released on 4/5/2022 and 4/7/2022 which totaled 339,094 fall-run (FR) of which 339,094 FR were marked (i.e., 100% marked) with an adipose fin clip and CWT. The COA 8.6.4 daily loss threshold for this group is 847.74. The average estimated fork length is 75 mm. This is the fourth release group from Coleman NFH associated with ITP COA 8.6.4. There has been loss this water year associated with the fourth release group.

- Approximately 480,000 brood year 2021 spring-run Chinook Salmon from Feather River Fish Hatchery will be released each day on 5/10/2022 and 5/11/2022 into San Pablo Bay at Mare Island. This group is 75% marked (CWT and adipose fin clip) and has an estimated fork length of 90mm. This final hatchery surrogate release group of SR for COA 8.6.4 from the FRFH will also be released in the Bay on 5/10/2022 and 5/11/2022 and will not meet the needs for the ITP COA 8.6.4.
- San Joaquin River Restoration Program (SJRRP) Salmon Conservation and Research Facility (SCARF) Chinook salmon:
 - Approximately 57,478 brood year 2021 spring-run Chinook salmon from the SJRRP SCARF were released on 2/14/2022 into the San Joaquin River at Fremont Ford Bridge (Highway 140). This release consisted of marked juveniles (adipose fin clip and CWR) and have an estimated average fork length of 85mm.
- The agencies in the SaMT discussed the thiamine vitamin deficiency that was observed in winter run Chinook salmon broodstock at the Livingston Stone NFH in BY 2021. Last year the thiamine deficiency appeared to negatively affect survival of juvenile fish as they migrate downstream towards the Delta. The thiamine deficiency issue is also likely impacting spring-run Chinook salmon. Although the egg take goals have been met at the Feather River Fish Hatchery, they are still experiencing fertility issues that are impacting production.

Central Valley Steelhead

Delta Life Stages:

- Spawning Adults, Kelts, Juveniles

Brood Year 2021 Productivity:

- Spawner abundance: There is limited information about the adult steelhead population. It is estimated to be small, contributing to the limited productivity of the population
- Natural steelhead: No JPE has been established for steelhead. Data are limited.
- Hatchery steelhead: Reclamation's Proposed Action has no hatchery steelhead triggers.
 - Approximately 610,911 steelhead from Coleman NFH were released at Battle Creek from 12/12/2021 to 12/13/2021. This group is 100% marked only (with an adipose-fin clip) and has an estimated average fork length of 195mm.

- Approximately 57,056 steelhead from Mokelumne River Fish Hatchery will be released into the Mokelumne River at Feist Ranch from 1/31/2022 to 2/1/2022. This group is 100% marked (CWT and adipose fin clip) and has an estimated average fork length of approximately 210mm.
- Approximately 424,000 steelhead from Nimbus Fish Hatchery will be released into the Lower American River at the Sunrise Boat Ramp from 1/31/2022 through 2/4/2022. This group is 100% marked (adipose fin clip) and has an average fork length of approximately 240mm.
- Approximately 475,000 steelhead from Feather River Fish Hatchery will be released into the Feather River at Boyd's Pump from 2/4/2022 through 2/11/2022. This group is 100% marked (adipose fin clip) and has an average fork length of approximately 210mm.
- Approximately 60,000 steelhead from Mokelumne River Fish Hatchery will be released into the Mokelumne River at Feist Ranch from 2/28/2022 through 3/1/2022. This group is 100% marked (adipose fin clip) and has an average fork length of approximately 210mm.
- Approximately 50,000 steelhead from Mokelumne River Fish Hatchery will be released into the Mokelumne River at Feist Ranch from 4/4/2022 through 4/5/2022. This group is 100% marked (adipose fin clip and right maxillary clip) and has an average fork length of approximately 230mm.

DISTRIBUTION

Winter-run Chinook Salmon

Current Distribution:

- On 5/24/2022, SaMT estimated 1-5% of juvenile winter-run Chinook salmon were present in the Delta and 95-99% were estimated to have exited the Delta (Table 1). The SaMT estimate is based both on historical trends and observations of recent monitoring. The SaMT expressed some concern that these numbers seem low especially given the precipitation events that occurred early in the Water Year (October and December 2021).
- Combined total natural winter-run Chinook salmon loss equals 73.04 fish (as of 5/22/2022).
- Combined total hatchery winter-run Chinook salmon loss equals 6.71 fish (as of 5/22/2022).
- Fish have been steadily arriving since the beginning of October 2021.

- Winter-run Chinook Salmon have not been observed in RST monitoring locations over the past week.
- Movement of winter-run Chinook Salmon juveniles into the lower reaches of the Sacramento River and upper Delta is continuing. Mill and Deer creeks daily flows were recorded higher than 95 cfs during the past week.

Historic Trends

- Based on historical trends in salvage, 99.9% of winter-run Chinook salmon should have been observed in salvage by this time of the water year (Table 3). Loss of natural winter-run Chinook salmon at the CVP and SWP fish collection facilities is unlikely to occur over the next week based on hydrology and recent observations. If historic trends in salvage were to continue, winter-run Chinook salmon loss is expected to remain the same over the next week.
- Based on historic loss patterns and historic cumulative loss, it was possible that in WY 2022 50% of the wild winter-run Chinook salmon Delta Performance single-year loss threshold (731.47 wild winter-run Chinook salmon, as set by the JPE estimate) could be exceeded (Figure 1, Operations Outlook Table 2a). However, as the season has progressed only a few years of historical loss would cause exceedance of the WY 2022 single-year loss threshold (WY 2011 and WY 2012). As of this point in the season no years of historical loss would cause exceedance of the WY 2022 single-year loss threshold. Members of the SaMT agree the likelihood of exceeding the next annual loss threshold and loss occurring in the next week is unlikely to lead to exceedance of the threshold.

Forecasted Distribution within Central Valley and Delta regions

- SaMT believes that the majority of winter-run Chinook salmon have left the system. The STARS model projects route-specific proportion of entrainment, survival, and travel times (Table 4). This model does not estimate entrainment into the lower Sacramento River sloughs (i.e., Three-Mile Slough). The DCC gates were closed 11/30/21 and will remain closed with the exception of Memorial Day Weekend (May 27 – May 31).
- The entrainment tool estimates a median loss of 0 fish and a maximum loss of 0 fish during this week (SacPAS last updated on 5/23/2022, Table 5, Figure 2).
- Genetic information from salvaged Chinook salmon results collected at Tracy Fish Collection Facility and Skinner through 4/25/2022. The majority of the samples that were classified, by LAD method, as winter-run Chinook salmon were genetically identified as fall-run Chinook salmon. The remaining few were identified as spring-run or late-fall run Chinook salmon. For results refer to Attachment C. A fish observed on 4/16/2022 originally classified as a natural winter-run Chinook salmon has been re-classified based on genetic ID as a natural steelhead. The WY 2022 natural winter-run Chinook salmon and

natural steelhead (April 1 – June 15) loss values have been updated accordingly.

- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.
- For preliminary results of acoustically tagged salmonids currently in the system see Table 6.

Spring-run Chinook salmon

Current Distribution

- On 5/24/2022, SaMT estimated 7-19% of young of year CV spring-run Chinook salmon were present in the Delta (Table 1).
- Spring-run chinook are being observed at monitoring sites (Lower Sacramento, GCID, Sacramento Trawls, and Chipps Island in Suisun Bay & West, Confluence, and Sac River System strata). As of the SaMT meeting, Tisdale data available through 5/13/2022.

Historical Trends

- Based on historical trends in salvage, 97.8% of YOY spring-run Chinook salmon were observed in salvage by this time of the water year (Table 3). If historic trends in salvage were to continue YOY spring-run Chinook salmon loss could possibly occur over the next week.

Forecasted Distribution within Central Valley and Delta regions

- On 5/24/2022 SaMT noted that some juvenile CV YOY spring-run Chinook salmon are still migrating downstream. Yearling CV spring run Chinook salmon are moving out from natal tributaries. Mill and Deer creek flows continue to exceed 95 cfs indicating that yearling spring-run Chinook salmon may begin to move and migrate into the mainstem Sacramento River (Table 4).
- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.
- For preliminary results of acoustically tagged salmonids in the system see Table 6.

Central Valley Steelhead

Current Distribution

- On 5/24/2022 SaMT estimated 5-20% of juvenile CCV steelhead were present in the Delta (Table 1).
- Combined total loss of hatchery steelhead equals 639.11 fish as of 5/22/2022. Previous weeks calculation was erroneous.
- Combined total loss of natural steelhead between December 1 and March 31 equals 84.28 fish.
- Combined total loss of natural steelhead between April 1 and June 15 equals 151.66fish as of 5/23/2022.

Historical Trends

- Based on historical trends in salvage, 100% (December through March) of juvenile CCV steelhead should have been observed in salvage by this time of the water year. Based on historical trends in salvage, 89.7% (April through June) of juvenile CCV steelhead should have been observed in salvage by this time of the water year. If historic trends in salvage were to continue, juvenile CCV steelhead loss could possibly increase over the next week.

Forecasted Distribution within Central Valley and Delta regions

- Natural steelhead were observed in key monitoring locations this past week (GCID and lower Feather River).
- SaMT estimated that 5-20% of the population of CCV steelhead may be present in the Delta at this time. Closure of the DCC gates for the season will reduce exposure and possible entrainment of juvenile CCV steelhead into the interior Delta via the DCC gates.
- The entrainment tool estimates a median loss of fish and a maximum loss of 63 fish during this week (SacPAS last updated on 5/23/2022, Table 5, Figure 2).
- For results of entrainment into Delta strata regions from DSM2 model runs (North Delta into Interior and Central Delta, San Joaquin River and Central Delta into South Delta, and South Delta into fish facilities) refer to Attachment A.

TABLE 1. Salmonid distribution estimates

| Location | Yet to Enter Delta (%) | In the Delta (%) | Exited Delta past Chipps Island (%) |
|---|------------------------|------------------|-------------------------------------|
| Young-of-year (YOY) winter-run Chinook salmon | 0 | 1-5 | 95-99 |
| YOY spring-run Chinook salmon | 1-3 | 7-19 | 80-90 |
| YOY hatchery winter-run Chinook salmon | 0 | 0 | 100 |
| Natural origin steelhead | 1-5 | 5-20 | 75-94 |

TABLE 2. Historic migration and salvage patterns.

| Date (5/22) | Red Bluff Diversion Dam | Tisdale RST | Knights Landing RST | Sac Trawl (Sherwood) Catch Index | Chipps Island Trawl Catch Index | Salvage |
|-------------------------------------|--|--|--|--|--|-------------------------|
| Chinook, Winter-run, Unclipped | 100.0% (100.0%,100.0%) BY: 2012 - 2020 | 100.0% (100.0%,100.0%) BY: 2012 - 2020 | 100.0% (100.0%,100.0%) BY: 2013 - 2020 | 100.0% (100.0%,100.0%) BY: 2012 - 2020 | 99.9% (99.7%,100.1%) BY: 2012 - 2020 | 99.9% (99.7%,100.1%) |
| Chinook, Spring-run, Unclipped | 99.7% (99.5%,99.9%) BY: 2012 - 2020 | 100.0% (100.0%,100.0%) BY: 2012 - 2020 | 100.0% (100.0%,100.0%) BY: 2013 - 2020 | 100.0% (99.9%,100.0%) BY: 2012 - 2020 | 99.6% (99.2%,100.0%) BY: 2012 - 2020 | 97.8% (94.8%,100.9%) |
| Steelhead, Unclipped (April - June) | N/A | N/A | N/A | N/A | N/A | 89.7% (79.5%,99.8%) |

TABLE 3. STARS model output.

| Date (5/22/2022) | DCC | Georgiana Slough | Sacramento River | Sutter and Steamboat | Interior Delta |
|---------------------------|-----|------------------|------------------|-------------------------------|----------------|
| Stock: Late Fall Run | N/A | N/A | N/A | N/A | N/A |
| Proportion of Entrainment | 0 | 0.32 | 0.44 | 0.24 | N/A |
| Survival | N/A | 15% | 47% | 35% | N/A |
| Travel Time | N/A | 19d | 11.8d | 12.3d | N/A |
| Stock: Winter Run | N/A | N/A | N/A | N/A | N/A |
| Proportion of Entrainment | N/A | N/A | 0.57 | 0.14 Sutter 0.15 Steamboat | 0.14 |
| Survival | N/A | N/A | 0% | 0% Sutter 0% Steamboat | 0% |
| Travel Time | N/A | N/A | 6.6d | 6.7d Sutter 6.3d Steamboat | 10.2d |

TABLE 4. Mean daily flow and percent change (Wilkins Slough, Deer Creek, Mill Creek; cfs from CDEC) and temperature and percent change (Knights Landing; °F from RST).

| Date | Mill Creek flow (MLM) | MLM Change | MLM Alert | Deer Creek flow (DCV) | DCV Change | DCV Alert | Wilkins Slough flow (WLK) | Knights Landing temperature (°F) | Alert Triggered |
|-----------|-----------------------|------------|------------|-----------------------|------------|------------|---------------------------|----------------------------------|-----------------|
| 5/22/2022 | 206.0 | -2.3% | Flow>95cfs | 163.8 | -1.8% | Flow>95cfs | 3654.2 | N/A | N/A |
| 5/21/2022 | 210.8 | -7.2% | Flow>95cfs | 166.9 | -0.6% | Flow>95cfs | 3683.2 | N/A | N/A |
| 5/20/2022 | 227.1 | 0.1% | Flow>95cfs | 167.8 | -1.3% | Flow>95cfs | 3786.5 | N/A | N/A |
| 5/19/2022 | 227.0 | 3.7% | Flow>95cfs | 170.1 | -1.4% | Flow>95cfs | 3877.2 | N/A | N/A |
| 5/18/2022 | 218.9 | 0.7% | Flow>95cfs | 172.5 | -1.8% | Flow>95cfs | 3974.6 | N/A | N/A |
| 5/17/2022 | 217.2 | -0.1% | Flow>95cfs | 175.8 | -1.2% | Flow>95cfs | 3929.9 | N/A | N/A |
| 5/16/2022 | 217.5 | 5.3% | Flow>95cfs | 177.8 | -1.9% | Flow>95cfs | 3991.8 | 52.6 | N/A |

TABLE 5. a) WY 2022 loss and salvage predictor data: Predicted weekly loss of winter-run Chinook salmon and steelhead at CVP and SWP facilities. b) Environmental details, current and forecast.

| Week | 33 | 34 |
|---|-----------|-----------|
| a) | Model | N/A |
| Steelhead median | 8 | 0 |
| Steelhead high | 63 | 43 |
| Winter-run Chinook median | 0 | 0 |
| Winter-run Chinook high | 0 | 0 |
| b) | Data | Forecast |
| Temperature (Mallard Island, C) | 17.7 | 18.0 |
| Precipitation (5-d running sum, inches) | 0.0 | 0.0 |
| Old + Middle River flows (cfs) | -1,820 | -1,820 |
| Sacramento River flow (Freeport, cfs) | 7,765 | 7,765 |
| DCC Gates | Open | Closed |
| San Joaquin River flow (Vernalis, cfs) | 673 | 673 |
| Export | 1,215 | 1,215 |

TABLE 6. Preliminary survival (Φ) results for acoustically tagged salmonids (Chinook and steelhead). Individual studies denoted by rows with bold lines.

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|---|------------------|-------------------------|-------------------------|---|---|--|---|
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 1 (200) | 12/15/2021 – 12/16/2021 | RBDD | 75.5% (SE = 3.0) | 17.1% (SE = 2.7) | 22.7% (SE = 3.4) | Fish detected at Old River and Benicia receivers |
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 2 (200) | 2/2/2022 – 2/3/2022 | RBDD | 6.0% (SE = 1.7) | 0.5% (SE = 0.5) | 8.4% (SE = 8.0) | Fish detected at Tower Bridge, below Georgiana, and Benicia receivers |
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 3 (200) | 2/23/2022 – 2/24/2022 | RBDD | 3.5% (SE = 1.3) | 1.6% (SE = 0.9) | 43.1% (SE = 18.8) | Fish detected at Tower Bridge and Georgiana receivers |
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 4 (200) | 3/17/2022 – 3/18/2022 | RBDD | 4.0% (SE = 1.4) | 0.5% (SE = 0.5) | 12.6% (SE = 11.8) | Fish detected at I-80 (3.5%) and Benicia (0.5%) receivers |
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 5 (367) | 4/7/2022 – 4/8/2022 | RBDD | 6.0% (SE = 1.2) | 1.9% (SE = 0.7) | 32.0% (SE = 10.0) | Fish detected at Tower Bridge (6%) and Benicia receivers (2%) |

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|---|-----------------|---------------------|------------------------------------|-----------------------------|---|--|---|
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 6 (361) | 4/27/2022-4/28/2022 | RBDD | 1.7% (SE = 0.7) | 0.3% (SE=0.3) | 16.7% (SE= 15.2) | Fish Detected at Benicia Receivers (0.28) |
| Hatchery-origin Chinook salmon: Seasonal Survival Study | Week 7 (375) | 5/11/2022-5-12-2022 | RBDD | 0.3% (SE=0.3) | 0.0% (SE= NA) | 0.0% (SE= NA) | Fish detected at I80 Bridge (0.27) |
| Hatchery-origin winter-run Chinook salmon | Week 1 (139) | 2/9/2022 | Caldwell Park, Redding | 15.8% (SE = 3.1) | 6.5% (SE = 2.1) | 40.9% (SE = 10.5) | Fish detected at Tower Bridge, I80 Bridge (15.8%), and Benicia (6.5%) receivers |
| Hatchery-origin winter-run Chinook salmon | Week 2 (430) | 3/2/2022 | Caldwell Park, Redding | 12.6% (SE = 1.6) | 5.6% (SE = 1.1) | 44.4% (SE = 6.8) | Fish detected at Tower Bridge, below Georgiana (4.9%), and Benicia (5.6%) receivers |
| Hatchery-origin Battle Creek winter-run Chinook salmon | Release 1 (312) | 3/16/2022 | North Fork Battle Creek, below ECD | 1.0% (SE = 0.6) | 0% | 0% | Fish detected at Tower Bridge (0.6%) and I-80 (0.6%) receivers |
| Hatchery-origin Battle Creek winter-run Chinook salmon | Release 2 (600) | 3/16/2022 | North Fork Battle Creek, below ECD | 0.8% (SE = 0.4) | 0.2% (SE = 0.2) | 20.0% (SE = 17.9) | Fish detected at Tower Bridge (0.7%) and I-80 (0.7%) receivers |

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|---|-------------------------|-----------------------|------------------------------------|-----------------------------|---|--|--|
| Hatchery-origin Battle Creek winter-run Chinook salmon | Release 3 (288) | 3/16/2022 | North Fork Battle Creek, below ECD | 1.4% (SE = 0.7) | 0% | 0% | Fish detected at Tower Bridge (1%) and I-80 (1%) receivers |
| 6-year study San Joaquin River steelhead – March releases | Head of Old River (144) | 3/15/2022 | Head of Old River | Not Enough Detections Yet | 7.6% (SE = 2.2) | N/A | Fish detected at Clifton Court Radial Gates, CVP trash rack, and Benicia receivers |
| 6-year study San Joaquin River steelhead – March releases | Durham Ferry (248) | 3/15/2022 | Durham Ferry | Not Enough Detections Yet | 4.8% (SE = 1.4) | N/A | Fish detected at Clifton Court Intake Canal, CVP trash rack, and Benicia receivers |
| 6-year study San Joaquin River steelhead – March releases | Stockton (100) | 3/15/2022 | Stockton | Not Enough Detections Yet | 12.1% (SE = 3.3) | N/A | Fish detected at Old River Quimby Island and Benicia (6%) receivers |
| 6-year study San Joaquin River steelhead – April releases | Head of Old River (144) | 4/19/2022 – 4/22/2022 | Head of Old River | Not Enough Detections Yet | 4.8% SE = 187 | N/A | Fish detected at Old River, Clifton Court Radial Gates, CVP trash rack, and Middle River receivers |

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|---|--------------------|-----------------------|-----------------------|-----------------------------|---|--|--|
| 6-year study San Joaquin River steelhead – April releases | Durham Ferry (240) | 4/19/2022 – 4/22/2022 | Durham Ferry | Not Enough Detections Yet | 7.9% SE = 1.7 | N/A | Fish detected at CVP trash rack and Clifton Court Radial Gates receivers |
| 6-year study San Joaquin River steelhead – April releases | Stockton (100) | 4/19/2022 – 4/21/2022 | Stockton | Not Enough Detections Yet | 20.0% SE = 4.0 | N/A | Fish detected at Middle River receiver |
| Hatchery-origin San Joaquin spring-run Chinook salmon | Fremont Ford (694) | 3/30/2022 | Fremont Ford | Not Enough Detections Yet | Not Enough Detections Yet | Not Enough Detections Yet | Not Enough Detections Yet |
| Hatchery-origin rice field reared fall-run Chinook salmon | Release 1 (315) | 3/7/2022 | Sutter Bypass | 9.2% (SE = 1.6) | 1% (SE = 0.5) | 10.3% (SE = 5.7) | Fish detected at Below Georgiana and Benicia (1%) receivers |
| Hatchery-origin fall-run Chinook salmon paired release | Upstream (300) | 4/5/2022 | Battle Creek | 6.0% (SE = 1.4) | 2.1% (SE = 0.8) | 34.5% (SE = 11.5) | Fish detected at Butte Bridge receiver (42%) |
| Hatchery-origin fall-run Chinook salmon paired release | Downstream (300) | 4/7/2022 | Sac River, Butte City | 11.3% (SE = 1.8) | 2.4% (SE = 0.9) | 21.3% (SE = 7.2) | Fish detected at Butte Bridge receiver (1%) |

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|--|-----------------|-----------------------|-------------------------|-----------------------------|---|--|--|
| Natural-origin Red Bluff RST captured Chinook salmon | Release 1 (72) | 4/13/2022 – 4/15/2022 | Altube Island | 25% (SE = 5.1) | 5.6% (SE = 2.7%) | 22.2% (SE = 9.8) | Fish detected at Tower Bridge, Sacramento River below Georgiana (6%), and Benicia (1.4%) receivers |
| Natural-origin Red Bluff RST captured Chinook salmon | Release 2 (91) | 4/27/2022 – 4/29/2022 | Altube Island | 0% (SE = 0) | NA | NA | Fish detected at Butte Bridge (3%) |
| Natural-origin Red Bluff RST captured Chinook salmon | Release 3 (68) | 5/4/2022 – 5/5/2022 | Altube Island | 0% (SE = 0) | NA | NA | Fish detected at Butte Bridge (6%) |
| Natural-origin Red Bluff RST captured Chinook salmon | Release 4 (71) | 5/11/2022 – 5/13/2022 | Altube Island | 0% (SE = 0) | NA | NA | Fish detected at Salt Creek (100%) and Butte Bridge (4%) receivers |
| Spring-Run Surrogate Upper Sac River Basin | Release 1 (149) | 4/15/2022 | Red Bluff Diversion Dam | 13.3% (SE = 2.8) | 2.7% (SE = 1.3) | 20.0% (SE = 8.9) | Fish detected at Benicia receiver |
| Spring-Run Surrogate Upper Sac River Basin | Release 2 (150) | 4/20/2022 | Red Bluff Diversion Dam | 7.3% (SE = 2.1) | 0% | 0% | Fish detected at Tower Bridge receiver |
| Spring-Run Surrogate Upper Sac River Basin | Release 3 (148) | 4/28/2022 | Red Bluff Diversion Dam | 0% | 0% | 0% | Fish detected at Meridian Bridge receiver |

| Project | Group (n) | Release Date(s) | Release Location | Min. Φ to Tower Bridge | Min. Φ to Benicia Bridge (East Span) | Min. through-Delta Φ (Sac to Benicia) | Notes |
|--|------------------|------------------------|-------------------------|---|---|--|---|
| Spring-Run Surrogate Upper Sac River Basin | Release 4 (152) | 5/5/2022 | Red Bluff Diversion Dam | 0% | 0% | 0% | Fish detected at Meridian Bridge receiver |

Table updated 5/23/2022. Additional information for all projects can be found online at

<https://oceanview.pfeg.noaa.gov/CalFishTrack/>




FIGURE 1. WY 2022 cumulative natural winter-run Chinook salmon loss (blue) and 2009 – 2018 historic cumulative loss (gray, different symbols). Historic daily mean plotted in black circles.




FIGURE 2. Predicted weekly loss of steelhead and winter-run Chinook salmon at the CVP and SWP facilities

EVALUATION

1. **After January 1, are more than 5% of juveniles from one or more salmonid species present in the Delta?**

Greater than 5% of juvenile winter-run Chinook salmon, CCV steelhead, and young of year spring-run Chinook salmon are present in the Delta.

2. **Does the operational outlook's ranges impact fish movement and change the potential distribution of fish?**

Potential effects within the 7 days (near-term) in the operations outlook. OMR flow is expected to remain at or more positive than -5,000 cfs this upcoming week.

Potential effects longer than the 7 days (longer-term) in the operations outlook. See response above to (2) (i).

3. **What is the likelihood of increased loss exceeding the next annual loss threshold (50%, 75% or 90% of threshold) resulting in OMR management actions based on population distribution, abundance, and behavior of fish in the Delta?**

Winter-run Chinook salmon

Total juvenile natural winter-run Chinook salmon (LAD) loss is 73.04 fish (as of 5/22/2022). Loss of juvenile winter-run Chinook salmon has not occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believe that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold set in the Proposed Action. SaMT anticipates potential loss occurring in the next week leading to exceedance of ITP COA 8.6.1 (winter run single year loss threshold) and ITP COA 8.6.3 (mid and late season natural winter-run daily loss threshold).

Total hatchery winter-run Chinook salmon loss is 6.71 fish (as of 5/22/2022). Loss of hatchery winter-run Chinook salmon has not occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believe that loss is unlikely to occur in the next week and is unlikely to lead to exceedance of the 50% loss threshold set in the Proposed Action.

Spring-run Chinook salmon

Total juvenile spring-run Chinook salmon (LAD) loss is 552.69 fish (as of 5/22/2022). Loss of juvenile spring-run Chinook salmon has occurred in the past week at the CVP and SWP fish salvage facilities. The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believes that loss occurring in the next

week is unlikely to lead to exceedance of the 50% single-year loss threshold set in the Proposed Action. SaMT does not anticipate loss occurring in the next week leading to exceedance of ITP COA 8.6.4 (daily spring run Chinook hatchery surrogate loss threshold).

Central Valley Steelhead

Total natural juvenile steelhead loss (December 1 through March 31) is 84.28 fish. Total natural juvenile steelhead loss (April 1 through June 15) is 151.66 fish (as of 5/22/2022). Loss of natural juvenile has occurred in the past week at the CVP and SWP fish salvage facilities. Total clipped steelhead loss is 639.11 fish (as of 5/22/2022). The agencies in the SaMT assessed the likelihood of exceeding the next annual loss threshold and believe that loss occurring in the next week is unlikely to lead to exceedance of the 50% single-year loss threshold set in the Proposed Action.

4. If an annual loss threshold has been exceeded, do continued OMR restrictions benefit fish movement and survival based on real-time information?

Winter-run Chinook salmon

The ITP COA 8.6.1 (winter run single year loss threshold) and ITP COA 8.6.3 (mid and late season natural winter-run daily loss threshold) has been exceeded in WY 2022. The annual loss threshold set in the Proposed Action for winter-run Chinook salmon has not been exceeded in WY 2022.

Spring-run Chinook salmon

The ITP COA 8.6.4 (daily spring run Chinook hatchery surrogate loss threshold) has not been exceeded in WY 2022. The annual loss threshold set in the Proposed Action for spring-run Chinook salmon has not been exceeded in WY 2022.

Central Valley Steelhead

The annual loss threshold set in the Proposed Action for steelhead (December 1 – March 31) has not been exceeded in WY 2022.

5. If OMR is more negative than -5,000 cfs are there changes in spawning, rearing, foraging, sheltering, or migration behavior beyond those anticipated to occur under OMR management at -5,000 cfs?

OMR flows are not expected to be more negative than -5,000 cfs during the next week.

Biology Distribution and Evaluation of Green Sturgeon POPULATION STATUS

Delta Life Stages:

- Adults and Juveniles

Juvenile Abundance:

- In 2021, 1037 larval green sturgeon and 5 juvenile green sturgeon were observed at the Red Bluff Diversion Dam during continuous fish monitoring using RSTs in the upper Sacramento River. 14 juveniles were captured and implanted with micro-acoustic tags during the month of October, and all were presumed to leave the upper Sacramento River during the unprecedented storm /run-off event that occurred on October 24, 2021 that resulted in flows at Bend Bridge of ~37,000 cfs.

DISTRIBUTION

Current Distribution

- Adults: Most abundant during spring spawning migration period of March through May, and post spawning out-migration periods May through June; October through January depending on first winter storm event resulting in significant Sacramento River flow increases. Adult presence year-round to a lesser extent mainly in San Pablo Bay. Recent sporadic occurrences of adult green sturgeon in the San Joaquin River system but spawning has not been documented. Unknown if spawning occurred historically in the San Joaquin River system. ARIS surveys confirmed adult sturgeon present in the Feather River.
- Juveniles: Age-1 through Age-3 juveniles present year-round and widely distributed. Juveniles tagged with acoustic tags in the main channel Sacramento River near Sherman Island detected in the Sacramento River as far upstream as the Cache Slough complex, in the San Joaquin River at the Antioch Bridge, in Threemile, Horseshoe Bend, and Montezuma Sloughs. Seasonal abundance at the primary sampling site (near Sherman Island) appears to be highest during summer in based on capture and telemetry data. Residence time at the primary sampling site for individual fish ranges from one day to over one year but telemetry data show outmigration from the primary sampling site to the Pacific Ocean ranges from 27 to 552 days. Recent capture data shows diurnal depth preference in the main channel of the Sacramento River. No recent documentation of shallow water habitat presence or foraging but likely. The remains of a juvenile/young adult green sturgeon were collected at CVP on 1/6/22; however, this does not count towards green sturgeon salvage totals.

Historical Trends

- Juvenile and adult green sturgeon are historically present in the San Joaquin and Sacramento rivers and Delta

Forecasted Distribution within Central Valley and Delta regions

- Juvenile and adult green sturgeon are present in the San Joaquin and Sacramento rivers and Delta during the next week.

EVALUATION

1. Is there likely to be salvage that may exceed the annual loss limit?

Green sturgeon salvage is 0 fish (as of 5/22/2022). The agencies in the SaMT assessed the likelihood of salvage occurring in the next week is unlikely to occur.

Biology, Distribution, and Evaluation of Delta Smelt POPULATION STATUS

Delta Smelt Life Stages:

Juveniles, Larvae and Adults

Brood Year 2021:

Abundance estimate:

The last Delta Smelt All Strata abundance estimate for post-larval/juveniles was 13,086 for data collected between 5/9/2022 – 5/12/2022. The most recent detections of Delta Smelt were on 5/10/2022 caught in the Sacramento River Deepwater Ship Channel.

Biological Conditions:

Juvenile Delta Smelt are present in the Sacramento Deep Water Ship Channel, and Lower Sacramento River stratum based on the survey detections with the previous two weeks. Larval Delta Smelt have been detected in the lower San Joaquin River stratum (station 815, 902), Lower Sacramento (704, 706, 707) and the north Delta. Delta water temperatures have reached the range in which Delta Smelt spawning occurs and larvae have been observed. Due to increased temperatures in the Delta, it is believed spawning has begun to decline for the season. (Damon et al 2016). The Smelt Monitoring Team discussed the most recent monitoring data (Table 4) and considered professional opinion on the historical trends in regional distribution.

DISTRIBUTION

Current Distribution

- Real time detection data is currently limited to EDSM sampling, 20-mm, Chipps Island, Bay Study, and LEPS.
- Since there are only a few recent detections of larval Delta Smelt, the Smelt Monitoring Team’s capacity to estimate where they are within the Delta is limited.
- The last Delta Smelt detection was on 5/10/2022 in Sacramento Deepwater Ship Channel.
- Larval sampling at the Skinner Fish Facility (SFF) and the Tracy Fish Collection Facility (TFCF) began on 2/7/2021.
- COA 8.5.2: No larval or juvenile Delta Smelt have been salvaged at the SFF or TFCF as of 5/23/2022 (Table 7).

TABLE 7. Summary of newly reported detections of Delta Smelt by Region and Salvage Facilities on 4/22/2022, 5/9/2022 and 5/10/2022. Regions are those defined by EDSM sampling. Delta Smelt >58mm FL are considered adults. Subadult fish are considered by the SMT to be fish from the previous year’s cohort based on size and timing of collection. Young of year are considered juveniles and larvae.

| Life Stage | North | South | West | Far West | Salvage |
|-----------------|-------|-------|------|----------|---------|
| Adult | 0 | 0 | 0 | 0 | 0 |
| Subadult | 0 | 0 | 0 | 0 | 0 |
| Larvae/Juvenile | 6 | 0 | 0 | 0 | 0 |

TABLE 8. Summary of recent Delta Smelt detections reported since last assessment and the total detections for the current water year. Notes reflect latest information on reported detections or completion of survey for the water year and include both larval and adult detections. Total Fish counts do not distinguish between hatchery origin and wild Delta Smelt. Table may include preliminary data that may not have received full QA/QC, but any corrections will be made the following week.

| Sampling Method | Frequency | New Detections | WY2022 | Notes |
|----------------------|-----------|----------------|--------|--|
| EDSM | Weekly | 6 | 72 | Phase 2 Ongoing Phase 3 Starts 7/5/22 |

| Sampling Method | Frequency | New Detections | WY2022 | Notes |
|------------------------------|--------------|-------------------|--------|--|
| SKT | Monthly | 0 | 18 | SKT#5 complete. Last SKT for WY2022 |
| SLS | Biweekly | 0 | 1 | Complete |
| 20-mm | Biweekly | 0 | 10 | Survey 2: Processing Survey 3: Processing Survey 4: Processing Survey 5: On water this week |
| Summer Townet | Biweekly | 0 | 0 | Complete |
| Bay Study | Monthly | 0 | 1 | Ongoing |
| FMWT | Monthly | 0 | 0 | Complete |
| Chipps Island Trawl | Weekly | 0 | 2 | Ongoing (3-days a week) |
| FCCL Brood Stock Collections | Weekly | 0 | 0 | Ended |
| LEPS | As available | 0 | 0 | Ongoing (20mm gear) |
| TFCF | Daily | 0 | 1 | Ongoing |
| FRP | Daily | 0 | 0 | Ongoing |
| Total | N/A | N/A | 105 | Sum of all Delta Smelt observed during the OMR Management Season |

Cultured Delta Smelt Experimental Releases

- On 12/14/2021-12/15/2021, the Experimental Release Technical Team trucked approximately ~12800 brood year 2021 Delta Smelt from the FCCL to Sacramento River at Rio Vista for release into the Delta. This release will include ~89% Adipose fin clipped individuals and ~11% Visible Implant Elastomer (VIE) tagged individuals.
- On 01/11/2022-01/12/2022, the Experimental Release Technical Team will truck approximately ~12,800 brood year 2021 Delta Smelt from the FCCL to Sacramento River at Rio Vista for release into the Delta. This release will include ~97% adipose fin clipped individuals and ~3% Visible Implant Elastomer tagged individuals.
- On 2/3/2022, the Experimental Release Technical Team trucked approximately ~6,400 brood year 2021 Delta Smelt from the FCCL to the Sacramento Deepwater Ship Channel for release into the Delta. This release

will include ~56% adipose fin clipped individuals and ~44% Visible Implant Elastomer (VIE) tagged individuals.

- On 2/9/2022 – 2/10/2022, the Experimental Release Technical Team trucked approximately ~12,800 brood year 2021 Delta Smelt from the FCCL to the Suisun Marsh for release into Montezuma Slough at Belden’s Landing. This release will include ~99.5% adipose fin clipped individuals and ~0.5% Visible Implant Elastomer (VIE) tagged individuals.
- On 02/16/2022-02/17/2022, the Experimental Release Technical Team will truck approximately ~10933 brood year 2021 Delta Smelt from the FCCL to Sacramento Deepwater Ship Channel near channel marker 53 for release into the Delta. This release will include ~100% Adipose fin clipped individuals and ~0% Visible Implant Elastomer tagged individuals.
- Genetic analysis was completed on the two potentially wild Delta Smelt and they were determined to be of hatchery origin with poor quality fin clips. A third smelt collect by the FCCL was identified as a hybrid and additional analysis is being run to determine parentage.
- The unmarked DSM caught by EDSM on 1/5/2022, in the Lower Sacramento River stratum did not assign to a known pair cross of cultured DSM by the GVL, so will be considered as being of wild origin.

TABLE 9. Weekly summary of the origin of Delta Smelt. These identifications are considered tentative and additional genetic testing will confirm the identity of individuals. Individuals with no tags are provided alive to the FCCL as potential additions to the FCCL Broodstock.

| Date | Survey | Stratum/Station | Total Caught | Ad. Clipped | VIE | No Tag |
|-----------|--------|---------------------|--------------|-------------|-----|--------|
| 4/22/2022 | EDSM | Sac DW Ship Channel | 1 | 0 | 0 | 1 |
| 5/9/2022 | EDSM | Sac DW Ship Channel | 1 | 0 | 0 | 1 |
| 5/10/2022 | EDSM | Sac DW Ship Channel | 4 | 0 | 0 | 1 |

Historical Trends

- Wild Delta Smelt detections in the Sacramento Deep Water Ship Channel indicate presence upstream of the confluence, but the fish may be freshwater residents and not representative of the migratory life history patterns in Delta Smelt (Hobbs 2019).

- Historically, the highest peak in salvage is in May and the second highest is in June (Grimaldo et al 2009; figure 5).

Forecasted Distribution within Central Valley and Delta regions

- Predicting the distribution of larval Delta Smelt is currently difficult because detection data is limited to a few individuals and historic patterns may not be representative of the low population levels.
- The SMT uses turbidity as a surrogate for Delta Smelt presence and in making assessments of the likelihood of entrainment for larval Delta Smelt after spawning begins.
- The potential of experimentally released Delta Smelt to distribute from their release site is unknown at this time and SMT cannot predict their distribution beyond the original release site and subsequent recaptures.

ABIOTIC CONDITIONS

Turbidity

- As of 5/23/2022 turbidity continues to be less than 12 FNU at Old River at Bacon Island (OBI) (See attachment B).
- Integrated Early Winter Pulse Protection ended on 1/2/2021.
- Precipitation is not expected in the next seven days.
- Three Station Daily Average water temperature on 5/23/2022 was 20.9° C. This temperature is above the range of Delta Smelt spawning; 9-18 ° C (Damon et al 2016) therefore spawning is likely winding down for the season.

TABLE 10. Relevant Environmental Factors to the current management actions for Delta Smelt.

| Date Reported | OBI Daily Turbidity (FNU) |
|----------------------|----------------------------------|
| 5/23/2022 | 3.06 |

X2 Conditions

- X2 is estimated to be at 90 km.
- When X2 is above 81 km, the SMT uses the X2_EC_Graph.xlsx tool to estimate the position of X2 for both the Sacramento and San Joaquin Rivers and assumes the average of the two is representative of an approximate X2 position.

Other Environmental Conditions

- The Fish and Water Operation Outlook OMR Index values are expected to range between -1,500 to -1,800 cfs from 5/17/2022 to 5/31/2022
- Real time tracking of environmental conditions, relevant thresholds and Delta Smelt catch data are updated daily at:
http://www.cbr.washington.edu/sacramento/workgroups/delta_smelt.html
- Clifton Court Forebay Temperature: 21.81° C

EVALUATION

1. Between December 1 and January 31, has any first flush condition been exceeded?

The running 3-day average flows and running 3-day average turbidity at Freeport exceeded the triggers for Integrated Early Winter Pulse Protection on 12/17/2021 (running 3-day averages: 27,152 cfs and 67 FNU). The CVP and SWP reduced exports from 12/20/2021 until 1/2/2022.

2. Do DSM have a high risk of migration and dispersal into areas at high risk of future entrainment? (December 1- January 31)

Based on recent detection data of experimentally released fish, Delta Smelt are unlikely to be prevalent in the South Delta. Delta Smelt are expected to have made their migration in response to first flush and temperatures are conducive to spawning. Delta Smelt are unlikely to move into the south and central delta since turbidity remains low.

3. Has a spent female been collected?

No spent female has been caught by SKT as of 5/10/2022. Two ripe females were observed in SKT 3. Two prespawn males were observed by SKT 4. SKT surveys have been completed for WY2022.

4. If OMR of -2000 cfs does not reduce OBI turbidity below 12NTU/FNU, what OMR target is deemed protective between -2000 and -5000 cfs?

OBI turbidity is currently below 12 FNU. Turbidity is expected to remain stable in the south delta.

5. If OBI is 12 NTU/FNU, what do other station locations show?

OBI turbidity is currently below 12 FNU. The daily average turbidities on 5/23/2022 at Prisoners Point (5.74 NTU), Holland Cut (5.98 FNU) and Victoria Canal (2.60 NTU), and are expected to remain stable over the next seven days.

6. If OBI is 12 NTU/FNU, is a turbidity bridge avoidance action not warranted? What is the supporting information?

The turbidity at OBI is below 12 FNU, and a turbidity bridge avoidance action ended on April 1st.

7. After March 15 and if QWEST is negative, are larval or juvenile DSM within the entrainment zone of the CVP and SWP pumps based on surveys?

As of 5/23/2022, QWEST is hovering around -400. However, the DCC are planned to be open 5/27/2022 – 5/31/2022. and the opening of the DCC has the potential to swing QWEST to 2,000. A larval Delta Smelt was last observed in the OMR corridor on 4/5/2022. Delta Smelt observed at station 902 have an increased likelihood of entrainment.

8. Based on real-time spatial distribution of Delta Smelt and currently available turbidity information, should OMR be managed to no more negative than -3,500?

Delta Smelt presence was last detected on 4/5/2022 in the OMR Corridor. Turbidity in the South Delta remains low across most stations (See Attachment B) and is expected to remain stable in the next seven days. The OMR index range is between -1,500 to -1,800 cfs for the next seven days. If there are still Delta Smelt in the OMR Corridor, they have an increased likelihood of entrainment.

9. What do hydrodynamic models, informed by EDSM or other relevant data, suggest the estimated percentage of larval and juvenile DSM that could be entrained may be?

The most recent detections of larval Delta Smelt in the lower San Joaquin River and the OMR Corridor were at station 815 (3/22) and 902 (4/5). Most Delta Smelt detections are limited to downstream locations in the San Joaquin and PTM results of nearby stations from 3/29/2022 along with the OMR index over the next seven days support the likelihood of entraining larval Delta Smelt into the OMR corridor remains low. Delta Smelt previously observed in the OMR corridor have an increased likelihood of entrainment and may continue to be present even if not observed by 20 mm surveys or detected by qualitative larval sampling.

DELTA SMELT REFERENCES

- Damon, L. J., S. B. Slater, R. D. Baxter, and R. W. Fujimura. 2016. Fecundity and reproductive potential of wild female Delta smelt in the upper San Francisco Estuary, California. *California Fish and Game* 102(4):188–210.
- Hobbs, J. A., Lewis, L. S., Willmes, M., Denney, C., & Bush, E. (2019). Complex life histories discovered in a critically endangered fish. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-52273-8>

- Lenny F. Grimaldo, Ted Sommer, Nick Van Ark, Gardner Jones, Erika Holland, Peter B. Moyle, Bruce Herbold & Pete Smith (2009) Factors Affecting Fish Entrainment into Massive Water Diversions in a Tidal Freshwater Estuary: Can Fish Losses be Managed? *North American Journal of Fisheries Management*, 29:5, 1253-1270, DOI: 10.1577/M08-062.1
- Polansky, L., Newman, K.B., Nobriga, M.L. et al. Spatiotemporal Models of an Estuarine Fish Species to Identify Patterns and Factors Impacting Their Distribution and Abundance. *Estuaries and Coasts* 41, 572–581 (2018).
<https://doi.org/10.1007/s12237-017-0277-3>
- Sommer, T., F. Mejia, M. Nobriga, and L. Grimaldo. 2011. The Spawning Migration of Delta Smelt in the Upper San Francisco Estuary. *San Francisco Estuary and Watershed Science* 9(2).

Attachment A

Objective

Background

DSM2 model results: summary tables

DSM2 model results: figures

DSM2 model interpretation entrainment into Delta strata regions

DSM2 channel locations information

Objective

Weekly modeling efforts are conducted to examine the effects of varying OMR conditions on the behavior of salmonids present in the Delta in a one-week “look ahead” or outlook. Members of the Salmon Monitoring Team (SaMT) use DSM2 modeling results to help answer how changing pumping regimes translates to differences in flows and velocities modeled at various channel locations within the Delta and what impact modeled environmental parameters may have on rearing, foraging, migrating, and holding salmonids.

Each series of runs consists of up to three OMR conditions: minimum and maximum scenarios bounded by expected OMR index values for that week (Ops Outlook, Table 1) and a baseline which represents an anticipated operational value. Assumptions are made to best estimate future hydrologic characteristics. These inputs are more confident for the future one, two, and three-day timeframes; days four through six have lower confidence. Model scenarios hold hydrology inputs between runs constant and adjust Delta export pumping rates to compare between scenario OMR index values (unless otherwise noted). Although hydrologic ensembles could be used, a single value or deterministic projection is used for efficiency.

SaMT members use weekly DSM2 model results from a range of scenarios as part of a suite of tools to help assess distribution and changes to behavior of salmonids. At each channel location over a six-day action period, environmental parameters are examined: modeled flow and velocity general statistics (e.g., magnitude, range, percent positive), differences in modeled flow and velocity values compared with the baseline scenario, etc. That information, in conjunction with channel location (e.g., close to the Delta pumping facilities, closer to areas with higher tidal influence, etc.) and other environmental considerations (e.g., tidal cycle, upcoming storms, etc.), is then interpreted from a biological perspective. The SaMT recognizes that interpretation of DSM2 model run results with respect to juvenile salmonids (non-neutrally buoyant particles) involves making some assumptions. SaMT explores the possible effects to salmonids of changing OMR index scenarios, assuming each of those potential operations could be that week’s controlling factor.

Background

| Category | Notes |
|---|--|
| <i>Process</i> | N/A |
| Weekly process | <ul style="list-style-type: none"> • DSM2 model runs use a historic and forecasted hydrological input dataset with no assumptions provided by DWR Thursdays and updated by Reclamation Mondays. • Input File updated Monday after initial distribution from DWR for removal of forecasted in lieu of historic input. • Reclamation provides scenarios based on expected OMR index values for the upcoming week. • DSM2 model runs produced Monday and distributed to SaMT members. |
| <i>Hydraulic footprint information</i> | N/A |
| Updated | 5/23/2022 (baseline and scenarios) |
| DSM2 modeling results range | 5/23/2022 – 5/29/2022 |
| <i>OMR index value scenarios</i> | N/A |
| Baseline | -1,700 cfs |
| Scenario 1 | NA |
| Scenario 2 | NA |
| <i>Changes between scenarios</i> | |
| Hydrology | No (see special considerations section below) |
| Delta Exports | Yes |
| <i>Common assumptions</i> | N/A |
| DSM2 run results based on the following assumptions | <ul style="list-style-type: none"> • CCFB Gates are operating to Priority 1 through the end of the forecast period. • The Delta Cross Channel gates were closed on November 30 and will remain closed through the end of the forecast period. • Suisun Marsh salinity control flashboards are in, and 2 of the Suisun Marsh salinity control gates are in open position from March 8, and the remaining 1 gate is closed for maintenance. • See Figure A1a for Sacramento River flow at Freeport during the forecast period. • See Figure A1b for San Joaquin River flow at Vernalis during the forecast period. • Clifton Court Forebay and Tracy Pumping Plant pumping is shown for model runs in Figure A2. |

| Category | Notes |
|---|---|
| <i>Additional information</i> | N/A |
| Considerations for current DSM2 model run | <ul style="list-style-type: none"> • The Base case is constrained by D1641 TUCO (4/4) meeting the X2 requirement via outflow and the IOP for April and May under ITP section 8.17 (Vernalis 1:1) and bound by the minimum combined export of 1,500 cfs. • Likely to be in this "configuration" through May 2022. |
| Caveats | <ul style="list-style-type: none"> • Time-step: DSM2 generates results at 15-minute time-steps. Visualizations of DSM2 model run results are aggregated over daily timesteps. Operations function on a more granular scale than daily time-step. <ul style="list-style-type: none"> ▪ Salmonid behavior: DSM2 provides flow fields which salmonids may encounter but salmonids are not neutrally buoyant particles. Models which incorporate behavior from acoustic tagged salmonids are being developed for South Delta (ePTM, ecoPTM). |

DSM2 model results: figures




FIGURE A1. Daily forecasted Freeport and Vernalis flows (cfs).




FIGURE A2. Daily forecasted pumping: Jones and Banks (cfs) for Baseline.

DSM2 model results: summary tables

TABLE A1. Summary of minimum, maximum, mean, and percent positive flows (cfs) and velocities (ft/s) by DSM2 channel for OMR scenarios over a 6-day time period. For scenario values refer to "Hydraulic Footprint Information" in the conditions / assumptions section above.

| Scenario (cfs) | DSM2 Channel | Flow Min. | Flow Max. | Flow Mean | Flow % Positive | Velocity Min. | Velocity Max. | Velocity Mean | Velocity % Positive |
|----------------|--------------|------------|-----------|-----------|-----------------|---------------|---------------|---------------|---------------------|
| Baseline-1600 | 6 | -932.89 | 1735.85 | 747.15 | 78.9 | -0.41 | 1.03 | 0.45 | 78.9 |
| Baseline-1600 | 21 | -7401.58 | 7101.47 | 250.33 | 53.19 | -0.5 | 0.48 | 0.02 | 53.19 |
| Baseline-1600 | 49 | -157342.56 | 148363.42 | -404.17 | 50.97 | -2 | 1.95 | 0.02 | 50.97 |
| Baseline-1600 | 81 | -4258.03 | 2621.63 | -1001.58 | 44.28 | -1.06 | 0.68 | -0.26 | 44.28 |
| Baseline-1600 | 94 | -13495.9 | 10452.89 | -800.79 | 49.48 | -1.75 | 1.49 | -0.07 | 49.48 |
| Baseline-1600 | 107 | -6321.7 | 4347.25 | -478.93 | 49.03 | -1.68 | 1.25 | -0.11 | 49.03 |
| Baseline-1600 | 124 | -19171.73 | 13098.51 | -2419.4 | 43.54 | -0.62 | 0.43 | -0.07 | 43.54 |
| Baseline-1600 | 148 | -8308.05 | 6204.77 | -534.33 | 49.63 | -0.88 | 0.69 | -0.04 | 49.63 |
| Baseline-1600 | 160 | -4994.31 | 3781.46 | -104.31 | 51.56 | -0.51 | 0.47 | 0.01 | 51.56 |
| Baseline-1600 | 434 | -168688.55 | 163984.44 | 2376.28 | 51.71 | -1.86 | 1.92 | 0.07 | 51.71 |




FIGURE A3: Spatial representation of DSM2 modeled flow by channel.




FIGURE A4: Spatial representation of DSM2 modeled velocity by channel.

DSM2 model interpretation entrainment in Delta strata regions, using modeled velocity results

North Delta into Interior and Central Delta

Channels 49 (San Joaquin River at Sherman Island) and 434 (Sacramento River at Sherman Island)

This week there was a single scenario modeled. There is no change to fish behavior with a lack of a 2nd (or 3rd) model to compare a baseline scenario to. Thus, the SaMT agreed discussing changes to fish behavior due to modeled conditions was not appropriate.

San Joaquin River and Central Delta into South Delta

Channels 6 (San Joaquin River at Head of Old River) and 21 (San Joaquin River upstream of Turner Cut).

This week there was a single scenario modeled. There is no change to fish behavior with a lack of a 2nd (or 3rd) model to compare a baseline scenario to. Thus, the SaMT agreed discussing changes to fish behavior due to modeled conditions was not appropriate.

South Delta into facilities

Channels 148 (Middle River) and 94 (Old River)

This week there was a single scenario modeled. There is no change to fish behavior with a lack of a 2nd (or 3rd) model to compare a baseline scenario to. Thus, the SaMT agreed discussing changes to fish behavior due to modeled conditions was not appropriate.

DSM2 channel locations information




FIGURE A5. Highlighted DSM2 channels by Delta Strata.



FIGURE A6. Aerial imagery of several DSM2 channels discussed in Proposed Action Assessment. North Delta (channels 434 and 49, top panel), San Joaquin River and Central Delta (channels 6 and 21, bottom panel on right), and South Delta (channels 94 and 148, bottom panel on left). Scale for all channels held constant.

TABLE A2. Description of channel location, by Delta Strata region. Not all listed channels have model results presented in every weekly Proposed Action Assessment.

| DSM2 Channel | Description |
|--|---|
| North Delta into Interior and Central Delta | N/A |
| CHAN049 | San Joaquin River at Sherman Island |
| CHAN310 | Three-Mile Slough |
| CHAN421 | Sacramento River at Delta Cross Channel |
| CHAN422 | Sacramento River at Delta Cross Channel |
| CHAN423 | Sacramento River at Delta Cross Channel |
| CHAN434 | Sacramento River at Sherman Island |
| San Joaquin River and Central Delta into South Delta | N/A |
| CHAN006 | San Joaquin River at Head of Old River (HOR) |
| CHAN021 | San Joaquin River downstream from confluence with Calaveras River |
| CHAN024 | San Joaquin River upstream of Turner Cut |
| CHAN054 | Old River at confluence with San Joaquin River (HOR) |
| CHAN107 | Old River north of Rock Slough |
| CHAN117 | Old River south of Franks Tract |
| CHAN124 | Old River between Franks Tract and San Joaquin River |
| CHAN160 | Columbia Cut |
| CHAN173 | Turner Cut |
| South Delta into Facilities | N/A |
| CHAN148 | Middle River |
| CHAN227 | Victoria Canal |
| CHAN081 | Grant Line Canal |
| CHAN094 | Old River |

Attachment B: Delta Turbidity Report

Department of Water Resources
Division of Operations and Maintenance
SWP Water Operations Office

Delta Turbidity Conditions Report

For conditions through: May 23, 2022

General Conditions:

Inflows:

| | |
|-------------|----------|
| Freeport | 7518 cfs |
| Yolo Bypass | 8 cfs |
| Vernalis | 576 cfs |
| Cosumnes | 162 cfs |
| Mokelumne | 161 cfs |
| Calaveras | 162 cfs |

Exports:

| | |
|---------------|---------|
| Clifton Court | 592 cfs |
| Jones | 901 cfs |

Other:

| | |
|-------------|-----------|
| OMR (Index) | -1706 cfs |
| QWEST | -327 cfs |
| NDOI | 4582 cfs |




FIGURE 2. Delta Daily Turbidity Trends Through 05/23/2022

Attachment C.

Genetic information from salvaged Chinook salmon results collected at Tracy Fish Collection Facility and Skinner through 4/25/2022. Underlined text in column headers (“CV ...”) represents aggregate of probability over four Chinook salmon runs for each sample.

| Sample Date | FL* | JD ** | ots28 | Sex | Assignment | Pos Prob1 | Grp | Pos Prob2 | <u>CV fall</u> | <u>CV late-fall</u> | <u>CV spring</u> | <u>CV winter</u> | Model | Facility |
|---------------------|-----|-------|-------|-----|------------|-----------|--------|-----------|----------------|---------------------|------------------|------------------|-----------|----------|
| 12/9/2021 14:00 | 233 | 162 | late | M | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Fall | CVP |
| 12/19/2021 12:00 | 132 | 172 | Late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Winter | CVP |
| 12/24/2021 7:00 | 153 | 177 | late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | CVP |
| 12/25/2021 12:00 | 158 | 178 | late | M | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | CVP |
| 12/25/2021 12:00 | 163 | 178 | late | M | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | CVP |
| 12/25/2021 12:00 | 130 | 178 | late | M | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Winter | CVP |
| 12/27/2021 6:00 | 180 | 180 | late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | CVP |
| 12/30/2021 14:00 | 185 | 183 | late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | SWP |
| 12/31/2021 22:00 | 128 | 184 | late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Winter | CVP |
| 1/1/2022 6:00 | 174 | 185 | late | M | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Late Fall | CVP |
| 1/7/2022 16:00 | 123 | 191 | late | F | Non-winter | 1.000 | Fall | 1.000 | 1.000 | 0.000 | 0.000 | 0.000 | Winter | CVP |
| 1/7/2022 22:00 | 130 | 191 | late | F | Non-winter | 1.000 | Spring | 0.549 | 0.451 | 0.000 | 0.549 | 0.000 | Winter | CVP |

Proposed Action Assessment

5/24/2022

| Sample Date | FL* | JD ** | ots28 | Sex | Assignment | Pos Prob1 | Grp | Pos Prob2 | CV fall | CV late-fall | CV spring | CV winter | Model | Facility |
|--------------------|-----|-------|-------|-----|----------------------------------|-----------|-----------|-----------|---------|--------------|-----------|-----------|--------|----------|
| 1/9/2022 18:00 | 135 | 193 | late | F | Non-winter | 1.000 | Fall | 0.932 | 0.932 | 0.000 | 0.068 | 0.000 | Winter | CVP |
| 1/12/2022 22:00 | 153 | 196 | late | M | Non-winter | 1.000 | Fall | 0.988 | 0.988 | 0.000 | 0.012 | 0.000 | Winter | CVP |
| 2/15/2022 13:00 | 160 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.761 | N/A | N/A | N/A | N/A | N/A | CVP |
| 3/29/2022 9:00 | 197 | N/A | N/A | N/A | Non-winter | 1.000 | Late Fall | 0.983 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/11/2022 15:00 | 98 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.972 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/11/2022 15:00 | 205 | N/A | N/A | N/A | Non-winter | 1.000 | Late Fall | 0.767 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/13/2022 8:00 | 99 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.987 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/13/2022 15:00 | 93 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.824 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/14/2022 11:00 | 84 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.934 | N/A | N/A | N/A | N/A | N/A | SWP |
| N/A | N/A | N/A | N/A | N/A | STEELHEAD - qPCR confirmed | n/a | N/A | N/A | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/17/2022 23:59 | 84 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.898 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/18/2022 15:30 | 120 | N/A | N/A | N/A | Non-winter | 1.000 | Spring | 0.856 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/19/2022 4:00 | 97 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.973 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/20/2022 17:00 | 102 | N/A | N/A | N/A | Non-winter | 1.000 | U *** | 0.620 | N/A | N/A | N/A | N/A | N/A | SWP |

Proposed Action Assessment

5/24/2022

| Sample Date | FL* | JD ** | ots28 | Sex | Assignment | Pos Prob1 | Grp | Pos Prob2 | CV fall | CV late-fall | CV spring | CV winter | Model | Facility |
|--------------------|-----|-------|-------|-----|------------|-----------|--------|-----------|---------|--------------|-----------|-----------|-------|----------|
| 4/20/2022 17:38 | 90 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.915 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/20/2022 17:38 | 98 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.863 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/20/2022 17:38 | 110 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.950 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/20/2022 23:59 | 78 | N/A | N/A | N/A | Non-winter | 1.000 | Spring | 0.557 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/21/2022 2:00 | 86 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.996 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/21/2022 2:00 | 87 | N/A | N/A | N/A | Non-winter | 1.000 | U *** | 0.529 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/22/2022 11:00 | 90 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.896 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/22/2022 12:00 | 83 | N/A | N/A | N/A | Non-winter | 1.000 | Spring | 0.642 | N/A | N/A | N/A | N/A | N/A | CVP |
| 4/22/2022 15:45 | 80 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.875 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/22/2022 15:45 | 95 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.738 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/23/2022 15:00 | 100 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.990 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/24/2022 13:00 | 93 | N/A | N/A | N/A | Non-winter | 1.000 | U *** | 0.524 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/24/2022 15:40 | 82 | N/A | N/A | N/A | Non-winter | 1.000 | Spring | 0.827 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/24/2022 15:40 | 86 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.981 | N/A | N/A | N/A | N/A | N/A | SWP |
| 4/25/2022 11:00 | 86 | N/A | N/A | N/A | Non-winter | 1.000 | Fall | 0.825 | N/A | N/A | N/A | N/A | N/A | SWP |

Proposed Action Assessment
5/24/2022

* FL: Fork Length

** JD: Julian Date

*** U: Unassigned