## **PARTICIPANTS**

- California Department of Fish and Wildlife (CDFW)
- California Department of Water Resources (DWR)
- State Water Resources Control Board (SWRCB)
- U.S. Bureau of Reclamation (USBR)
- U.S. Fish and Wildlife Service (USFWS)
- Kearns & West (K&W)

### **ACTION ITEMS**

- SMT to set meeting summary review deadline as noon on Monday, November 28<sup>th</sup>.
- DWR to request weekly brookstock collection summaries from the Fish Conservation and Culture Laboratory (FCCL).

### **MEETING SUMMARY**

# PART 1: Updates on Water Operations and Biological Updates

## Relevant Actions & Triggers

There are currently no relevant actions or triggers for Old and Middle River (OMR) management. The first one will be the Integrated Early Winter Pulse Protection action, and this cannot be initiated until December 1, 2022. There have been no changes since last week in the Incidental Take Permit (ITP) Conditions of Approval (COA) that are currently in effect. Starting December 1, 2022, COA 8.3.1 (Integrated Early Winter Pulse Protection) and 8.3.3 (Adult Longfin Smelt Entrainment Protection) can be considered. The descriptions below are intended as summaries and do not provide all the details related to each action or trigger. For full descriptions, please see the OMR Guidance Document or ITP as needed.

# Proposed Action

OMR				
Management				
Measures	Requirement	Time Frame	Trigger	Triggered?
Integrated Early Winter Pulse Protection ("First Flush" Turbidity Event)	Reduce exports for 14 consecutive days so that the 14-day averaged OMR index for the period shall not be more negative than -2,000 cubic feet per second (cfs).	Dec 1 to Jan 31	(1) Running 3-day average of daily flows at Freeport >25,000 cfs; and (2) Running 3-day average of daily turbidity at Freeport ≥50 Nephelometric Turbidity Units (NTU¹); or (3) Real-time monitoring indicates a high risk of migration and dispersal into areas at high risk of future entrainment or a spent DS has been collected in monitoring surveys.	Not active
OMR Management	Manage to a more positive OMR than -5,000 cfs.	From the onset of OMR management to the end.	N/A	Not active
Turbidity Bridge Avoidance ("South Delta Turbidity")	If the daily average turbidity at Bacon Island cannot be maintained less than 12 NTU, manage exports to achieve an OMR no more negative than -2,000 cfs until the daily average turbidity at Bacon Island drops below 12 NTU.	After the first flush or Feb 1 (whichever comes first) and until a ripe or spent female DS is detected or April 1 (whichever is first).	Average daily turbidity in Old River at Bacon Island (OBI) at a level of more than 12 NTU.	Not active
Larval and Juvenile Delta Smelt	Run hydrodynamic models and forecasts of entrainment, informed by the Enhanced Delta Smelt Monitoring (EDSM) or other relevant survey data to estimate the percentage of larval and juvenile DS that could be entrained. If necessary, manage exports to limit entrainment to be protective based on the modeled recruitment levels.	On or after March 15 of each year until off- ramp criteria are met.	If QWEST is negative AND larval or juvenile DS are within the entrainment zone of the pumps based on real-time sampling of spawning adults or young of year life stages.	Not active

<sup>&</sup>lt;sup>1</sup> The current instrumentation measures turbidity in Formazin Nephelometric Units (FNUs).

OMR				
Management				
Measures	Requirement	Time Frame	Trigger	Triggered?
End of OMR	OMR criteria may control	During OMR	DS: when the daily mean water	Not active
Management	operations until June 30	management	temperature at Clifton Court	
	(for DS and Chinook	to June 30,	Forebay (CCF) reaches 77°F for 3	
	salmon), until June 15	or when the	consecutive days	
	(for steelhead/rainbow	DS		
	trout), or when the	temperature		
	species-specific off ramps	off ramp has		
	have occurred, whichever	been		
	is earlier.	reached.		

# ITP Conditions of Approval

Condition of				
Approval	Requirement	Time Frame	Trigger	Triggered?
8.1.5.2 (Smelt	Outlines contents for weekly	Nov 1 <sup>st</sup>	N/A	Active
Monitoring	risk assessments of DS and LFS	through June		
Team Risk	required under 8.1.5 and 8.1.1.	30 <sup>th</sup> or until		
Assessment)		off-ramped		
		by 8.8		
8.3.1	Reduce south Delta exports for	Dec 1 to Jan	3-day running average	Not active
(Integrated	14 consecutive days to	31	daily flows at Freeport	
Early Winter	maintain a 14-day average		greater than, or equal to,	
Pulse	OMR index no more negative		25,000 cfs, AND Three-	
Protection)	than -2,000 cfs, and convene		day running average of	
	the SMT within one day of		daily turbidity at Freeport	
	triggering. After maintaining a		is greater than, or equal	
	14-day average OMR index no		to, 50 FNU OR The SMT	
	more negative than -2,000 cfs		determines that real-	
	for 14 days, Permittee shall		time monitoring of	
	maintain a 14-day average		abiotic and biotic factors	
	OMR index no more negative		indicates a high risk of DS	
	than -5,000 cfs, initiating the		migration and dispersal	
	OMR Management season.		into areas at high risk of	
			future entrainment.	
8.3.3 (Adult	After December 1, if an	Dec 1	Salvage threshold for	Not active
Longfin Smelt	Integrated Early Winter Pulse	through Feb	water year (WY) 2022 is	
Entrainment	Protection (COA 8.3.1) has not	28th	one.	
Protection)	yet initiated, Permittee shall			
	reduce south Delta exports to			
	maintain a 14-day average			
	OMR index no more negative			
	than -5,000 cfs and initiate			
	OMR Management if:			
	Cumulative expanded salvage,			
	Dec 1 <sup>st</sup> through Feb 28 <sup>th</sup> ,			
	exceeds most recent Fall			
	Midwater Trawl (FMWT) Index			
	divided by 10, or SMT			
	determines that there is a high			
	risk of entrainment.			

Condition of				
Approval	Requirement	Time Frame	Trigger	Triggered?
8.4.1 (OMR Management for Adult Longfin Smelt)	The SMT shall conduct weekly risk assessments and decide whether to recommend an OMR flow requirement to minimize entrainment of adult LFS. The SMT may provide advice to restrict south Delta exports for seven consecutive days to achieve a seven-day average OMR index within three risk categories:  Low risk: OMR between -4,000 cfs to -5,000 cfs Medium risk: OMR between -2,500 cfs to -4,000 cfs High risk: OMR between -1,250 cfs to -2,500 cfs	Onset of OMR management through Feb 28 <sup>th</sup>	SMT recommendation based on weekly risk assessment.	Not active
8.4.2 (Larval and Juvenile Longfin Smelt Entrainment Protection)	If triggered, it will restrict south Delta exports for seven consecutive days in order to maintain a seven-day average OMR index no more negative than -5,000 cfs and convene the SMT to recommend an OMR flow limit between -1,250 and -5,000 cfs.	January 1st through June 30th or until the temperature offramp occurs	(1) LFS larvae or juveniles are found in four or more of the 12 Smelt Larvae Survey (SLS) or 20 mm stations in the central or south Delta, or (2) LFS catch per tow exceeds five larvae or juveniles in two or more of the 12 stations in the central or south Delta. The relevant stations are: 809, 812, 815, 901, 902, 906, 910, 912, 914, 915, 918 and 919.	Not active
8.4.3 High flow offramp for Longfin Smelt	If triggered, COA 8.4.1 and 8.4.2 are not required or would cease if previously required.	Throughout OMR management	When river flows are (a) greater than 55,000 cfs in the Sacramento River at Rio Vista or (b) greater than 8,000 cfs in the San Joaquin River at Vernalis. If flows subsequently drop below 40,000 cfs in the Sacramento River at Rio Vista or below 5,000 cfs in the San Joaquin River at Vernalis, the OMR limit previously required as a part of Conditions of Approval 8.4.1 and 8.4.2 shall resume.	Not active

Condition of				
Approval	Requirement	Time Frame	Trigger	Triggered?
	Requirement  Maintain daily average turbidity at OBI at a level of less than 12 NTU. If the daily average turbidity at OBI is greater than 12 NTU, Permittee shall restrict south Delta exports to achieve an OMR flow that is no more negative than -2,000 cfs until the daily average turbidity at OBI is less than 12 NTU.  If triggered, this COA will restrict south Delta exports for seven consecutive days in order to maintain a seven-day average OMR index no more negative than -5,000 cfs and SMT members will meet to assess the risk of entrainment. The SMT may provide further advice to restrict exports in order to maintain an OMR index more positive than -5,000 cfs. In their assessment, SMT members will determine if risk of entrainment is low, medium, or high; subsequent OMR restrictions will be based on	After the first flush or Feb 1 until end of OMR management or until CDFW agrees that the action may be ended or modified.  Nov 1st through June 30th or until off-ramped by 8.8	Trigger  Turbidity at OBI > 12 FNU  When the five-day salvage of juvenile DS is greater than or equal to one plus the average prior three years' FMWT index (rounded down). The 2022September FMWT index for DS was zero.	Active, not triggered
8.8 (End of OMR Management)	level of risk. Furthermore, if salvage of DS exceeds 11 in three days, this COA will restrict south Delta exports for seven consecutive days in order to maintain a seven-day average OMR index no more negative than -3,500 cfs.  If triggered, OMR Management would be offramped for LFS and DS.	From the onset of OMR management through	Daily mean water temperature at CCF is >25° C for three consecutive days.	Not active
8.12 (Barker Slough Pumping Plant Longfin and Delta Smelt Protection)	Barker Slough Pumping Plant will reduce exports so the maximum 7-day average is <60 cfs.	June 30 <sup>th</sup> From January 15 through March 31 in dry and critical water years for LFS, and from March 1 <sup>st</sup> through June 30 <sup>th</sup> for DS	Larval Smelt are detected at SLS Station 716 during the period identified for each species, and/or when recommended by the SMT.	Not active

## **Current Operations & Outlook**

USBR and DWR shared operations updates from the Outlook. Their observations included:

- USBR reported that releases from Whiskeytown Dam on Clear Creek are currently 200 cfs. No changes expected for the next seven-day period.
- Releases from Keswick Dam on the Sacramento River are 3,250 cfs.
- Releases at Trinity River are 300 cfs with the power bypass concluding last weekend.
- Releases from Nimbus Dam on the American River are 1,300 cfs. No changes expected for the next seven-day period.
- Releases from Goodwin Dam on the Stanislaus River have returned to a baseflow of 200 cfs after fall pulse flows. No anticipated changes.
- The federal facility is exporting two units with a change order to decrease to one unit on Monday, November 28<sup>th</sup>.
- A 400 cfs power bypass is in effect at Folsom, with a modification to 200 cfs planned for the afternoon of November 22<sup>nd</sup>.
- Delta Cross Channel (DCC) gates closed on Monday, November 21<sup>st</sup>. Gates will re-open Friday the 25<sup>th</sup> and close again the following Monday and likely remain closed unless the Delta encounters salinity issues.
- The last quarter of the neap tidal cycle concludes mid-November with a stronger tidal sequence initiating on the 23<sup>rd</sup>.
- DWR reported that Feather River releases are 1,600 cfs.
- November 21<sup>st</sup> Sacramento River flows were 1,500 cfs.
- San Joaquin River flows at Vernalis are around 675 cfs.
- State facility exports are 500 cfs and may increase to 1,000 cfs on November 24<sup>th</sup>.
- Delta outflows peaked at 3,700 cfs on November 21st.
- QWEST, with the DCC gates open, reached 1,500 cfs and will decrease to -500 cfs with the gates closed.
- Rio Vista flows are increasing with the DCC gates closed reaching nearly 5,000 cfs and dropping to 2,000 cfs when DCC gates open over the weekend.
- Freeport flows are ranging from 5,500 to 7,000 cfs.
- The daily OMR Index is approximately -2,000 cfs and may trend more positive as combined exports decrease after November 24<sup>th</sup>.
  - November 19<sup>th</sup> OMRI
    - Daily: -2,600 cfs
    - Five-Day: -2,500 cfs
    - 14-Day: -2,700 cfs
  - November 19<sup>th</sup> Index Calculations
    - Daily: -2,100 cfs
    - Five-Day: -2,200 cfs
    - 14-Day: -2,300 cfs
  - November 21<sup>st</sup> Index Calculations
    - Daily: -2,100 cfs
    - Five-Day: -2,100 cfs
    - 14-Day: -2,300 cfs

- USBR clarified that a power bypass is when water travels through the lowest outlet valves bypassing the powerhouse in order to draw out the coldest water.
- Water quality is a topic of interest as spring tides push into the Delta. The neap cycle next week should improve water quality by reducing salinity.
- Sacramento River X2 is 95.2 km as of November 20<sup>th</sup>.
- The following updates were made to the survey table:
  - o Knights Landing started trapping on November 16<sup>th</sup>.
  - The lower Sacramento rotary screw trap is active.
  - o San Joaquin River Restoration Program USFWS and USBR Field Monitoring is not active.
  - CVP regular counts, CWT reading is active.

## Review of Environmental Conditions and Survey Updates

CDFW delivered catch updates on relevant surveys to the SMT.

- FMWT Survey October LFS data:
  - Life Stage:

Juvenile: 97Adult: Two

- o Regions:
  - San Pablo Bay: 95 (44 95 mm)
  - Suisun Bay: Four (61 83 mm)
- October LFS index: 261
- November index is on track for December 1<sup>st</sup> distribution.
- San Francisco Bay Study November LFS data:
  - Life Stage:

Juvenile: 73Adult: Three

- Regions:
  - San Pablo Bay: 41 (51 84 mm)
  - Suisun Bay: 10 (58 88 mm)
  - Lower Sacramento River: One (64 mm)

USFWS provided catch updates on the Enhanced Delta Smelt Monitoring Program (EDSM) and Chipps Island Trawl.

- EDSM sampled Monday through Thursday the week of November 14<sup>th</sup> completing all 36 sites.
  - o DS: Zero
  - LFS: 20 in Suisun Marsh (60 to 104 mm).
    - Two LFS adults were transferred to FCCL for broodstock.
      - No information exists on the status of the transferred fish.
  - o EDSM will be sampling Monday to Friday this week with Thursday off as a holiday.
- The week of November 14<sup>th</sup> Chipps Island crews completed 25 of 30 scheduled tows. Half a sampling day was cancelled.
  - o DS: Zero
  - LFS: One (72 mm)
  - Chipps Island will sample this week on Monday, Tuesday, and Wednesday due to the holiday.
- The DS abundance estimate for the week of November 14<sup>th</sup> was zero due to no detections.

• The last non-zero abundance estimate is from the week of November 7<sup>th</sup> at 1,240.

CDFW provided a salvage update (November 14<sup>th</sup> to November 20<sup>th</sup>).

• No DS or LFS have been detected at either facility this WY.

USBR shared environmental data updates.

No pertinent environmental conditions to note other than possible precipitation early next week.

# PART 2: Open Discussion on Species Status (Structured-Unstructured Time)

USBR and CDFW noted no significant changes since last week for DS.

CDFW informed that the first experimental release of DS of the WY begins Monday, November 28<sup>th</sup> and concludes the following Friday. Additional information will be covered during next week's SMT meeting.

CDFW highlighted that water temperatures are entering the range consistent with LFS spawning. With X2 upstream of the confluence it is expected that LFS may travel further east than observed in previous years. Current detections are centered around San Pablo Bay, but the fish will likely move towards fresher water as the season progresses.

No actions or conditions of approval were triggered this week and the SMT did not make any recommendations.

### PART 3: Live-edit Assessments

# Proposed Action Weekly Evaluation of Delta Smelt, including Distribution, Abiotic Conditions, Risk Assessment Questions, and Executive Summary

USBR reviewed proposed changes to the PA Assessment, which include the latest dates, detections, data, and the following:

- Language was amended in the Biological Conditions and Executive Summary to note recent DS detections are limited to the Lower Sacramento River.
- Noted that the San Joaquin River CDEC X2 tool is down.

## ITP Longfin Smelt Risk Assessment

The SMT reviewed and discussed updates to the ITP Risk Assessment for LFS, which include the latest dates, detections, and data as well as:

### Advice to WOMT

• No items for elevation to WOMT.

#### Sections 1-A and 1-B

 No modifications to risk, however LFS rationale now captures that water temperatures are conducive to LFS spawning.

#### **Executive Summary**

• No modifications to the executive summaries beyond specifying locations of LFS detections in the Delta and current conditions being conducive for LFS spawning.

## Part 4: Additional Considerations/Discussion

Agencies reported no items for elevation to WOMT.