

Contents

	Page
Acknowledgments	iii
Acronyms and Abbreviations	xvii
Executive Summary	xix
Chapter 1. Introduction.....	1
1.1. Background.....	1
1.2. Development, Implementation, and Scope of the Land Retirement Demonstration Project	3
1.3. Tranquillity Project Site Location and Description	5
1.4. Literature Cited	7
Chapter 2. Physical Impacts of Agricultural Land Retirement in the San Joaquin Valley	9
2.1. Introduction.....	9
2.2. Tranquillity Site	10
2.2.1. Geology	10
2.2.2. Soils	10
2.2.3. Groundwater	13
2.2.4. Climate and Weather	14
2.2.5. Irrigation	16
2.2.6. Monitoring of Selenium Levels.....	16
2.2.6.1. Surface Water Monitoring	16
2.2.6.2. Soil Monitoring.....	17
2.2.6.3. Soil Chemical Analyses	19
2.2.6.4. Groundwater Level Monitoring	19
2.2.6.5. Groundwater Quality Monitoring	21
2.2.7. Groundwater Response to Land Retirement.....	22
2.2.8. Groundwater Quality	32
2.2.8.1. Groundwater Salinity	32
2.2.8.2. Groundwater Major Ion Chemistry.....	32
2.2.8.3. Trace Elements in Groundwater	35
2.2.8.4. Selenium Trend in Groundwater.....	38
2.2.8.5. Boron in Groundwater	44
2.2.8.6. Origin and Isotopic Composition of Groundwater	44
2.2.8.7. Groundwater Age.....	45
2.2.9. Evaporative Concentration of Shallow Groundwater.....	46
2.2.10. Trend Analysis for Soils.....	48
2.2.11. Change Detection Analyses.....	51
2.2.12. Soil Interpretive Summary.....	52
2.2.13. Conclusions	59

2.3. Atwell Island Site.....	60
2.3.1. Geology	60
2.3.2. Soils	60
2.3.3. Weather.....	67
2.3.4. Irrigation	67
2.3.5. Hydrology and Surface Water Monitoring.....	68
2.3.6. Groundwater Level Monitoring.....	69
2.3.7. Groundwater Quality Monitoring.....	69
2.3.7.1. Groundwater Response to Land Retirement.....	71
2.3.7.2. Groundwater Salinity.....	71
2.3.7.3. Groundwater Major Ion Chemistry.....	72
2.3.7.4. Selenium in Groundwater	73
2.3.8. Conclusions	73
2.4. Literature Cited.....	74
Chapter 3. Selenium Levels of Biota from Retired Agricultural Lands in the San Joaquin Valley, California	79
3.1. Introduction.....	79
3.2. Methods.....	81
3.3. Results.....	83
3.3.1. Plants	83
3.3.2. Invertebrates	90
3.3.3. Small Mammals.....	96
3.4. Discussion and Conclusions	101
3.5. Literature Cited.....	103
Chapter 4. Results of a Habitat Restoration Study on Retired Agricultural Lands in the San Joaquin Valley, California	107
4.1. Introduction.....	107
4.2. Methods.....	108
4.2.1. Study Design	108
4.2.2. Restoration Methods.....	108
4.2.3. Plot-Level Monitoring.....	111
4.2.3.1. Vegetation Survey Methods.....	111
4.2.3.2. Invertebrates.....	112
4.2.3.3. Amphibians and Reptiles	114
4.2.3.4. Birds.....	114
4.2.3.5. Small Mammals	115
4.2.4. Site-Wide Surveys.....	115
4.2.4.1. Spotlighting Surveys.....	115
4.2.4.2. Track Station Surveys.....	116
4.2.4.3. Winter Raptor Surveys.....	117
4.2.4.4. Statistical Analysis of Vegetation Data	118
4.2.4.5. Analysis of Data for the Other Biotic Groups	119

4.3. Results.....	120
4.3.1. Plot-Level Monitoring.....	120
4.3.1.1. Vegetation.....	120
4.3.1.2. Invertebrates.....	129
4.3.1.3. Amphibians and Reptiles.....	134
4.3.1.4. Birds.....	135
4.3.1.5. Small Mammals.....	144
4.3.2. Site-Wide Surveys.....	146
4.3.2.1. Spotlighting Surveys.....	146
4.3.2.2. Track Station Surveys.....	151
4.3.2.3. Winter Raptor Surveys.....	155
4.4. Discussion.....	156
4.4.1. Vegetation.....	156
4.4.2. Invertebrates.....	158
4.4.3. Amphibians and Reptiles.....	160
4.4.4. Birds.....	161
4.4.5. Small Mammals.....	163
4.4.6. Site-Wide Surveys.....	163
4.4.6.1. Spotlighting Surveys.....	163
4.4.6.2. Track Station Surveys.....	164
4.4.6.3. Winter Raptor Surveys.....	164
4.5. Conclusions and Recommendations.....	165
4.5.1. Conclusions.....	165
4.5.2. Recommendations.....	168
4.6. Literature Cited.....	170
Chapter 5. Overview of the Implications and Concerns of Agricultural	
Land Retirement in the San Joaquin Valley, California.....	177
5.1. Declining Groundwater.....	178
5.2. Selenium in Groundwater.....	178
5.3. Selenium in Soil.....	178
5.4. Selenium Concentrations in Biota.....	179
5.5. Surface Water Ponding.....	179
5.6. Wildlife Abundance, Diversity, and Recovery of Sensitive Species.....	179
5.7. Restoration Challenges and Techniques.....	180
5.8. Agricultural Pests.....	183
5.9. Recommendation.....	183
Appendix 1. CVPIA Land Retirement Program Chronology	
Appendix 2. Compliance with Biological Opinion	
Appendix 3. Research and Restoration	

Tables

Table 2-1. Estimates of Natural drainage rates and vertical permeability at the Tranquillity site. 29

Table 2-2. Groundwater level decline observed in fifteen wells at the Tranquillity site for the time period from October 1999 to October 2004..... 30

Table 2-3. Baseline year groundwater quality data for shallow wells at the Tranquillity site — major ions and field parameters. 33

Table 2--4. Baseline year groundwater quality data for deep wells at the Tranquillity site — major ions and field parameters. 33

Table 2--5. Baseline groundwater quality data for shallow wells at the Tranquillity site — trace elements and tritium. 34

Table 2-6. Baseline groundwater quality data for deep wells at the Tranquillity site— trace elements and tritium. 34

Table 2-7. Summary statistics for selenium concentrations measured in groundwater samples taken from Coast Range deposits for the 5 years of groundwater monitoring at the Tranquillity Site. 38

Table 2-8. Analysis of seasonality of selenium data based on Kruskal-Wallis 1-way ANOVA..... 39

Table 2-9. Parametric and nonparametric correlations of selenium concentration in the Tranquillity wells on time 40

Table 2-10. Summary of results of linear and nonlinear trends without and with adjustment for outliers. 41

Table 2-11. Change detection analyses for COCs in soil at the Tranquillity site..... 52

Table 2-12. Comparison of total selenium concentrations at the Tranquillity site to Western U.S. and San Joaquin Valley soils. 52

Table 2-13. Average soluble to total selenium percentages all sites at Tranquillity..... 53

Table 2-14. Total selenium (mg/kg dry soil) 63

Table 2-15. Soluble selenium 1-5 extract (mg/kg dry soil) 63

Table 2-16. A comparison of total selenium concentrations with soluble selenium concentrations..... 64

Table 2-17. Soil salinity (EC_e dS/m) at the Atwell Island site 65

Table 2-18. Soluble boron 1-5 extract (mg/kg dry soil) at the Atwell Island site. 66

Table 2-19. Estimated 2002 net crop water requirement and deep percolation losses at the Atwell Island site. 68

Table 2-20. Baseline groundwater quality data for shallow wells at the Atwell Island site - major ions, field parameters, and selenium. 72

Table 3-1. List of biotic groups monitored, corresponding performance standards (USFWS 1999), and an indication whether performance standards were met. 80

Table 3-2. Geometric means (Mean), sample size (N), number of non-detects (ND), minimum (Min), maximum (Max), and standard deviation factor (SD factor) of selenium concentrations (mg/kg dry weight) in plant species collected from Tranquillity, 1999 to 2003. 85

Table 3-3. Geometric means (Mean), sample size (N), number of non-detects (ND), minimum (Min), maximum (Max), and standard deviation factor (SD factor) of selenium concentrations (mg/kg dry weight) in plant species collected from Atwell Island, 2000 to 2002. 86

Table 3-4. Results of analysis of variance tests of selenium concentrations (mg/kg dry weight) between sample years in plant species collected from Tranquillity, 1999 to 2003. 87

Table 3-5. Results of Analysis of Variance tests of selenium concentrations (mg/kg dry weight) between sample years in plant species collected from Atwell Island, 2000 to 2002. 87

Table 3-6. List of plant samples from Tranquillity that exceeded the population-level performance standard (2.0 mg/kg). 88

Table 3-7. Results of Analysis of Variance tests of selenium concentrations (mg/kg dry weight) between landform types in plant species collected from Tranquillity, 1999 to 2003. 88

Table 3-8. Results of Analysis of Variance tests of selenium concentrations (mg/kg) between landform types in plant species collected from Atwell Island, 2000 to 2002. 89

Table 3-9. Results of Analysis of Variance tests of selenium concentrations (mg/kg) between plant parts in plant species collected from Tranquillity, 1999 to 2003. 90

Table 3-10. Results of Analysis of Variance tests of selenium concentrations (mg/kg) between plant parts in plant species collected from Atwell Island, 2000 to 2002. 90

Table 4-1. Overview of shrub island planting, Tranquillity HRS. Figures in the Initial Planting columns indicate the number of individuals that were initially transplanted to each island. 110

Table 4-2. Plot-level survey dates for the various biotic groups, 1999 to 2003. 112

Table 4-3. Dates of spotlighting surveys, track station surveys, and winter raptor surveys conducted from 1999 to 2003. 116

Table 4-4. Overall vegetation species richness in the Tranquillity HRS study plots.	120
Table 4-5. Species imprinted in the Tranquillity HRS Plots (1999), and an overview of those imprinted species noted in the quadrats during spring vegetation monitoring, 2000-2003.....	121
Table 4-6. Non-imprinted, native species (excluding <i>Atriplex argentea</i>) noted in the quadrats during spring vegetation monitoring in the Tranquillity HRS, 1999-2003.	122
Table 4-7. All species noted in the quadrats during spring vegetation monitoring in the Tranquillity HRS, 2003, including species code, common name, life-form, origin, mean percent cover, and relative abundance.	123
Table 4-8. Invertebrate families collected on the Tranquillity study plots that contain agriculturally beneficial species, 1999 to 2003.....	133
Table 4-9. Invertebrate families collected on the Tranquillity study plots that contain agricultural pest species, 1999 to 2003.	134
Table 4-10. Diversity of amphibians and reptiles observed at Tranquillity 1999-2003.	135
Table 4-11. Special status bird species observed on the Tranquillity study plots, 1999 to 2003.	135
Table 4-12. Blocks with the highest mean richness of birds, season and year of occurrence, and the magnitude of difference from other blocks. ...	136
Table 4-13. Blocks with the highest mean abundance of birds, season and year of occurrence, and the magnitude of difference from other blocks. ...	138
Table 4-14. Blocks with the highest mean abundance of grassland obligate birds, grassland facultative birds, and other birds, season and year of occurrence, and the magnitude of difference from other blocks.....	139
Table 4-15. Nest abundance and species richness, 2002 and 2003. Parentheses indicate nest numbers off study plot areas and asterisks indicate nests not monitored.	140
Table 4-16. Mean nest initiation dates, mean fledging dates, mean clutch size, and nesting success rates of bird species nesting on the Tranquillity study site, 2002 and 2003.	142
Table 4-17. Vegetative composition and ground cover at mallard and red-winged blackbird nests.	143
Table 4-18. The number of small mammals captured in live traps and pitfall traps on the Habitat Restoration Study plots, Tranquillity site, 1999 to 2003.	144
Table 4-19. The number of ornate shrews (<i>Sorex ornatus</i>), California voles (<i>Microtus californicus</i>), and Botta's pocket gophers captured in the pitfall traps at the Tranquillity study site, 1999 to 2003.....	145

Table 4-20. Rates of observation (mean number per kilometer) of wildlife species observed during spotlight surveys conducted near Tranquillity, California, 1999 to 2003. Asterisks denote zero observations..... 148

Table 4-21. Annual numbers and mean rates of visitation of wildlife at track stations established at the Tranquillity site, 1999 to 2003. Asterisks denote zero observations. 152

Table 4-22. Seasonal mean rates of visitation of wildlife at track stations established at the Tranquillity site, 1999 to 2003. 153

Table 4-23. Frequency and rate of bird species encountered at the Tranquillity site, 1999 to 2003 during winter raptor surveys. 156

Table A3-1. Annual monitoring activities on the Habitat Restoration Study plots (Tranquillity site), 1999-2003, with the number of times each survey was conducted per year. 4

Table A3-2. Annual site-wide monitoring activities at the Tranquillity site, 1999-2003, with the number of times each survey was conducted per year..... 4

Table A3-3. Annual monitoring activities on the Habitat Restoration Study plots (Atwell Island site) 2001-02, with the number of times each survey was conducted per year. 5

Figures

Figure 1-1. Drainage-impaired lands in the San Joaquin Valley..... 2

Figure 1-2. Location of the Tranquillity site including the 178 ha (440 ac) North Avenue parcel and the 16 ha (40 ac) Manning Avenue parcel..... 6

Figure 2-1. U.S. Department of Agriculture soil mapping units, Tranquillity site..... 12

Figure 2-2. Conceptual site model for groundwater conditions at the Tranquillity site..... 14

Figure 2-3. Monthly precipitation at CIMIS Station 105, Westlands Water District, near the Tranquillity site (1993-2003). 15

Figure 2-4. Annual precipitation at CIMIS Station 105, Westlands Water District, near the Tranquillity site (1993-2003). 15

Figure 2-5. Soil sample locations at the Tranquillity site. 18

Figure 2-6. Monitor well and sump location map, Tranquillity site..... 20

Figure 2-7. Hydrographs for three agricultural drain sumps at the Tranquillity site showing a declining shallow groundwater trend..... 23

Figure 2-8. Hydrographs for the monitor well cluster site 325 and 326 at the Tranquillity site showing a declining shallow groundwater trend and vertical (downward) hydraulic gradient. 24

Figure 2-9. Hydrographs for monitor wells 15M1 and 16A1 at the Tranquillity site showing a declining shallow groundwater trend..... 24

Figure 2-10. Depth to shallow groundwater, October 1999. The project area underlain by shallow groundwater within 2.1 m (7 feet) of the land surface is approximately 243 ha (600 acres)..... 25

Figure 2-11. Depth to shallow groundwater, October 2000. The project area underlain by shallow groundwater within 2.1 m (7 feet) of the land surface is approximately 22 ha (55 acres)..... 26

Figure 2-12. Depth to shallow groundwater, October 2001. The project area underlain by shallow groundwater within 2.1 m (7 feet) of the land surface is approximately 14 ha (34 acres)..... 27

Figure 2-13. Contour map showing spatial distribution of net water table declines at the Tranquillity site from October 1999 to October 2004. 31

Figure 2-14. Comparison of dominant major ion concentrations and electrical conductivity for groundwater samples from shallow and deep wells. 35

Figure 2-15. Ratio of selenium to manganese (Se/Mn) concentration in groundwater samples plotted verses well depth. The Se/Mn ratio shows a decreasing trend with depth indicating oxidizing geochemical conditions in the shallow wells and reducing conditions in the deep wells. 37

Figure 2-16. Boxplots of selenium concentrations in Coast Range deposits for the 5 years of groundwater sampling at the Tranquillity site. The inner black boxes represent the 95 percent confidence interval of the median concentration. 38

Figure 2-17. Selenium data and trend in well 16B1 along with “best fit” nonlinear regression [Sinusoidal Fit: $y=a+b*\cos(cx+d)$]. 42

Figure 2-18. Selenium data and trend in well 16B1 without outlier datum along with “best fit” nonlinear regression - [Sinusoidal Fit: $y=a+b*\cos(cx+d)$]. 42

Figure 2-19. Selenium data and trend in well 16B1 along with “best fit” nonlinear regression - [Harris Model: $y=1/(a+bx^c)$]. 43

Figure 2-20. Selenium data and trend in well 16Q1 along with ‘best fit’ nonlinear regression - [Rational Function: $y=(a+bx)/(1+cx+dx^2)$]. 43

Figure 2-21. Plot of stable isotope data for groundwater samples from the Tranquillity site indicating the shallow groundwater has undergone significant evaporation. 45

Figure 2-22. Photos of open surface cracks in Tranquillity clay soils at the Tranquillity site (Jan 2005). 47

Figure 2-23. EC_e Trends in top foot of soil from 1999 to 2004..... 48

Figure 2-24. EC_e Trends in 2 to 3 foot depth of soil from 1999 to 2004..... 49

Figure 2-25. EC_e Trends in 4 to 5 foot depth of soil from 1999 to 2004..... 49

Figure 2-26. Soluble selenium trends in top foot of soil from 1999 to 2004..... 49

Figure 2-27. Soluble selenium trends in 2 to 3-foot depths of soil from 1999 to 2004. 50

Figure 2-28. Soluble selenium trends in 4 to 5 foot depths of soil from 1999 to 2004. 50

Figure 2-29. Total selenium trends in top foot of soil from 1999 to 2004. 50

Figure 2-30. Total selenium trends in 2 to 3 foot depth from 1999 to 2004..... 51

Figure 2-31. Total selenium trends in 4 to 5 foot depth from 1999 to 2004..... 51

Figure 2-32. Total selenium concentrations in surface soils 0-30 cm (0-1 foot) at the Tranquillity site from the 2004 sampling event. 55

Figure 2-33. Soluble selenium in surface soils 0-30 cm (0-1 foot) at the Tranquillity site from the 2004 sampling event..... 56

Figure 2-34. Soluble boron concentrations in the surface soils 0-30 cm (0-1 foot) at the Tranquillity site from the 2004 sampling event. 57

Figure 2-35. Soil salinity (ECe) concentrations (dS/m) in the surface soils 0-30 cm (0-1 foot) at the Tranquillity site from the 2004 sampling event..... 58

Figure 2-36. U.S. Department of Agriculture soil mapping units and soil sample locations at the Atwell Island site..... 62

Figure 2-37. Monitor well locations at the Atwell Island site. 70

Figure 2-38. Hydrographs of groundwater levels observed in Wells 5B3 and 5B4 showing pre-project and post-project groundwater levels..... 71

Figure 3-1. Geometric means (solid circles) and error bars (mean multiplied by and divided by one standard deviation factor) of selenium concentrations in crickets collected from the Tranquillity and Atwell Island study sites. 92

Figure 3-2. Geometric means (solid circles) and error bars (mean multiplied by and divided by one standard deviation factor) of selenium concentrations in beetles collected from the Tranquillity and Atwell Island study sites. 93

Figure 3-3. Geometric means (solid circles) and error bars (mean multiplied by and divided by one standard deviation factor) of selenium concentrations in spiders collected from the Tranquillity and Atwell Island study sites. 94

Figure 3-4. Geometric means (solid circles) and error bars (mean multiplied by and divided by one standard deviation factor) of selenium concentrations in isopods collected from the Tranquillity and Atwell Island study sites. 95

Figure 3-5. Geometric means (solid circles) and error bars (mean multiplied by and divided by one standard deviation factor) of selenium concentrations in deer mouse (PEMA) bodies collected from Tranquillity and Atwell Island. 97

Figure 3-6. Geometric means (solid circles) and error bars (geometric mean multiplied by and divided by one standard deviation factor) of selenium concentrations in deer mouse (PEMA) livers collected from Tranquillity and Atwell Island. 98

Figure 3-7. Geometric means of selenium concentrations in deer mouse (PEMA) bodies and livers collected from irrigated and non-irrigated lands at Tranquillity, 1999 to 2003. 99

Figure 3-8. Geometric means (solid circles) and error bars (geometric mean multiplied by and divided by one standard deviation factor) of selenium concentrations in shrew (SOOR) bodies collected from Tranquillity. 100

Figure 3-9. Geometric means (solid circles) and standard deviations of selenium concentrations in shrew (SOOR) livers collected from the Tranquillity study site. 101

Figure 4-1. Map of the Tranquillity HRS plots showing treatments and blocks. 109

Figure 4-2. Locations of pitfall arrays, amphibian and reptile coverboards, avian transects and point-count locations, and small mammal trapping lines on a study plot at the Tranquillity site. 113

Figure 4-3. Locations of the spotlight survey route, track stations, and winter raptor survey route, 1999 to 2003. 117

Figure 4-4. Distribution of *Bromus madritensis* (red brome) at the Tranquillity HRS, 2003. Each small rectangle represents a ‘sextant’; i.e., the portion of the plot in which a particular group of four sampling quadrats was located. 124

Figure 4-5. Distribution of *Sisymbrium irio* (London rocket) at the Tranquillity HRS. Each small rectangle represents a ‘sextant’, i.e., the portion of the plot in which a particular group of four sampling quadrats was located. 125

Figure 4-6. Species rank-abundance by treatment, Tranquillity HRS Plots 2003. Species letter codes correspond to those presented in Table 4-7. ... 127

Figure 4-7. Species rank-abundance by treatment, Tranquillity HRS Plots 2003. 128

Figure 4-8. Ordinal richness of invertebrates for all treatments in 1999 to 2003. 130

Figure 4-9. Mean invertebrate abundance for all treatments in 1999 to 2003. 131

Figure 4-10. Mean abundance of Coleoptera and Araneae for all treatments in 1999 to 2003. 132

Figure 4-11. Mean richness of birds by treatment in winter, spring, summer, and fall 1999 to 2003. 136

Figure 4-12. Mean abundance of birds by treatment in winter, spring, summer, and fall 1999 to 2003. 137

Figure 4-13. Mean abundance (mean number of birds observed per plot per day, all seasons combined) of obligate grassland birds, facultative grassland birds, and other birds, 1999 to 2003. 138

Figure 4-14. Mean abundance of grassland obligates and facultative birds by treatment, 1999 to 2003. 139

Figure 4-15. Locations of bird nests found in 2002 at the Tranquillity study site. 141

Figure 4-16. Locations of bird nests found in 2003 at the Tranquillity study site. 141

Figure 4-17. Mean abundance of small mammals per 120 trap nights in winter, spring, summer, and fall at the Tranquillity study site, 1999 to 2003. 146

Figure 4-18. Annual wildlife abundance observed during spotlight surveys conducted near Tranquillity, California, 1999 to 2003. 147

Figure 4-19. Seasonal abundance of wildlife observed during spotlight surveys conducted near Tranquillity, California, 1999 to 2003. 149

Figure 4-20. Abundance of barn owls (*Tyto alba*) and western burrowing owls (*Athene cunicularia hypugea*) observed during spotlight surveys conducted near Tranquillity, California, 1999 to 2003. 150

Figure 4-21. Abundance of desert cottontails (*Sylvilagus auduboni*) and black-tailed hares (*Lepus californicus*) observed during summer spotlight surveys conducted near Tranquillity, California, 1999 to 2003. 150

Figure 4-22. Seasonal abundance of coyotes (*Canis latrans*) and domestic dogs (*Canis familiaris*) observed during spotlight surveys conducted near Tranquillity, California, 1999 to 2003. 151

Figure 4-23. The yearly abundance (mean rate of visitation) of wildlife during track station surveys conducted at the Tranquillity site, 1999 to 2003. 154

Figure 4-24. The seasonal abundance (mean rate of visitation) of wildlife at track stations established at the Tranquillity site, 1999 to 2003. 154