

Final

Fisheries and Aquatic Ecosystems Technical Report

Shasta Lake Water Resources Investigation, California

Prepared by:

**United States Department of the Interior
Bureau of Reclamation
Mid-Pacific Region**



December 2014

Contents

Chapter 1	Affected Environment	1-1
1.1	Environmental Setting	1-1
1.1.1	Aquatic Habitat	1-1
1.1.2	Fisheries Resources.....	1-13
1.1.3	Aquatic Macroinvertebrates.....	1-53
Chapter 2	Impact Assessment.....	2-1
Chapter 3	References.....	3-1
3.1	Printed Sources	3-1

Tables

Table 1-1.	Fish Species Known to Occur in Primary Study Area	1-13
Table 1-2.	Frequency and Estimated Area of Suitable Coldwater Fish Spawning Habitat in Shasta Lake Tributaries Estimated to Occur Within Projected Varial Zones of Reservoir Enlargement Alternatives ^a	1-20
Table 1-3.	Frequency of Occurrence of Shasta Lake Tributaries with Fish Passage Impediments, by Elevation Range, Among Streams Surveyed in 2012.....	1-21
Table 1-4.	Estimated Percent Occurrence of Partial and Complete Fish Passage Barriers in Shasta Lake Tributaries Within the Projected Varial Zones of Reservoir Enlargement Alternatives	1-22
Table 1-5.	Frequency and Proportion of Fish-Bearing Tributaries to Shasta Lake Surveyed in 2012	1-23
Table 1-6.	Frequency and Proportion of Shasta Lake Tributaries Inhabited by Hardhead ^a and Foothill Yellow-Legged Frog ^b Based on Streams Surveyed in 2011 to 2012.....	1-24
Table 1-7.	Central Valley Fish Species Potentially Affected by Project Alternatives	1-26
Table 1-8.	Life History and Distributions of Evaluation Fish Life Stages in Primary and Extended Study Areas.....	1-29

Figures

Figure 1-1. San Francisco Bay and Sacramento-San Joaquin Delta.....	1-9
Figure 1-2. Seasonal Occurrence of Different Life Stages of Four Chinook Salmon Runs	1-33

Attachments

Attachment 1	Assessment of Fisheries Impacts Within the Sacramento-San Joaquin Delta
--------------	---

Abbreviations and Acronyms

°C	degrees Celsius
°F	degrees Fahrenheit
ACID	Anderson-Cottonwood Irrigation District
Bay	San Francisco Bay
Bay-Delta	San Francisco Bay-Delta
BMI	benthic macroinvertebrates
BO	Biological Opinion
CALFED	CALFED Bay-Delta Program
CDFG	California Department of Fish and Game
CDFW	California Department of Fish and Wildlife
CESA	California Endangered Species Act
cfs	cubic feet per second
CVP	Central Valley Project
CVPIA	Central Valley Project Improvement Act
DCC	Delta Cross Channel
Delta	Sacramento-San Joaquin Delta
DEM	Digital Elevation Model
DO	dissolved oxygen
DPS	distinct population segment
DWR	California Department of Water Resources
ESA	Federal Endangered Species Act
ESU	Evolutionarily Significant Unit
FR	Federal Register
GIS	geographic information system
KMP	Klamath Mountains Province
MAF	million acre-feet
msl	mean sea level
NGO	nongovernmental organization
NMFS	National Marine Fisheries Service
ppt	parts per thousand
RBPP	Red Bluff Pumping Plant
Reclamation	U.S. Department of the Interior, Bureau of Reclamation
RK	river kilometer
RM	river mile

Shasta Lake Water Resources Investigation
Biological Resources Appendix – Fisheries and Aquatic Ecosystems Technical Report

SONCC	Southern Oregon/Northern California Coast
State	State of California
STNF	Shasta-Trinity National Forest
SWP	State Water Project
TCD	temperature control device
UKT	Upper Klamath Trinity
USFS	U. S. Forest Service
USFWS	U.S. Fish and Wildlife Service
WSEL	water surface elevation
X2	salinity isopleth

Chapter 1

Affected Environment

1.1 Environmental Setting

This chapter describes the affected environment as it relates to fisheries and the aquatic ecosystems in the study area.

The primary study area includes Shasta Lake and the lower reaches of its major and minor tributaries, and the Sacramento River from Shasta Dam to Red Bluff Pumping Plant (RBPP). Because of the potential for a project at Shasta Dam to affect resources outside the primary study area, information on an extended study area is also included. For the purpose of fisheries and the aquatic ecosystems, this extended study area includes the Sacramento River downstream to the Sacramento-San Joaquin Delta (Delta). It also includes portions of the lower Feather River, lower American River, and lower San Joaquin River basins, and the water service areas of the Central Valley Project (CVP) and State Water Project (SWP). The Trinity River is also included in the affected environment because CVP and SWP operations in response to project operation alternatives have the potential to affect Trinity River flows.

Descriptions of fisheries and the aquatic ecosystems were derived primarily from the following sources:

- *Assessment of Fisheries Impacts Within the Sacramento-San Joaquin Delta* (Attachment 1)
- *Shasta Lake Water Resources Investigation Mission Statement Milestone Report* (Reclamation 2003)
- *Shasta Lake Water Resources Investigation Initial Alternatives Information Report* (Reclamation 2004)
- *Tributary Fisheries Characterization Report* (Reclamation 2014)
- Chapter 3, “Biological Environment,” in *Draft Shasta Lake Water Resources Investigation Plan Formulation Report* (Reclamation 2007)

1.1.1 Aquatic Habitat

This section briefly describes the aquatic habitats in the primary and extended study areas and CVP and SWP service areas. Factors affecting the abundance and distribution of fish populations are described under a separate section titled “Fisheries Resources” below.

Primary Study Area

The primary study area includes Shasta Lake and primary upstream tributaries and the Sacramento River from Shasta Dam downstream to Keswick Dam. The Sacramento River supports the largest contiguous riverine and wetland ecosystems in the Central Valley and provides 35 percent of the State of California's (State) water supply. Most of the Sacramento River flow is controlled by the U.S. Department of the Interior, Bureau of Reclamation's (Reclamation), Shasta Dam, and river flow is augmented in average water years by transfer of up to 1 million acre-feet (MAF) of Trinity River water through Clear Creek and Spring Creek tunnels to Keswick Reservoir (Reclamation 2004). Water resources development, including the construction of dams and diversions, has affected the hydrology, geomorphology, and ecology of the watershed. Many of these effects have been detrimental to local aquatic habitats and species.

Shasta Lake and Vicinity Shasta Dam and Shasta Lake are located on the upper Sacramento River in northern California. Shasta Dam is located about 9 miles northwest of the city of Redding, and the dam and entire reservoir are within Shasta County.

Today, the current composition and distribution of fish species inhabiting Shasta Lake reflect the historic fisheries, the operational effects of Shasta Dam as well as dams on several of the upstream tributaries, and the introduction of nonnative fish species. Shasta Lake fish species include native and nonnative species, which are dominated by mostly introduced warm-water and cold-water species (Weidlein 1971; CDFG, unpublished data). Shasta Lake tributary fish species comprise several native and nonnative species and have been managed to favor naturally produced ("wild") and stocked (hatchery-cultured) native and nonnative trout species (Rode 1989, Moyle 2002, Rode and Dean 2004, CDFG unpublished data). Major assemblages of non-fish aquatic animal species include benthic macroinvertebrates and zooplankton communities.

The distribution and productivity of organisms and aquatic habitats of Shasta Lake are greatly affected by the reservoir's dynamic seasonal surface elevation fluctuations and thermal stratification. The reservoir's flood control, water storage, and water delivery operations typically result in declining water elevations during summer through fall, rising or stable elevations during winter, and rising elevations during spring and sometimes into the early summer, while storing precipitation and snowmelt runoff. During summer, the epilimnion (relatively warm surface layer) is 30 to 50 feet deep and warms up to 80 degrees Fahrenheit (°F). Water temperatures above 68°F favor warm-water fishes such as bass and catfish. Deeper water layers, which include the hypolimnion (cold bottom layer) and the metalimnion (transition zone between epilimnion and the hypolimnion), are cooler and are suitable for cold-water species. Shasta Lake is classified as a cool-water, mesotrophic, monomictic reservoir because it is moderately productive and has one period of mixing each year, although it never completely turns over (Bartholow et al. 2001).

The amount of warm-water and cold-water habitat provided by Shasta Lake is a function of the total storage volume and associated surface area provided by Shasta Lake. This relationship is influenced by variation in the water surface elevation (WSEL) throughout the year. Variation in WSEL is a function of water demand, water quality requirements, and inflow, and WSEL can change based on the water year type. Typically, primary production in reservoirs is associated with storage volumes when all other factors are held constant (Stables et al. 1990). Increased storage and the corresponding increase in surface area results in a greater total biomass and a greater abundance of plankton and fish, because available habitat area is increased.

Within the watersheds surrounding Shasta Lake, there are approximately 1,300 ephemeral, intermittent, and perennial stream channels totaling about 2,903 miles. Aquatic habitats provided by perennial and intermittent streams that are tributaries to Shasta Lake are important to the overall productivity of the cold-water and warm-water fisheries supported by the lake. Most of these tributaries are relatively short and steep and may be classified as confined headwater channels that contribute water, sediment, and organic and inorganic material to Shasta Lake. The lower reaches of the tributaries draining into the reservoir may provide spawning habitat for adfluvial fishes (i.e., fish that spawn in streams, but rear and grow to maturity in lakes) residing in Shasta Lake, as well as stream-resident fishes and other aquatic species that either directly or indirectly support aquatic resources in Shasta Lake.

Upper Sacramento River (Shasta Dam to Red Bluff) The upper Sacramento River flows for approximately 10 miles between Shasta Dam and Keswick Dam and 59 miles between Keswick Dam and RBPP. The river in this reach has cool water temperatures because of regulated releases from Shasta and Keswick dams, and a stable, largely confined channel with little meander. Immediately downstream from Keswick Dam, the river is deeply incised in bedrock with very limited riparian vegetation and limited functioning riparian ecosystems. Near Redding, the river flows into the valley and the floodplain broadens. Historically, this area appeared to have had wide expanses of riparian forests, but much of the river's riparian zone is currently subject to urban encroachment and noxious weed problems. This encroachment becomes quite extensive in the Anderson/Redding area, with homes placed directly in or adjacent to the riparian zone.

Despite net losses of gravel since construction of Shasta Dam, substrates in much of this reach contain gravel needed for spawning by salmonids, mostly derived from the Central Valley Project Improvement Act (CVPIA) gravel augmentation program. This reach provides much of the remaining spawning and rearing habitat of several listed anadromous salmonids. For this reason, it is one of the most sensitive and important stream reaches in the State.

Three water control structures, Keswick Dam, Anderson-Cottonwood Irrigation District (ACID), and RBPP, are located along the Sacramento River in this

reach. The main tributaries to the Sacramento River between Shasta Dam and Red Bluff are Battle, Bear, Clear, Cow, and Cottonwood creeks. The primary land uses along the Sacramento River between Shasta Dam and RBPP are urban, residential, and agricultural.

Before the construction of Shasta Dam, the Sacramento River typically experienced large fluctuations in flow driven by winter storms, with late-summer flows averaging 3,000 cubic feet per second (cfs) or less. These fluctuations and periodic high flows moved large amounts of sediment and gravel out of the mountainous tributaries and down the Sacramento River. The completion of Shasta Dam in 1945 resulted in general dampening of historic high and low flows, reducing the timing, magnitude, and duration of winter floods while maintaining higher summer flows between 7,000 and 13,000 cfs. The annual volume of flow in the Sacramento River continues to vary significantly from year to year. However, average monthly flows following the construction of Shasta Dam no longer exhibit pronounced seasonal winter highs and summer lows. This is primarily because of winter flood control operations that have reduced peak flood flows, and summer releases made for water supply purposes.

Extended Study Area

The extended study area consists of the lower Sacramento River (including major tributaries and floodplain bypasses) and Delta, Trinity River, lower San Joaquin River, and the CVP and SWP service areas. Each of these areas/water bodies is described separately below.

Lower Sacramento River The roughly 300 miles of the lower Sacramento River can be subdivided into distinct reaches. These reaches are discussed separately because of differences in morphology, water temperature regime, and aquatic habitat functions. This section focuses on the reaches of the lower Sacramento River from RBPP to Colusa and from Colusa to the Delta. Each of these reaches is discussed individually along with the main tributaries and floodplain bypasses to the Sacramento River.

Red Bluff Pumping Plant to Colusa In this reach, the Sacramento River functions as a large alluvial river with active meander migration through the valley floor. The river is classified as a meandering river, where stable, straight sections confined by levees or riprap alternate with more sinuous, dynamic sections (Sacramento River Conservation Area Forum 2003). The active channel is fairly wide in some stretches and the river splits into multiple forks at many different locations, creating gravel islands, often with riparian vegetation. Historic bends in the river are visible throughout this reach and appear as scars of the historic channel locations with the riparian corridor and oxbow lakes still present in many locations. The channel remains active and has the potential to migrate in times of high water. Point bars, islands, high and low terraces, instream woody cover, early successional riparian plant growth, and other evidence of river meander and erosion are common in this reach. The channel

takes on varying widths, and aquatic habitats consist of shallow riffles, deep runs, deep pools at the bends, glides in the straight reaches, and shallow vegetated floodplain areas that become inundated during high flows.

Colusa to the Delta The general character of the Sacramento River changes drastically downstream from Colusa from a dynamic and active meandering channel to a confined, narrow channel restricted from migration. While setback levees exist along portions of the river upstream from Colusa, the levees become much narrower along the river edge as the river continues south to the Delta. Surrounding agricultural lands encroach directly adjacent to the levees, which have cut the river off from the majority of its riparian corridor, especially on the eastern side of the river. The majority of the levees in this reach are lined with riprap, allowing the river no erodible substrate. The channel width is fairly uniform and river bends are static as a result of confinement by levees. Therefore, aquatic habitats are fairly homogenous because depth profiles and substrate composition are fairly uniform throughout the reach. Two primary tributaries (i.e., Feather and American rivers) enter the lower Sacramento River in this reach (see additional discussion below). Multiple water diversion structures in this reach move floodwaters into floodplain bypass areas during high-flow events. Primary floodplain bypass areas include the Butte Basin, Sutter Bypass, and Yolo Bypass, all of which are fed by overflow weirs along the Sacramento River (see additional discussion below).

Primary Tributaries to Lower Sacramento River The lower reaches of two primary tributaries (i.e., Feather and American rivers) are included because of the project's potential for the effecting flows and associated flow-related effects on fish species of management concern within these tributaries. However, potential project-related flow changes in these areas are limited because of other influencing factors including operation of upstream CVP and SWP reservoirs (i.e., Oroville and Folsom dams, respectively) and inflows from in-basin tributaries, as well as diversions and flood bypasses.

Lower Feather River Aquatic habitats found in the lower Feather River, downstream from Oroville Dam, vary as the river flows from releases at the California Department of Water Resources (DWR) Oroville Dam facilities down to the confluence with the Sacramento River at Verona. At the upper extent, the approximate 8-mile low-flow (about 600 cfs) section contains mainly riffles and runs, which provide spawning habitat for the majority of Feather River Chinook salmon (*Oncorhynchus tshawytscha*) and steelhead (*O. mykiss*). Also present in the low-flow channel is a series of remnant gravel pit pools/ponds that connect to the main channel. This stretch is fairly confined by levees as it flows through the city of Oroville. From the downstream end of the low-flow channel, the Feather River is fairly active and meanders its way south to Marysville. However, this stretch is bordered by active farmland, which confines the river into an incised channel in certain stretches. Several areas of adjacent farmlands are in the process of being restored to floodplain habitat with setback levees.

Lower American River Flows in the lower American River (downstream from Folsom and Nimbus dams) provide habitat for anadromous and resident fish species. The river is fairly low gradient and is composed of riffle, run, glide, and pool habitats. Dams along the watershed have reduced gravel inputs to the system, but the lower American River contains large gravel bars and forks in many locations, leaving gravel/cobble islands within the channel. The majority of the lower American River is surrounded by the American River Parkway, preserving the surrounding riparian zone. The river channel does not migrate to a large degree because of the geologic composition that has allowed the river to incise deep into sediments, leaving tall cliffs and bluffs adjacent to the river.

Sacramento River Floodplain Bypasses As described above, there are three major floodplain bypasses – Butte Basin, Sutter Bypass, and Yolo Bypass – with a total of 10 overflow structures along the lower Sacramento River (six weirs, three flood relief structures, and an emergency overflow roadway) that provide access to broad, inundated floodplain habitat during wet years.

Unlike other Sacramento River and Delta habitats, floodplains and floodplain bypasses are seasonally dewatered (as high flows recede) during late spring through autumn. This prevents introduced fish species from establishing year round dominance except in perennial water sources (Sommer et al. 2003). Moreover, many of the native fish are adapted to spawn and rear in winter and early spring (Moyle 2002) during the winter flood pulse. Introduced fish typically spawn during late spring through summer when the majority of the floodplain is not available to them.

Butte Basin The Butte Basin lies east of the Sacramento River and extends from the Butte Slough outfall gates near Meridian to Big Chico Creek near Chico Landing. Flood flows are diverted out of the Sacramento River into the Butte Basin and Sutter Bypass via several designated overflow areas (i.e., low points along the east side of the river) that allow high flood flows to exit the Sacramento River channel.

Sutter Bypass The Sutter Bypass is a narrow floodwater bypass conveying Sacramento River flood flows from the Butte Basin and the Tisdale Weir. The bypass area is an expansive land area in Sutter County used mainly for agriculture. In times of high water, Sacramento River water enters the bypass through the Butte Slough outfall and the Tisdale Weir (when the river stage exceeds 45.5 feet) and inundates the bypass with as much as 12 feet of water. The Sutter Bypass, in turn, conveys flows to the lower Sacramento River region at the Fremont Weir near the confluence with the Feather River and into the Sacramento River and the Yolo Bypass (USACE and the Reclamation Board 2002).

Yolo Bypass The Yolo Bypass is an approximate 59,000-acre land area that conveys Sacramento River flood waters around Sacramento during times of high runoff. Flow is diverted from the Sacramento River into the bypass when

the river stage exceeds 33.5 feet (corresponding to 56,000 cfs at Verona). Diversion of the majority of Sacramento River, Sutter Bypass, and Feather River floodwaters to the Yolo Bypass from Fremont Weir controls Sacramento River flood stages at Verona. During large flood events, up to 80 percent of Sacramento River flows are diverted into the bypass.

All six weirs (Moulton, Colusa, Tisdale, Fremont, Sacramento, and Cache Creek) have a fixed-level, concrete overflow section, followed by a concrete, energy-dissipating stilling basin, with a rock and/or concrete erosion blanket across the channel beyond the stilling basin and a pair of training levees that define the weir's flow escape channel. All overflow structures except the Sacramento Weir pass floodwaters by gravity once the river reaches the overflow WSEL. The Sacramento Weir has gates on top of the overflow section that hold back floodwaters until opened manually by DWR's Division of Flood Management.

Lower San Joaquin River The lower San Joaquin River downstream from the Merced River confluence is characterized by a relatively wide (approximately 300 feet) channel with little canopy or overhead vegetation and minimal bank cover. Aquatic habitat in the San Joaquin River is characterized primarily by slow-moving glides and pools, is depositional in nature, and has limited water clarity and habitat diversity. Several fish species use this lower segment of the river to some degree, even if only as a migratory pathway to and from upstream spawning and rearing areas. The lowermost portion of the river also is used by certain fish species (e.g., delta smelt (*Hypomesus transpacificus*)) that make little to no use of areas in the upper segment of the river (see Delta discussion below).

Sacramento-San Joaquin Delta The Delta and San Francisco Bay (Bay) make up the largest estuary on the west coast (EPA 1993). The Delta and Suisun Bay, on the western edge of the Delta, are located at the confluence of the Sacramento and San Joaquin rivers and may be considered to represent the most important, complex, and controversial geographic area for both anadromous and resident fisheries production and distribution of California water resources for numerous beneficial uses (Hanson, pers. comm., 2009). The Delta consists of a large network of channels through which water, nutrients, and aquatic food resources are moved and mixed by tidal action. The Delta is shown in Figure 1-1.

The San Francisco Bay-Delta (Bay-Delta) is a complex estuarine ecosystem, a transition zone between inland sources of freshwater and saltwater from the ocean. Along the salinity gradient extending from the Golden Gate upstream into the central Delta and tributaries, the species composition of the aquatic community changes dramatically, although the basic functional relationships among organisms (e.g., predator-prey) remain similar throughout the system.

The Delta's channels are used to transport water from upstream reservoirs to the south Delta, where Federal and State facilities (C.W. "Bill" Jones Pumping Plant and Harvey O. Banks Delta Pumping Plant, respectively) pump water into CVP and SWP canals, respectively.

Environmental conditions in the Delta depend primarily on the physical structure of Delta channels, inflow volume and source, Delta Cross Channel (DCC) operations, Delta exports and diversions, and tides. The CVP affects Delta conditions primarily through control of upstream storage and diversions, Delta exports and diversions, and DCC operations. These factors also determine outflow and the location of the entrapment zone, which is an area of high organic carbon that is critically important to a number of fish and invertebrate species, as well as to the overall ecology of the Delta and Suisun Bay. In addition to these physical factors, environmental conditions such as water temperature, predation, food production and availability, competition with introduced exotic fish and invertebrate species, and pollutant concentrations all contribute to interactive, cumulative conditions that have substantial effects on Delta fish populations.




Figure 1-1. San Francisco Bay and Sacramento-San Joaquin Delta

Delta habitat is of key importance to fish, as illustrated by the more than 120 fish species that rely on its unique habitat characteristics for one or more of their life stages (EPA 1993). Fish species found in the Delta include anadromous species, as well as freshwater, brackish water, and saltwater species. The Delta provides spawning and nursery habitat for more than 40 resident and anadromous fish species, including delta smelt, Sacramento splittail, American shad, and striped bass. The Delta is also a migration corridor and seasonal rearing habitat for all four runs of Chinook salmon, steelhead, and green sturgeon.

Adult Chinook salmon move through the Delta during most months of the year. Chinook salmon and steelhead juveniles depend on the Delta as transient rearing habitat during their migration to the ocean, and may remain for several months, feeding in marshes, tidal flats, and sloughs. All life stages of striped bass and American shad (*Alosa sapidissima*) are found in the Delta; approximately 45 percent of striped bass (*Morone saxatilis*) spawn in the Delta, as do some American shad. Numerous resident species live in the Delta year-round, including delta smelt, Sacramento splittail (*Pogonichthys macrolepidotus*), and introduced threadfin shad (*Dorosoma petenense*).

Delta inflow and outflow are important for fishes residing primarily in the Delta (e.g., delta smelt, longfin smelt) (USFWS 2008), as well as juveniles of anadromous species (e.g., Chinook salmon) that rear in the Delta before ocean entry. Seasonal Delta inflows affect several key ecological processes, including (1) the migration and transport of various lifestages of resident and anadromous fishes using the Delta; (2) salinity levels at various locations within the Delta, as measured by the location of the salinity isopleths (X2) (i.e., the position in kilometers eastward from the Golden Gate Bridge of the 2 parts per thousand (ppt) near bottom isohaline); and (3) the Delta's primary (phytoplankton) and secondary (zooplankton) production.

The Bay region is predominantly developed for urban and industrial uses. The region contains numerous small streams and reservoirs used primarily for domestic water supply. All anadromous species use these habitats, with the exception of some American shad and striped bass that complete their entire life cycles within the Delta and upstream. The four runs of Chinook salmon and steelhead migrate as adults from the Pacific Ocean, through the Bay and into their natal rivers, while Chinook salmon and steelhead smolts migrate downstream through the Bay on their way to the ocean.

More than 200 fish species, mostly marine, exist in the Bay (Miller and Lea 1972). The Bay is an important nursery area for marine and estuarine species, including bay shrimp (*Cragon* spp.), dungeness crab (*Cancer magister*), northern anchovy (*Engraulis mordax*), Pacific herring (*Clupea harengus*), and English sole (*Pleuronectes vetulus*). The Bay provides a protective, highly productive habitat that enhances early survival and growth of these species.

Delta outflow influences abundance and distribution of fish and invertebrates in the Bay through changes to salinity, currents, nutrient levels, and pollutant concentrations. The response of organisms to outflow depends on species and life stage. The variability in the response of organisms to variable outflow volumes is important in the dynamics of the estuarine community. The effect of Delta outflow on aquatic organisms is determined by its timing, magnitude, and duration. The cause-and-effect relationship between Delta outflow and organism abundance and distribution is complex and often dictated by a chain or web of events rather than by specific, direct effects. Although correlations between flows and organism abundance have been identified, the mechanisms of the relationships are largely unknown. Water residence time in the Bay, determined by tides, local inflow, Delta outflow, and bathymetry, also affects fish species abundance (Smith 1987).

In many segments of the estuary, but particularly in Suisun Bay and the Delta, salinity is controlled by the balance of salt water intrusion from the Bay and freshwater flow from the tributaries to the Delta. By altering the timing and volume of flows, water development has affected salinity patterns in the Delta and in parts of the Bay. Historically, under natural conditions, the Carquinez Strait/Suisun Bay region marked the approximate boundary between salt water and fresh water in the estuary during much of the year. In the late summer and fall of drier years, when Delta outflow was minimal, seawater moved into the Delta from the Bay. Beginning in the 1920s, following several dry years, and because of increased upstream storage and diversions, salinity intrusions moved farther upstream and became more frequent.

Since the 1940s, releases of fresh water from upstream storage facilities have increased Delta outflows during summer and fall. These flows have correspondingly limited the extent of salinity intrusion into the Delta. Reservoir releases have helped to ensure that the salinity of water diverted from the Delta is acceptable during summer and late fall for agricultural, municipal, and industrial uses.

Salinity is an important habitat component in the estuarine environment of the Delta. All estuarine species are assumed to have optimal salinity ranges, and their survival may be affected by the amount of habitat available within the species' optimal salinity range. Because the salinity field in the Bay-Delta is largely controlled by freshwater inflows, Delta outflow may determine the surface area of optimal salinity habitat that is available to the species (Hieb and Baxter 1993, Unger 1994).

The transition area between saline waters within the Bay and freshwater within the rivers, frequently referred to as the low salinity zone, is located within Suisun Bay and the western Delta. The low salinity zone has also been associated with the entrapment zone, a region of the Bay-Delta characterized by higher levels of particulates, higher abundances of several types of organisms, and a turbidity maximum. It is commonly associated with the position of X2,

but actually occurs over a broader range of salinities (Kimmerer 1992). Originally, the primary mechanism responsible was thought to be gravitational circulation, a circulation pattern formed when freshwater flows seaward over a dense, landward-flowing marine tidal current. However, gravitational circulation does not occur in the entrapment zone in all years, nor is it always associated with X2 (Burau et al. 1998). Lateral circulation within the Bay-Delta or chemical flocculation may play a role in the formation of the turbidity maximum of the entrapment zone.

As a consequence of higher levels of particulates, the entrapment zone may be biologically significant to some species. Mixing and circulation in this zone concentrates plankton and other organic material, thus increasing food biomass and production. Larval fish such as striped bass, delta smelt, and longfin smelt may benefit from enhanced food resources. Since about 1987, however, the introduced Asian overbite clam population has cropped much of the primary production in the Bay-Delta and there has been virtually no enhancement of phytoplankton production or biomass in the entrapment zone (CUWA 1994). Although the base of the food chain may not have been enhanced in the entrapment zone during the past decade, this region continues to have relatively high levels of invertebrates and larval fish.

X2 and the entrapment zone are not as closely related as previously believed (Burau et al. 1998), X2 continues to be used as an index of the location of the entrapment zone and area/or of increased biological productivity. Historically, X2 has varied between San Pablo Bay (river kilometer (RK) 50) during high Delta outflow and Rio Vista (RK 100) during low Delta outflow. X2 has typically been located between approximately Honker Bay and Sherman Island (RK 70 to 85). X2 is controlled directly by the volume of Delta outflow, although changes in X2 lag behind changes in outflow. Minor modifications in outflow do not greatly alter X2.

Trinity River Sacramento River flow is augmented in average water years by transfer of up to 1 MAF of Trinity River water through the Clear Creek and Spring Creek tunnels to Keswick Reservoir (Reclamation 2004). Flows in the Trinity River (below Lewiston Dam) are generally cold, providing habitat for anadromous and resident fish species. Aquatic habitats in the river consist of riffle, run, glide, and pool habitats. Fish habitat values have increased in quantity and quality through restoration activities that have taken place over the last several years. Implementation of the Trinity River Restoration Program is expected to further increase the value of the habitat below Lewiston Dam over the next 10 to 15 years (NMFS 2000).

CVP/SWP Service Areas The CVP and SWP service areas contain several highly altered aquatic habitat types, including reservoirs, canals, ditches, and other manmade water conveyance structures/facilities. Agricultural land and urban development are the dominate land uses within these service areas. As a result of all these factors, the aquatic communities that occupy the habitats are

highly adapted to these disturbed environments and are dominated by nonnative species.

1.1.2 Fisheries Resources

This section describes the life history, habitat requirements, and factors that affect the abundance of species selected for the assessment of impacts of the proposed project alternatives. A separate discussion on aquatic macroinvertebrates in the primary and extended study areas is presented after this section.

Primary Study Area

Water bodies within the primary study area contain a large and diverse assemblage of resident and anadromous fish species, including recreationally and commercially important species, and species that are listed as threatened and endangered (Table 1-1).

Table 1-1. Fish Species Known to Occur in Primary Study Area

Common Name	Scientific Name	Distribution Within Primary Study Area		
		Shasta Lake Tributaries	Shasta Lake/ Keswick Reservoir	Sacramento River— Keswick to Red Bluff
Chinook salmon	<i>Oncorhynchus tshawytscha</i>		X	
winter-run				X
spring-run				X
fall-run				X
late fall-run				X
Rainbow trout	<i>Oncorhynchus mykiss</i>	X	X	X
Steelhead trout	<i>Oncorhynchus mykiss</i>			X
Brown trout	<i>Salmo trutta</i>	X	X	X
Green sturgeon	<i>Acipenser medirostris</i>			X
White sturgeon	<i>Acipenser transmontanus</i>	X	X	X
Pacific lamprey	<i>Entosphenus tridentatus</i>			X
Western brook lamprey	<i>Lampetra richardsoni</i>			X
Sacramento sucker	<i>Catostomus occidentalis</i>	X	X	X
Sacramento pikeminnow	<i>Ptychocheilus grandis</i>	X	X	X
Hardhead	<i>Mylopharodon conocephalus</i>	X	X	X
Sacramento blackfish	<i>Orthodon microlepidotus</i>	X	X	

Table 1-1. Fish Species Known to Occur in Primary Study Area (contd.)

Common Name	Scientific Name	Distribution Within Primary Study Area		
		Shasta Lake Tributaries	Shasta Lake/ Keswick Reservoir	Sacramento River— Keswick to Red Bluff
California roach	<i>Lavinia symmetricus</i>	X		X
Speckled dace	<i>Rhinichthys osculus</i>	X	X	
Golden shiner	<i>Notemigonus crysoleucas</i>	X	X	
Carp	<i>Cyprinus carpio</i>	X	X	X
Channel catfish	<i>Ictalurus punctatus</i>	X	X	X
White catfish	<i>Ameiurus catus</i>		X	X
Brown bullhead	<i>Ameiurus nebulosus</i>		X	X
Black bullhead	<i>Ameiurus melas</i>		X	X
Riffle sculpin	<i>Cottus gulosus</i>	X	X	
Prickly sculpin	<i>Cottus asper</i>	X		X
Rough sculpin	<i>Cottus asperrimus</i>	X		
Pit sculpin	<i>Cottus pitensus</i>	X		
Bigeye marbled sculpin	<i>Cottus klamathensis macrops</i>	X		
Largemouth bass	<i>Micropterus salmoides</i>		X	
Smallmouth bass	<i>Micropterus dolomieu</i>	X	X	X
Spotted bass	<i>Micropterus punctulatus</i>	X	X	
Black crappie	<i>Pomoxis nigromaculatus</i>		X	
White crappie	<i>Pomoxis annularis</i>		X	
Bluegill sunfish	<i>Lepomis macrochirus</i>		X	
Green sunfish	<i>Lepomis cyanellus</i>	X	X	
Threadfin shad	<i>Dorosoma petenense</i>		X	
Tule perch	<i>Hysterocarpus traski</i>	X	X	X
Tui chub	<i>Siphateles bicolor</i>	X	X	

Source: Moyle 2002; Reclamation 2004 ; Reclamation 2014

Shasta Lake and Vicinity Shasta Lake fish species include native and nonnative species, which are dominated by mostly introduced warm-water and cold-water species (Weidlein 1971; CDFG, unpublished data) (Table 1-1). Major assemblages of aquatic non-fish animal species include benthic macroinvertebrates and zooplankton communities.

Cold-Water Species Shasta Lake and its tributaries provide very productive habitats for cold-water fish species, which typically prefer or require water temperatures cooler than 70°F. During the cooler months, cold-water species such as rainbow trout, brown trout, and landlocked Chinook salmon may be found throughout the lake; however, these species do not spawn in the lake, preferring to spawn in tributary streams. During summer, these cold-water species may be found rearing in association with the cold, deep hypolimnion and metalimnion layers within the reservoir, although the fish may make frequent forays into the epilimnion to feed on small prey fish and return to cooler depths to digest their prey (Finnell and Reed 1969, Koski and Johnson 2002, Moyle 2002, Quinn 2005).

Native species such as white sturgeon, hardhead, riffle sculpin, Sacramento sucker, and Sacramento pikeminnow tend to reside in cooler water strata in the reservoir and in and near tributary inflows (Moyle 2002). Trout may also congregate near the mouths of the reservoir's tributaries, including the upper Sacramento River, McCloud River, Pit River, and Squaw Creek, at various times of the year for various purposes, including thermal refuge, foraging, and spawning, when conditions are favorable for these species.

Hatchery- and pen-reared trout and salmon are stocked in Shasta Lake several times each year to support the sport fishery. About 60,000 pounds of juvenile rainbow trout and about 50,000 subcatchable Chinook salmon are planted annually (Baumgartner, pers. comm., 2008).

Climate conditions and reservoir storage volume are the two most influential factors affecting cold-water habitat and primary productivity in Shasta Lake (Bartholow et al. 2001). Cold-water habitat provided by Shasta Lake is a function of the total storage and associated surface area provided by Shasta Lake. This relationship is influenced by variation in the WSEL throughout the year. Variation in WSEL is a function of water demand, water quality requirements, and inflow, and WSEL can change based on the water year type. Typically, primary production in reservoirs is associated with storage volumes when all other factors are held constant (Stables et al. 1990). Increased storage and the corresponding increase in surface area results in a greater total biomass and a greater abundance of plankton and fish, because available habitat area is increased.

Warm-Water Species The warm-water fish habitats of Shasta Lake occupy two ecological zones: the littoral (shoreline/rocky/vegetated) and the pelagic (open water) zones. The littoral zone lies along the reservoir shoreline down to the maximum depth of light penetration on the reservoir bottom, and supports populations of spotted bass, smallmouth bass, largemouth bass, black crappie, bluegill, channel catfish, and other warm-water species.

The upper, surface layer of the pelagic zone is the principal plankton-producing region of the reservoir. Plankton comprises the base of the food web for most of the reservoir's fish populations. Operation of the Shasta Dam temperature control device (TCD), which helps conserve the reservoir's cold-water pool by accessing warmer water for storage releases in winter, spring, and early summer, may reduce zooplankton biomass in the epilimnion. However, operations of the TCD may result in some increased plankton production at deeper levels as a result of a slight warming of the hypolimnetic layers within the reservoir during fall (Bartholow et al. 2001).

Warm-water species, such as largemouth bass, smallmouth bass, spotted bass, and several sunfishes, were introduced into Shasta Lake and have become well established with naturally sustaining populations. Spotted bass are currently the dominant warm-water species in Shasta Lake (S. Baumgartner, pers. comm.,

2006). These warm-water fishes feed primarily on invertebrates while young and become predaceous on other fishes, including engaging in some cannibalism, as they grow. In Shasta Lake, threadfin shad and crayfish and other invertebrates are most abundant in the diets of these fish (Saito et al. 2001). Spawning activity usually begins during late March or April when temperatures rise to around 60°F. Males generally build the nests in sand, fine gravel, rubble, or debris-covered bottoms at depths between 1 and 20 feet, which varies by species. Spotted bass and catfishes typically spawn at greater depths than the other warm-water species in Shasta Lake. Eggs generally hatch in 3 to 5 days at the predominant springtime water temperatures in Shasta Lake, and males guard the eggs and larvae for up to 4 weeks (Moyle 2002). Fry and juveniles disperse into shallow water and prefer areas with vegetation and large rubble as protective cover from predators (Moyle 2002, Ratcliff 2006).

The primary factors affecting warm-water fish abundance and production in Shasta Lake include seasonal reservoir fluctuations, availability of high-quality littoral habitat, and annual climate variations (Ratcliff 2006). The effect of sport fishery harvests on Shasta Lake warm-water fish populations is not well understood; it is believed to be small because of catch-and-release practices, although it is generally thought that overfishing of naturally reproducing populations by sport fisheries seldom limits fish abundance (Moyle 2002).

The magnitude and timing of reservoir level fluctuations, associated shoreline erosion, and suppression of shoreline and emergent vegetation are thought to generally be the most significant factors affecting warm-water fish production in reservoirs, including Shasta Lake (Moyle 2002, Ratcliff 2006). Water level variations influence physical, chemical, and biological processes, which in turn affect fish populations. Reservoir drawdowns reduce water depths and influence thermal stratification and the resulting temperature, dissolved oxygen (DO), and water chemistry profiles.

The typical seasonality of reservoir fluctuations on Shasta Lake can affect year-to-year reproductive success of littoral-spawning fishes, especially the black bass species, by influencing nesting behavior (e.g., abandonment of nests) and dewatering of nests containing eggs in years when reservoir levels decline during the spring and early summer months. Under these same conditions, juveniles may be forced to move to areas with less protection from predation or lower food production. In years when the reservoir rises rapidly and/or extensively during spring and early summer, submergence of active bass nests by more than 15 to 20 feet often results in high egg mortality (Stuber et al. 1982, Lee 1999, Moyle 2002).

Shoreline and littoral vegetation are important warm-water fish habitat components for sustainable fisheries (Ratcliff 2006). Structural diversity (e.g., submerged trees, brush, rock, boulders, and rubble) provides shelter and feeding areas for fish. During construction of the reservoir, many trees and brush fields were cleared before inundation. Portions of the Pit River and Squaw Creek arms

were not cleared, as evidenced by the large number of inundated trees observable in certain areas. Clearing efforts reduced the potential structural diversity of the inundated habitat. Vegetative clearing in many reservoirs has resulted in rocks, boulders, and man-made features (e.g., bridge pilings, riprap, marinas) being the only structural habitat features available, especially for bass and other warm-water fishes.

Annual reservoir fluctuations create highly variable conditions for establishment and maintenance of shoreline and littoral-zone vegetation and aquatic invertebrate communities that subsequently impose limitations on warm-water fish production. Exposed shoreline reservoir areas generally require 3 to 4 years to reestablish terrestrial vegetation. The absence of established, rooted aquatic vegetation is a common aquatic habitat factor that limits populations and fishery production for many fish species in reservoirs (Ploskey 1986, Moyle 2002).

The Shasta-Trinity National Forest (STNF), in cooperation with other Federal and State agencies and local nongovernmental organizations (NGO), has implemented a habitat improvement program at Shasta Lake. The objective of this program is to increase cover for warm-water fish. As the fishery management agency for Shasta Lake, the California Department of Fish and Wildlife (CDFW formerly known as the California Department of Fish and Game [CDFG]) prepared a Draft Management Plan for Shasta Lake in 1991. This plan, which has not been finalized, acknowledges the benefit to warm-water fish of structural enhancement projects.

STNF, CDFW, and NGOs have used a variety of materials and techniques to construct structural enhancements (e.g., willow planting, brush structures) to provide warm-water fish habitat within the drawdown zone of Shasta Lake. The materials and techniques have varied because of differences in funding, available materials, site conditions (reservoir levels), longevity, and desired outcome.

According to STNF aquatic biologists, brush structures constructed from whiteleaf manzanita (*Arctostaphylos manzanita*) have been the STNF's preferred means of structural enhancement since about 1990. These structures have been constructed in areas where manzanita is available near the shoreline, typically in manner that provides varying degree of structural habitat as water levels change over time. The biologists have indicated that these structures have typically resulted in a threefold to tenfold increase in the abundance of warm-water fish in the treated areas (Joe Zustak, pers. comm. 2007).

Tributary Species The lower reaches of the tributaries draining to the reservoir provide spawning habitat for adfluvial fishes (i.e., fish that spawn in streams, but rear and grow to maturity in lakes) residing in Shasta Lake, as well as stream-resident fishes, with rainbow trout the principal game species. Accessible and suitable cold-water fish spawning habitat, including appropriate seasonal flows, depths, and gravel substrates, was observed in only 5 percent of

intermittent and nearly 70 percent of perennial tributaries to Shasta Lake surveyed in 2011–12 (Table 1-2).

Most native fish species found in Shasta Lake may also inhabit the lower reaches of the tributaries. Several tributaries to Shasta Lake (e.g., Little Squaw Creek, Little Backbone Creek) have been subjected to discharge from abandoned upslope copper mines. The Shasta Lake West Watershed Analysis (Bachmann 2000) suggests that these creeks are “biologically dead” as a result of acid mine discharge from these mines. This watershed analysis also stated that “fish kills” have occurred in Shasta Lake in the vicinity of such tributaries during high runoff conditions. No fish were observed during 2012 in tributaries known to be affected by a legacy of mining and acidic, metal-laden mine drainage. These tributaries include Little Squaw Creek and Little Backbone Creek, both located in the watershed to the immediate northwest of Shasta Dam, and Town Creek, located near the Bully Hill Mine in the Squaw Creek arm (Reclamation 2014).

The four main tributaries to Shasta Lake are the Sacramento River, McCloud River, Squaw Creek, and Pit River, which are renowned for their high-quality recreational trout fisheries. Each of these streams drains considerable watershed areas comprising mixed conifer forests in the reaches above Shasta Lake. With the exception of the Pit River, which has a series of hydroelectric project dams that begin immediately upstream from Shasta Lake, each of these tributaries has more than 30 miles of high-quality, fish-bearing riverine habitat between Shasta Lake and upstream dams on the Sacramento and McCloud rivers and steep headwater reaches on Squaw Creek.

For the most part, land use along the main Shasta Lake tributaries upstream from the reservoir is a mix of Federal and privately managed forest and timberlands and except for sparse residential development, several small municipalities, and the hydropower projects on the Pit, McCloud, and Sacramento rivers, much of the area is lightly developed. The Sacramento River above Shasta Lake is paralleled by a major interstate highway and railroad transportation corridor. In July 1991, a railroad accident spilled 19,000 gallons of the fumigant pesticide metam sodium into the Sacramento River near the town of Dunsmuir, approximately 35 stream miles upstream from Shasta Lake. Metam sodium is highly toxic and killed aquatic and riparian vegetation, aquatic macroinvertebrates, and fish and amphibians along the entire length of the river to Shasta Lake, where a massive chemical containment and neutralization effort was mounted. Ecological recovery efforts were implemented shortly after this spill incident and populations of fish, aquatic macroinvertebrates, and the vegetation adjacent to the stream have attained levels that appear to be in a natural dynamic equilibrium consistent with full recovery, although some amphibian and mollusk population remained depressed at least 15 years later (Cantara Trustee Council 2007).

In addition to the four primary tributaries, there are 1,232 intermittent and perennial stream channels totaling about 2,962 miles of channel that contribute seasonal or year round flows to Shasta Lake. Most of these channels are relatively short and steep and may be classified as confined headwater channels that contribute water, sediment, and organic and inorganic material to Shasta Lake. Many (64 percent) of these channels are intermittent and have stream slopes greater than 10 percent (mean gradient of 27 percent). About 14 percent, or 154, of the stream channels are perennial, with slopes of less than 7 percent. In the Pacific coast and Cascade ranges, stream channels with gradients up to about 4 to 7 percent and possessing sufficient flows typically exhibit a good potential to support habitation by fish and other aquatic organisms, although steeper slopes do not necessarily, in and of themselves, preclude habitation by fish, particularly trout, sculpins, and dace (Naiman 1998; Reeves et al. 1998). About 79 percent of the tributaries with good fish-bearing potential in the study area occur on the Sacramento River, McCloud River, Squaw Creek, and Pit River arms.

Aquatic habitat for resident and adfluvial fishes is generally limited in the intermittent tributaries to Shasta Lake because a large percentage (92 percent) of these channels do not possess suitable hydrologic conditions (i.e., sufficient duration and amount of discharge) and/or are too steep to provide accessible habitat, even seasonally, for fish (Reclamation 2014). The gradient of most of these tributaries rapidly increases upstream from the shoreline, and natural barriers to fish are common. These barriers are most often created by cascades, waterfalls, and steep reaches of stream channel (i.e., greater than 7 percent slope) that are more than one-quarter mile in length. Stream channel data generated from field inventories and analysis using Reclamation's geographic information system (GIS) Digital Elevation Model (DEM) indicate that most barriers on the perennial tributaries occur near the reservoir. Fifty-four percent of all of the intermittent (both non-fish bearing and potentially fish-bearing) and 30 percent of perennial tributaries contained partial or complete barriers to fish migration within stream reaches occurring in the 1,070 feet and 1,090 feet mean sea level (msl) elevation range (Table 1-3). Estimated percent occurrence of partial and complete fish passage barriers in Shasta Lake tributaries within the projected varial zones of reservoir enlargement alternatives of 6 feet (1,070–1,076 feet msl), 12 feet (1,077–1,082 feet msl), and 18 feet (1,083–1,090 feet msl), based on streams surveyed in 2011–12 is shown in (Table 1-4).

Aquatic habitat for resident and adfluvial fishes in intermittent tributaries to Shasta Lake is generally limited, whereas the aquatic habitat composition in perennial tributaries is more diverse across all elevation strata (Reclamation 2014). Two percent of intermittent and 87 percent of perennial tributaries to Shasta Lake sampled in 2011–12 were found to be inhabited by fish (Table 1-5). Only cold-water species (trout) were observed in intermittent streams during periods of surface flow and in isolated pools after cessation of flow. Cold-water species inhabited 83 percent and warm-water species inhabited 48 percent of the sampled perennial tributaries (Reclamation 2014). Warm-water species were

Table 1-3. Frequency of Occurrence of Shasta Lake Tributaries with Fish Passage Impediments, by Elevation Range, Among Streams Surveyed in 2012

Lake Arm	No. Streams Sampled		Number of Streams with Fish Passage Impediments by Elevation Range					
	Total	With Passage Impediments	1,070–1,076 ft msl		1,077–1,082 ft msl		1,083–1,090 ft msl	
			Partial	Complete	Partial	Complete	Partial	Complete
Intermittent (Non Fish-Bearing)								
Main Body	12	10	1	2	2	0	0	5
Big Backbone	3	3	0	1	0	2	0	0
Sacramento River	20	10	1	3	0	2	0	4
McCloud River	13	6	1	4	0	0	0	1
Squaw Creek	12	5	0	3	0	2	0	0
Pit River	29	17	1	6	0	4	0	6
Total	89	51	4	19	2	10	0	16
Intermittent (Potentially Fish-Bearing)								
Main Body	1	1	0	0	1	0	0	0
Big Backbone	2	0	0	0	0	0	0	0
Sacramento River	5	1	0	0	0	0	0	1
McCloud River	1	0	0	0	0	0	0	0
Squaw Creek	1	0	0	0	0	0	0	0
Pit River	10	6	3	0	0	1	0	2
Total	20	8	3	0	1	1	0	3
Perennial								
Main Body	2	2	2	0	1	0	0	0
Big Backbone	2	1	0	0	1	0	0	0
Sacramento River	7	1	1	0	1	0	0	1
McCloud River	5	2	0	1	1	1	0	1
Squaw Creek	3	1	1	0	0	0	0	0
Pit River	4	0	0	0	0	0	0	0
Total	23	7	4	1	4	1	0	2

Key:
msl = mean sea level
ft = foot

Table 1-4. Estimated Percent Occurrence of Partial and Complete Fish Passage Barriers in Shasta Lake Tributaries Within the Projected Varial Zones of Reservoir Enlargement Alternatives^a

Stream Hydrology	Lake Arm	Streams Sampled		Lakewide Total Number of Streams	Percent of Streams with Barriers		
		No.	No. Potential Fish- bearing		1,070- 1,076 ft msl	1,077- 1,082 ft msl	1,083- 1,090 ft msl
Intermittent	Main Body	13	1	134	23%	8%	38%
	Big Backbone	5	2	51	20%	40%	0%
	Sacramento River	25	5	247	20%	8%	24%
	McCloud River	14	1	137	21%	14%	7%
	Squaw Creek	13	1	128	23%	15%	0%
	Pit River	39	10	381	28%	13%	18%
	Total	109	20	1,078	24%	13%	17%
Perennial	Main Body	2	2	13	100%	50%	0%
	Big Backbone	2	2	13	0%	50%	0%
	Sacramento River	7	7	55	14%	14%	14%
	McCloud River	5	5	29	20%	40%	20%
	Squaw Creek	3	3	26	33%	0%	0%
	Pit River	4	4	18	0%	0%	0%
	Total	23	23	154	22%	22%	9%

Note:

^a Based on a sample of streams surveyed in 2011–12. A 6-foot raise corresponds to an elevation of 1,070–1,076 feet (msl), an 8-foot raise corresponds to an elevation of 1,077–1,082 feet (msl), and an 18-foot raise corresponds to an elevation of 1,083–1,090 feet (msl).

Key:

msl = mean sea level

ft = foot

Table 1-5. Frequency and Proportion of Fish-Bearing Tributaries to Shasta Lake Surveyed in 2012

Stream Hydrology	Lake Arm	No. of Streams Sampled					Percent of Streams		
		Total	With Fish	Without Fish	With Warm-Water Fish	With Cold-Water Fish	Fish-Bearing	Warm-Water Fish	Cold-Water Fish
Intermittent	Main Body	13	0	13	0	0	0%	0%	0%
	Big Backbone	5	0	5	0	0	0%	0%	0%
	Sacramento River	25	0	25	0	0	0%	0%	0%
	McCloud River	14	0	14	0	0	0%	0%	0%
	Squaw Creek	13	0	13	0	0	0%	0%	0%
	Pit River	39	2	37	0	2	5%	0%	5%
	Total	109	2	107	0	2	2%	0%	2%
Perennial ^a	Main Body	2	1	1	1	0	50%	50%	0%
	Big Backbone	2	2	0	1	2	100%	50%	100%
	Sacramento River	7	6	1	3	6	86%	43%	86%
	McCloud River	5	5	0	2	5	100%	40%	100%
	Squaw Creek	3	2	1	1	2	67%	33%	67%
	Pit River	4	4	0	3	4	100%	75%	100%
	Total	23	20	3	11	19	87%	48%	83%

Note:

^a Includes recent documented occurrences of fish in the Sacramento, Pit, and McCloud Rivers that were not sampled during 2012 tributary surveys.

Table 1-6. Frequency and Proportion of Shasta Lake Tributaries Inhabited by Hardhead^a and Foothill Yellow-Legged Frog^b Based on Streams Surveyed in 2011 to 2012

Stream Hydrology	Lake Arm	No. of Streams Sampled				Percent of Streams Sampled		
		Total	With Hardhead	With FYLF	Total No. with Special-Status Species	With Hardhead	With FYLF	Overall Percent with SS species
Intermittent	Main Body	13	0	0	0	0%	0%	0%
	Big Backbone	5	0	0	0	0%	0%	0%
	Sacramento River	25	0	0	0	0%	0%	0%
	McCloud River	14	0	0	0	0%	0%	0%
	Squaw Creek	13	0	0	0	0%	0%	0%
	Pit River	39	0	0	0	0%	0%	0%
	Total	109	0	0	0	0%	0%	0%
Perennial	Main Body	2	0	0	0	0%	0%	0%
	Big Backbone	2	0	0	0	0%	0%	0%
	Sacramento River	7	1	4	5	14%	57%	71%
	McCloud River	5	1	2	3	20%	40%	60%
	Squaw Creek	3	0	1	1	0%	33%	33%
	Pit River	4	1	2	3	25%	50%	75%
	Total	23	3	9	12	13%	39%	52%

Notes:

^a Hardhead is a USFS sensitive species.

^b Foothill yellow-legged frog is a USFS and CDFW sensitive species.

Key:

CDFW = California Department of Fish and Wildlife

FYLF = Foothill yellow-legged frog

SS = special-status

USFS = U.S. Forest Service

Upper Sacramento River (Shasta Dam to Red Bluff) The upper Sacramento River (Shasta Dam to Red Bluff) provides vital fish spawning, rearing, and/or migratory habitat for a diverse assemblage of native and nonnative species. Native species present in this reach of the river can be separated into anadromous (i.e., species that spawn in freshwater after migrating as adults from marine habitat) and resident species. Native anadromous species include four runs of Chinook salmon, steelhead trout, green and white sturgeon (*Acipenser medirostris* and *A. transmontanus*), and Pacific lamprey (*Lampetra tridentata*). Native resident species include Sacramento pikeminnow (*Ptychocheilus grandis*), Sacramento splittail, Sacramento sucker (*Catostomus occidentalis*), hardhead (*Mylopharodon conocephalus*), California roach (*Lavinia symmetricus*), and rainbow trout (*O. mykiss*). Nonnative resident

species include largemouth bass (*Micropterus salmoides*), smallmouth bass (*Micropterus dolomieu*), white and black crappie (*Pomoxis annularis* and *P. nigromaculatus*), channel catfish (*Ictalurus punctatus*), white catfish (*Ameiurus catus*), brown bullhead (*Ictalurus nebulosus*), bluegill (*Lepomis macrochirus*), green sunfish (*Lepomis cyanellus*), and golden shiner (*Notemigonus crysoleucas*).

Keswick Reservoir The U.S. Fish and Wildlife Service (USFWS) conducts a propagation and captive broodstock program for endangered winter-run Chinook salmon at the Livingston Stone National Fish Hatchery located at the base of Shasta Dam on the Sacramento River upstream from Keswick Reservoir. The program consists of collecting adult winter-run Chinook salmon from the mainstem Sacramento River, holding and spawning the adults, rearing the juveniles in the hatchery environment, then releasing them back into the mainstem Sacramento River downstream from Keswick Dam. The overriding goal of the programs is to supplement the endangered population and provide an “insurance policy” against extinction. The propagation program (initiated in 1989), and the captive broodstock program (initiated in 1991) are recognized in the Sacramento River winter-run Chinook salmon Recovery Plan (National Marine Fisheries Service (NMFS) 2014) for this endangered species. Water is supplied to the hatchery from Shasta Dam.

Keswick Reservoir is operated by Reclamation as a reregulating facility. Levels in Keswick Reservoir are subject to operational changes at Whiskeytown and Shasta lakes. The reservoir provides habitat for a variety of aquatic organisms, including native and nonnative fish. Table 1-1 includes the fish species known to occur in Keswick Reservoir. The aquatic habitat in the upper reach of Keswick reservoir is mostly riverine; in the lower half of the reservoir, it is slow current, run-of-the-river habitat. In addition to water released from Shasta Dam and Whiskeytown Lake, this reservoir is the recipient of water and sediment from Spring Creek, emanating from the Iron Mountain Mine. Additional information on the relationship between Spring Creek and Keswick Reservoir is provided in Chapter 9 of the Environmental Impact Statement.

Extended Study Area

Lower Sacramento River and Delta The extended study area includes the middle and lower Sacramento River, tributaries, Delta, and CVP and SWP water service areas. Like the primary study area, habitats in the extended study area also provide vital fish spawning, rearing, and/or migratory habitat for a diverse assemblage of native and nonnative species, many of which are the same as those found in the primary study area (Table 1-7).

Table 1-7. Central Valley Fish Species Potentially Affected by Project Alternatives

Common Name	Scientific Name	Distribution
Native Species		
Hitch	<i>Lavinia exilicauda</i>	Central Valley rivers; Delta
Blackfish	<i>Orthodon microlepidotus</i>	Central Valley rivers; Delta
California roach	<i>Lavinia symmetricus sp.</i>	Central Valley rivers; Delta
Hardhead	<i>Mylopharodon conocephalus</i>	Central Valley rivers; Delta
Sacramento splittail	<i>Pogonichthys macrolepidotus</i>	Central Valley rivers; Delta
Pikeminnow	<i>Ptychocheilus grandis</i>	Central Valley rivers; Delta
Sacramento sucker	<i>Catostomus occidentalis</i>	Central Valley rivers; Delta
Delta smelt	<i>Hypomesus transpacificus</i>	Delta; San Francisco Bay
Longfin smelt	<i>Spirinchus thaleichthys</i>	Delta; San Francisco Bay
Steelhead/rainbow trout	<i>Oncorhynchus mykiss</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Threespine stickleback	<i>Gasterosteus aculeatus</i>	Central Valley rivers; Delta
Prickly sculpin	<i>Cottus asper</i>	Central Valley rivers; Delta
Tule perch	<i>Hysteroecarpus traski</i>	Central Valley rivers; Delta; San Francisco Bay
White sturgeon	<i>Acipenser transmontanus</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Green sturgeon	<i>Acipenser medirostris</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Introduced Species		
American shad	<i>Alosa sapidissima</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Threadfin shad	<i>Dorosoma petenense</i>	Central Valley rivers; Delta
Goldfish	<i>Carassius auratus</i>	Central Valley rivers; Delta
Red shiner	<i>Cyprinella lutrensis</i>	Central Valley rivers; Delta
Carp	<i>Cyprinus carpio</i>	Central Valley rivers; Delta
Golden shiner	<i>Notemigonus chrysoleucas</i>	Central Valley rivers; Delta
Rosyface shiner	<i>Notropis rubellus</i>	Central Valley rivers; Delta
Fathead minnow	<i>Pimephales promelas</i>	Central Valley rivers; Delta
White catfish	<i>Ameiurus catus</i>	Central Valley rivers; Delta
Black bullhead	<i>Ameiurus melas</i>	Central Valley rivers; Delta
Channel catfish	<i>Ictalurus punctatus</i>	Central Valley rivers; Delta
Wakasagi	<i>Hypomesus nipponensis</i>	Delta; San Francisco Bay
Western mosquitofish	<i>Gambusia affinis</i>	Central Valley rivers; Delta
Inland silverside	<i>Menidia beryllina</i>	Central Valley rivers; Delta; San Francisco Bay
Striped bass	<i>Morone saxatilis</i>	Central Valley rivers; Delta; San Francisco Bay; Pacific Ocean
Bluegill	<i>Lepomis macrochirus</i>	Central Valley rivers; Delta
Redear sunfish	<i>Lepomis microlophus</i>	Central Valley rivers; Delta
Smallmouth bass	<i>Micropterus dolomieu</i>	Central Valley rivers; Delta
Largemouth bass	<i>Micropterus salmoides</i>	Central Valley rivers; Delta
White crappie	<i>Pomoxis annularis</i>	Central Valley rivers; Delta
Black crappie	<i>Pomoxis nigromaculatus</i>	Central Valley rivers; Delta
Bigscale logperch	<i>Percina macrolepida</i>	Delta; San Francisco Bay
Yellowfin goby	<i>Acanthogobius flavimanus</i>	Delta; San Francisco Bay
Shimofuri goby	<i>Tridentiger bifasciatus</i>	Delta; San Francisco Bay
Chameleon goby	<i>Tridentiger trigonocephalus</i>	Delta; San Francisco Bay

Source: Moyle 2002, California Department of Fish and Wildlife unpublished data

Key:

Delta = Sacramento-San Joaquin Delta

Trinity River

The Trinity River provides habitat for Southern Oregon/Northern California Coast (SONCC) Coho salmon (*Oncorhynchus kisutch*), SONCC Chinook salmon, Klamath Mountains Province (KMP) steelhead, green sturgeon, white sturgeon, Pacific lamprey, resident rainbow trout, speckled dace, three-spine stickleback, Klamath small scale sucker (*Catostomus rimiculus*), prickly sculpin, and riffle sculpin (*Cottus gulosus*), brook trout (*Salvelinus fontinalis*), brown trout (*Salmo trutta*) American shad, brown bullhead, golden shiner, and green sunfish. Coho salmon and KMP steelhead are included in this discussion because they are special-status species and CVP and SWP operations in response to changes at Shasta Dam have the potential to affect Trinity River flows.

CVP/SWP Service Areas The CVP and SWP water service areas contain several highly altered aquatic habitat types, including reservoirs, canals, ditches and other manmade water conveyance structures/facilities. Agricultural land and urban development are the dominant land uses within these service areas. As a result of all these factors, the aquatic communities that occupy the habitats are highly adapted to these disturbed environments and are dominated by nonnative species, some of which are detrimental to survival of native species.

Special-Status Species

Special-status fish species are legally protected or are otherwise considered sensitive by Federal, State, or local resource conservation agencies and organizations. Special-status fish species addressed in this section include:

- Species listed as threatened or endangered under the Federal Endangered Species Act (ESA) or California Endangered Species Act (CESA).
- Species identified by USFWS, NMFS, or CDFW as species of special concern.
- Species fully protected in California under the California Fish and Game Code.
- Species identified as priorities for recovery under the CALFED Bay-Delta Program (CALFED) Multi-Species Conservation Strategy (CALFED 2000).
- Species considered sensitive or endemic by the U.S. Forest Service (USFS).
- Species considered a survey and manage species by USFS.

A total of nine special-status fish species occur or have the potential to occur in the primary and extended study areas and are described below (see also Table 1-

1). Of the nine species, Central Valley steelhead distinct population segment (DPS), Sacramento River winter-run evolutionarily significant unit (ESU), Central Valley spring-run Chinook salmon ESU, Southern DPS of North American green sturgeon, and delta smelt are Federally listed as threatened or endangered species. USFWS delisted Sacramento splittail from its Federally listed-as-threatened status on September 22, 2003. NMFS determined that listing is not warranted for Central Valley fall-/late fall-run Chinook salmon. However, it is still designated as a Species of Concern because of concerns over specific risk factors. The two remaining species (hardhead and Sacramento perch) are considered Species of Special Concern by CDFW and/or Federal Species of Concern by USFWS. Brief descriptions follow for the special-status species with potential to occur in the primary and extended study areas.

Fish Species of Primary Management Concern

Evaluating potential project alternative-related impacts on fish and aquatic resources requires an understanding of fish species' life histories and life-stage-specific environmental/habitat requirements. Therefore, this information is provided below for fish species of primary management concern that occur within the primary and extended study areas. Species of primary management concern include special-status species likely to occur in the potentially affected portions of the Sacramento River and tributaries and Delta (e.g., Chinook salmon, steelhead, green sturgeon, delta smelt, longfin smelt, Sacramento splittail, hardhead) and species that are recreationally and/or commercially important (e.g., striped bass).

Because these species collectively represent a diversity of life histories and environmental/habitat requirements, and because they are among the most sensitive to environmental perturbation, the findings from assessments made for these species can be effectively used to make inferences to other fish species using the primary and extended study areas. Species of primary management concern with the greatest potential to be affected by implementation of the proposed project alternatives are discussed below. The seasonal timing of important life stages for these species in the study areas is presented in Table 1-8.

Table 1-8. Life History and Distributions of Evaluation Fish Life Stages in Primary and Extended Study Areas

Life Stage/Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Steelhead												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Fall-run Chinook salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Late fall-run Chinook salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Winter-run Chinook salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Spring-run Chinook salmon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Green sturgeon												
Adult migration												
Spawning												
Egg incubation												
Rearing/emigration												
Delta smelt												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Estuarine rearing												
Longfin smelt												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Estuarine rearing												
Sacramento splittail												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												

Table 1-8. Life History and Distributions of Evaluation Fish Life Stages in Primary and Extended Study Areas (contd.)

Life Stage/Location	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Hardhead												
Adult foraging and spawning												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												
Striped Bass												
Adult migration												
Spawning												
Larvae and juvenile rearing												
Adult and juvenile rearing												

Sources: Vogel and Marine 1991, Moyle 2002, Wang 1986, National Marine Fisheries Service 2005

Key:

= period of potential occurrence

Central Valley Steelhead On March 19, 1998, naturally spawned Central Valley steelhead were listed as threatened by NMFS (63 Federal Register (FR) 13347, March 19, 1998). The Central Valley ESU includes all naturally spawned populations of steelhead (and their progeny) in the Sacramento and San Joaquin Rivers and their tributaries. Resident rainbow trout were previously included as part of the protected fish, but in January 2006, NMFS directed that only the anadromous form should be listed as threatened, and the resident form did not warrant listing (71 FR 834, January 5, 2006).

The original critical habitat designation for the Central Valley steelhead was withdrawn pending review. The consent decree (U.S. District Court of the District of Columbia Civil Action No. 00-2799 CKK) resulted in the withdrawal of the critical habitat designation for this ESU. On December 10, 2004, NMFS published a new proposal to designate critical habitat for Central Valley steelhead that includes the lower Feather River; Battle, Cottonwood, Antelope, Mill, Deer, Big Chico, and Butte creeks; Sacramento, Yuba, American, Cosumnes, Mokelumne, Calaveras, San Joaquin, Merced, Tuolumne, and Stanislaus rivers; and the Delta. The final designation for Central Valley steelhead critical habitat was published on September 2, 2005, and was in effect on January 2, 2006 (70 FR 52488, September 2, 2005).

In July 2014, NMFS published the Recovery Plan for Central Valley steelhead, which identifies recovery goals, objectives, and criteria, as well as proposed management actions aimed at bringing the populations to a point at which they can be delisted.

Central Valley steelhead historically migrated upstream into the high gradient upper reaches of Central Valley streams and rivers for spawning and juvenile

rearing. Construction of dams and impoundments on the majority of Central Valley rivers has created impassable barriers to upstream migration and substantially reduced the geographic distribution of steelhead. Although quantitative estimates of the number of adult steelhead returning to Central Valley streams to spawn are not available, anecdotal information and observations indicate that population abundance is low (NMFS 1996). Steelhead distribution is currently restricted to the mainstem Sacramento River downstream from Keswick Dam, the Feather River downstream from Oroville Dam, the Yuba River downstream from Englebright Dam, the American River downstream from Nimbus Dam, the Mokelumne River downstream from Comanche Dam, Cosumnes River, and a number of smaller tributaries to the Sacramento River system, Delta, and the Bay. Steelhead have also been reported from tributaries to the San Joaquin River; however, the status of these populations is under investigation.

The Central Valley steelhead population is composed of both naturally spawning steelhead and steelhead produced in hatcheries. NMFS is continuing to evaluate the status of steelhead and has recently finalized a recovery plan for the species.

Adult steelhead migrate upstream during fall and winter (September through approximately February) with steelhead migration into the upper Sacramento River typically occurring during fall, and adults migrating into lower tributaries typically during late fall and winter. Steelhead spawn in areas characterized by clean spawning gravels, cold-water temperatures, and moderately high velocity. Spawning typically occurs during winter and spring (December through April) with the majority of spawning activity occurring during January and March. Unlike Chinook salmon, which die after spawning, adult steelhead may migrate downstream after spawning and return to spawn in subsequent years.

Steelhead spawn by creating a depression in the spawning gravels where eggs are deposited and fertilized (redd). Steelhead require relatively clean, cool (less than 57°F (13.9 degrees Celsius (°C))) water in which to spawn successfully. The eggs hatch anywhere from 19 to 80 days after spawning, depending on water temperature (warmer temperatures result in faster hatching times), and the young remain in the gravel for several weeks before emerging as fry (Raleigh et al. 1984). The young steelhead emerge from the gravel redd as fry, and rear in the stream system, foraging on insects for 1 to 2 years or longer before migrating to the ocean.

Juvenile steelhead undergo a physiological transformation (i.e., smoltification) that allows the juvenile steelhead to migrate from the freshwater rearing areas downstream to coastal marine waters. Downstream migration of steelhead smolts typically occurs during late winter and early spring, (January through May), although based on salvage data at the Federal and State pumping plants in the Delta, the peak months for emigration appear to be March and April in most years. The seasonal timing of downstream migration of steelhead smolts may

vary in response to a variety of environmental and physiological factors, including changes in water temperature, changes in streamflow, and increased turbidity resulting from stormwater runoff. The juvenile steelhead rear within the coastal marine waters for approximately 2 to 3 years before returning to their natal stream as spawning 4- or 5-year-old adults.

Because steelhead have a mandatory freshwater residency period, it is critical that suitable conditions for juvenile rearing exist year-round. Requirements for optimal juvenile rearing include adequate cover (i.e., greater than 25 percent of stream area), food supply (i.e., enough to sustain growth), and water temperatures of 43°F to 65°F (6°C to 18°C) (Raleigh et al. 1984). Although juveniles are known to withstand temperatures of up to 77°F (25°C), survival at these higher temperatures depends on a number of factors, including exposure duration, acclimation factors, food availability, water quality (specifically DO concentrations), and groundwater dynamics.


The steelhead life cycle is characterized by a high degree of flexibility (plasticity) in the duration of both their freshwater and marine rearing phases. The steelhead life cycle is adapted to respond to environmental variability in stream hydrology and other environmental conditions.

Factors affecting steelhead abundance are similar to those described for winter-run and spring-run Chinook salmon. One of the primary factors affecting population abundance of steelhead has been the loss of access to historical spawning and juvenile rearing habitat within the upper reaches of the Sacramento River and its tributaries and within the San Joaquin River as a result of the migration barriers caused by construction of major dams and reservoirs. Water temperatures within the rivers and creeks, particularly during summer and early fall, have also been identified as a factor affecting growth and survival of juvenile steelhead. Juvenile steelhead are vulnerable to entrainment at a large number of unscreened water diversions located along the Sacramento River and within the Delta, in addition to entrainment and salvage mortality at CVP and SWP export facilities. Changes in habitat quality and availability for spawning and juvenile rearing, exposure to contaminants, predation mortality, passage barriers and impediments to migration, changes in land use practices, and competition and interactions with hatchery-produced steelhead have all been identified as factors affecting steelhead abundance. Unlike Chinook salmon, steelhead are not vulnerable to recreational and commercial fishing within the ocean, although steelhead support a small inland recreational fishery for hatchery-produced fish. Ocean survival is affected by climatic and oceanographic conditions, and adults are vulnerable to predation mortality by marine mammals.

In recent years a number of changes have been made to improve the survival and habitat conditions for steelhead. Several large previously unscreened water diversions have been equipped with positive barrier fish screens. Improvements

to fish passage facilities have also been made to improve migration and access to spawning and juvenile rearing habitat.

Chinook Salmon The Sacramento River supports four separate runs of Chinook salmon: fall-run, late fall-run, winter-run, and spring-run, denoting when adults enter freshwater and begin their upstream migration. Figure 1-2 shows the seasonal occurrence of Chinook salmon in the Delta and tributary waters.



LEGEND
 DENOTES PRESENCE AND RELATIVE MAGNITUDE
 DENOTES ONLY PRESENCE

Source: Vogel and Marine 1991

Figure 1-2. Seasonal Occurrence of Different Life Stages of Four Chinook Salmon Runs

Fall-Run Chinook Salmon Fall-run Chinook salmon represent about 80 percent of the total Chinook salmon produced in the Sacramento River drainage (Kjelson et al. 1982). On March 9, 1998 (63 FR 11481), NMFS issued a proposed rule to list fall-run Chinook salmon as threatened, but determined the species did not warrant listing, and identified it as a candidate species (64 FR 50393, September 16, 1999). It was then changed to a species of concern in 2004. NMFS also determined that both late fall-run and fall-run comprise a single ESU, but because they are separate in timing and effects, they are distinguished as separately for the purposes of this document.

Although fall-run and late fall-run Chinook salmon inhabit a number of watersheds within the Central Valley for spawning and juvenile rearing, the largest populations occur within the mainstem Sacramento River, Feather River, Yuba River, American River, Mokelumne River, Stanislaus River, Tuolumne

River, and Merced River. Through the San Joaquin River Restoration Program, fall-run Chinook salmon captured at the Hills Ferry Barrier just upstream from the Merced River confluence, have been transported upstream to help reestablish Chinook salmon populations in the San Joaquin River downstream from Friant Dam. Fall-run Chinook salmon, in addition to spawning in these river systems, are also produced in fish hatcheries located on the Sacramento River, Feather River, American River, Mokelumne River, and Merced River.

Since 1999, hatchery spawners average more than 86,000 adults, and natural spawners average about 281,000 adults for the Sacramento and San Joaquin river system (CDFW 2014). Hatchery operations are intended to mitigate for the loss of access to upstream spawning and juvenile rearing habitat resulting from construction of dams and reservoirs within the Central Valley, in addition to producing fall-run Chinook salmon as part of the ocean salmon enhancement program to support commercial and recreational ocean salmon fishery. Fall-run Chinook salmon also support an inland recreational fishery.

Adult fall-run Chinook salmon migrate into the Sacramento River and its tributaries from July through December. Fall-run Chinook salmon spawn during early October through late December and incubation takes place during October through March. The peak of spawning is in October and November as water temperature drops. Fall-run Chinook salmon move upstream from the ocean in the late summer and early fall in mature condition and spawn soon after arriving at their spawning grounds. Juvenile Chinook salmon emerge from the gravel and migrate downstream to the ocean soon after emerging, rearing in the streams for only few months.

Temperature requirements vary according to life stage of Chinook salmon and habitat conditions. The following describes some general requirements for Chinook salmon, including all four runs occupying the Sacramento River. Most adult Chinook salmon migrate upstream when water temperatures are between 51 and 60°F (10.5 to 15.5°C) (Bell 1990, Hinze et al. 1956, as cited in McCullough et al. 2001). Spring-run Chinook salmon hold in waters typically under 60°F (15.5°C) (NMFS 1997), but because they hold in deep cold pools, surface water temperatures can reach as high as 73°F (22.8°C) (Beauchamp et al. 1983). Adults tend to spawn when water temperatures drop to between 41°F and 57°F (5°C to 13.9°C) (McCullough 1999, Moyle 2002, NMFS 2002, Slater 1963, Reiser and Bjornn 1979). During spawning, the female digs a nest (redd) with her tail before depositing her eggs while the male(s) alongside her fertilizes them.

The duration of egg incubation is temperature-dependent. Eggs will hatch sooner in warmer water, but water that is too warm during incubation can either kill the eggs directly or result in deformities and/or mortality post-hatching. The optimal range of water temperatures during egg incubation is between 41°F and 57°F (5°C to 13.9°C) (USFWS 1995, NMFS 1997, Slater 1963). Upon hatching, the young fish (alevins) will remain in the nest until their yolk sac has

been absorbed, at which time the young fish (now called fry) emerge from the redds. A portion of the fry population migrate downstream soon after emergence, where they rear within the lower river channels, Delta, and Suisun Bay during the spring months (Baker and Morhardt 2001). The remaining portion of juvenile Chinook salmon continue to rear in the upstream stream systems through the spring months, until they undergo smoltification, which typically takes place between April and early June. A small proportion of the fall-run Chinook salmon juveniles may, in some systems, rear through summer and fall, migrating downstream during fall, winter, or early spring as yearlings.

Water temperatures for rearing fry and juvenile Chinook salmon are optimal between 53°F and 60°F (11.7°C to 15.5°C) (NMFS 2000, 2002). Chinook salmon smolts begin to migrate downstream and through the Delta and the Bay to the ocean. Studies have shown that smoltification can be hindered and survival compromised when water temperatures exceed 62°F (16.7°C) (Zedonis and Newcomb 1997, Marine and Cech 2004).

The juvenile and adult Chinook salmon rear in coastal marine waters, foraging on fish and macroinvertebrates (e.g., northern anchovy, Pacific herring, squid, krill), until they reach maturation. Adult Chinook salmon spawn at ages ranging from approximately 2 to 5 years old, with the majority of adult fall-run Chinook salmon returning at 3 years old. Chinook salmon, unlike steelhead, die after spawning.

Late Fall-Run Chinook Salmon Late fall-run Chinook salmon mostly inhabit the Sacramento River, with spawning occurring upstream from RBPP. Late fall-run Chinook salmon migrate into the Sacramento River from October through April and spawn from January through April. Peak spawning activity in February and March is followed by egg incubation from January through June, and fry emergence from April through June. Rearing and emigration of fry and smolts occur from April through December. Juvenile Chinook salmon rear in the streams during summer, and in some streams they remain throughout the year.

Sacramento River Winter-Run Chinook Salmon With the possible exception of Battle Creek, the Sacramento River upstream from RBPP is the only spawning stream for winter-run Chinook salmon, which have been in a major decline since the 1960s. The abundance of winter-run Chinook salmon before the construction of Shasta Dam is unknown. Some biologists believe the run was relatively small, possibly consisting of a few thousand fish (Slater 1963). Others, relying on anecdotal accounts, believe the run could have numbered more than 200,000 fish (NMFS 1993a). The population during the mid-1960s, more than 20 years after the construction of Shasta Dam, exceeded 80,000 fish (Reclamation 1986). The population declined substantially during the 1970s and 1980s.

In 1989, Sacramento River winter-run Chinook salmon escapement was estimated at 695 adults. Escapement continued to decline, ranging between 185 and 1,240 fish between 1989 and 1994, with an average of 525 adults (CDFW 2014). The sharp decline in escapement during the late 1980s and early 1990s prompted listing of the winter-run Chinook salmon as endangered under ESA (59 FR 440, January 4, 1994) and CESA. Immediately following the listing, the numbers slowly began to increase. However, construction of the TCD in Shasta Dam in 1999 helped provide the necessary cold water for winter-run Chinook salmon. Since the completion of the TCD, the number of returning spawners has ranged between 827 in 2011 and 17,296 in 2006, averaging more than 6,000 adults (CDFW 2014).

The portion of the Sacramento River from Keswick Dam to Chipps Island, all waters westward from Chipps Island to the Carquinez Strait Bridge, all waters of San Pablo Bay, and all waters of the Bay north of the San Francisco-Oakland Bay Bridge have been designated as critical habitat for winter-run Chinook salmon (58 FR 33212, June 16, 1993). Critical habitat includes the river water, river bottom, and adjacent riparian zone (i.e., those adjacent terrestrial areas that directly affect a freshwater aquatic ecosystem).

As with other Chinook salmon stocks, NMFS is continuing to evaluate the status of the winter-run Chinook salmon population and the effectiveness of various management actions implemented within the Sacramento River, Delta, and ocean to provide improved protection and reduced mortality for winter-run salmon, in addition to providing enhanced habitat quality and availability for spawning and juvenile rearing. In July 2014, NMFS published the Recovery Plan for Sacramento River winter-run Chinook salmon, which identifies recovery goals, objectives, and criteria, as well as proposed management actions aimed at bringing the populations to a point at which they can be delisted.

Adult winter-run Chinook salmon spend 1 to 3 years in the ocean. Adult escapement consists of 67 percent 3-year-olds, 25 percent 2-year-olds, and 8 percent 4-year-olds (Hallock and Fisher 1985). Adult winter-run Chinook salmon leave the ocean and migrate through the Delta into the Sacramento River from November through July, passing RBPP on the Sacramento River from mid-December through July, with peak migration occurring during March (Moyle 2002). Most migrating adults have passed RBPP by late June (Moyle 2002). Winter-run Chinook salmon adults prefer water temperatures ranging between 57 and 67°F (14 to 19°C) for upstream migration (NMFS 2014).

Winter-run Chinook salmon spawn from mid-April through August (Moyle 2002). When water temperatures range between 50 and 59°F (10 to 15°C), egg incubation continues through mid-October. The primary spawning habitat in the Sacramento River is between the RBPP and Keswick Dam. Some fish may spawn below the RBPP, but warm-water temperature below the RBPP kills the eggs during most summers (Yoshiyama et al. 1998).

Juvenile winter-run Chinook salmon rear in the Sacramento River from July through March (Hallock and Fisher 1985). Most winter-run Chinook salmon fry pass the RBPP by October; most emigrating pre-smolts and smolts pass the RBPP by March (Martin et al. 2001).

Juvenile Chinook salmon move downstream from spawning areas in response to many factors, including inherited behavior, habitat availability, flow, competition for space and food, and water temperature. The number of juveniles that move, and the timing of movement, are highly variable. Storm events and the resulting high flow and turbidity appear to trigger downstream movement of substantial numbers of juvenile Chinook salmon. Juveniles have been observed in the Delta from November through May (NMFS 2014). In general, juvenile abundance in the Delta increases in response to increased Sacramento River flow (USFWS 1995).

Winter-run Chinook salmon smolts (i.e., juveniles that are physiologically ready to enter seawater) may migrate through the Delta and the Bay to the ocean from November through May (NMFS 2014).

A variety of environmental and biological factors have been identified that affect the abundance, mortality, and population dynamics of winter-run Chinook salmon. One of the primary factors that have affected population abundance of winter-run Chinook salmon has been the loss of access to historical spawning and juvenile rearing habitat within the upper reaches of the Sacramento River and its tributaries as a result of the migration barriers caused by Shasta and Keswick dams (Brandes and McLain 2001; Baker and Morhardt 2001). Operation of the Red Bluff Diversion Dam previously impeded adult upstream migration and vulnerability of juvenile winter-run Chinook salmon to predation mortality. However, in 2010, construction of the Red Bluff Fish Passage Improvement Project at the Red Bluff Diversion Dam began, and was completed in 2011, thus improving fish passage.

Water temperatures in the mainstem Sacramento River have been identified as a factor affecting incubating eggs, holding adults, and growth and survival of juvenile winter-run Chinook salmon rearing in the upper Sacramento River (Baker and Morhardt 2001). Juvenile winter-run Chinook salmon are also vulnerable to entrainment at a large number of unscreened water diversions located along the Sacramento River and within the Delta in addition to entrainment and salvage mortality at the CVP and SWP export facilities (Reclamation 2008). Changes in habitat quality and availability for spawning and juvenile rearing, exposure to contaminants and acid mine drainage, predation mortality by Sacramento pikeminnow, striped bass, largemouth bass, and other predators, and competition and interactions with hatchery-produced Chinook salmon have all been identified as factors affecting winter-run Chinook salmon abundance. In addition, subadult and adult winter-run Chinook salmon are vulnerable to recreational and commercial fishing; ocean survival is affected

by climatic and oceanographic conditions; and adults are vulnerable to predation mortality by marine mammals (Brandes and McLain 2001).

A number of changes have been made to improve survival and habitat conditions for winter-run Chinook salmon. The NMFS biological opinion (BO) for winter-run Chinook salmon (NMFS 1993) established water temperature objectives for the upper Sacramento River. Subsequent NMFS BOs in 2004 and 2009 reinforced these objectives, with the 2009 NMFS BO requiring water temperatures in the Sacramento River below 56°F at compliance locations between Balls Ferry and Bend Bridge from April 15 through September 30 to protect winter-run Chinook salmon (RPA Actions I.2.3 and I.2.4).

Recent changes in reservoir operations, including greater carryover storage, increased imports of cold water from the Trinity River system, and, most importantly, installation of a TCD on Shasta Dam, have substantially improved water temperature conditions in the reach. Modifications to CVP and SWP export operations have also been made in recent years to improve survival of juvenile salmon during migration through the Delta.

Spring-Run Chinook Salmon On September 16, 1999, the Central Valley spring-run Chinook salmon ESU was listed as threatened under the ESA by NMFS. The Central Valley spring-run Chinook salmon ESU includes all naturally spawned populations of spring-run Chinook salmon in the Sacramento River and its tributaries, as well as artificially propagated Feather River spring-run Chinook salmon (70 FR 37177, June 28, 2005).

Critical habitat for Central Valley spring-run was designated on February 16, 2000, but on April 30, 2002, the U.S. District Court for the District of Columbia approved a NMFS consent decree (National Association of Home Builders v. Evans) withdrawing the critical habitat designation for this and 18 other ESUs of salmon and steelhead. The consent decree challenged the process by which NMFS established the critical habitat designations, citing that the agency did not take into consideration the economic impacts on the interested parties, as required.

On December 10, 2004, NMFS published a new proposal to designate critical habitat for seven ESUs of Chinook salmon and steelhead in California, including the Central Valley spring-run Chinook salmon. The final designation for critical habitat was published on September 2, 2005, but was in effect on January 2, 2006. The critical habitat includes roughly 1,272 miles of occupied stream habitat and 427 square miles of estuarine habitat, and encompasses the lower Feather River; the Sacramento and Yuba rivers; Beegum, Battle, Clear, Cottonwood, Antelope, Mill, Deer, Butte, and Big Chico creeks; the north Delta (the central and south Delta were excluded); and Suisun, San Pablo, and north San Francisco bays (70 FR 52488, September 2, 2005).

In July 2014, NMFS published the Recovery Plan for Central Valley spring-run Chinook salmon, which identifies recovery goals, objectives and criteria, as well as proposed management actions aimed to bring the populations to a point at which they can be delisted. In the 2009 NMFS BO, RPA Action I.2.4 put protective measures in place for spring-run Chinook salmon with respect to water temperature, that, when possible, water temperatures should not exceed 56°F at the same compliance locations between Balls Ferry and Bend Bridge from October 1 through October.

Historical records indicate that adult spring-run Chinook salmon enter the mainstem Sacramento River in February and March. Adults hold in deep, cold pools near spawning habitat until spawning commences in late summer and fall. Spring-run Chinook salmon are sexually immature during upstream migration (Fisher 1994). Spawning occurs in gravel substrates in late August through October. Considerable overlap occurs between spring-run and fall-run Chinook spawning on the mainstem Sacramento River and most of the major tributaries. This overlap has likely resulted in genetic introgression (i.e., loss of genetic purity) of the spring-run stocks (Slater 1963). Genetically pure spring-run Chinook salmon occur mostly only in two spawning tributaries, Mill and Deer creeks.

Juveniles emerge during November and December in most locations but may emerge later when water temperature is cooler. Spring-run Chinook salmon may migrate downstream as young-of-year juveniles or yearlings. Based on observations in Butte Creek and the Sacramento River, young-of-year juveniles migrate during November to June. Yearling spring-run Chinook salmon migrate during October through March, with peak migration in November (Cramer and Demko 1997, Hill and Webber 1999). The downstream migration of both spring-run Chinook salmon fry and yearlings during the late fall and winter typically coincides with increased flow and turbidity associated with winter stormwater runoff.

Juvenile spring-run Chinook salmon rear in their natal streams, the mainstem Sacramento River, and the Delta. Juveniles that remain in their natal streams, especially small, cold tributary streams, may migrate downstream as yearlings. Juveniles migrate downstream to the ocean as yearlings with the onset of the storm season in October of the year following spawning, and migration may continue through March (CDFG 1998).

A variety of environmental and biological factors have been identified that affect the abundance, mortality, and population dynamics of spring-run Chinook salmon. The main factor affecting population abundance of spring-run Chinook salmon is the loss of access to historical spawning and juvenile rearing habitat within the upper reaches of the Sacramento River and its tributaries and San Joaquin River as a result of the migration barriers caused by construction of major dams and reservoirs. Water temperatures have been identified as

affecting incubating eggs, holding adults, and growth and survival of juvenile spring-run Chinook salmon.

Juvenile spring-run Chinook salmon are also vulnerable to entrainment at a large number of unscreened water diversions located along the Sacramento River and within the Delta, in addition to entrainment and salvage mortality at the CVP and SWP export facilities. Changes in habitat quality and availability for spawning and juvenile rearing, exposure to contaminants, predation mortality by Sacramento pikeminnow, striped bass, largemouth bass, and other predators, and competition and interactions with hatchery-produced Chinook salmon have all been shown to affect spring-run Chinook salmon abundance. In addition, as for winter-run Chinook salmon, subadult and adult spring-run Chinook salmon are vulnerable to recreational and commercial fishing; ocean survival is affected by climatic and oceanographic conditions; and adults are vulnerable to predation mortality by marine mammals.

A number of changes have been made to improve the survival and habitat conditions for spring-run Chinook salmon. Several large previously unscreened water diversions have been equipped with positive barrier fish screens. Changes to ocean salmon fishing regulations have been made to improve the survival of adult spring-run Chinook salmon. Modifications to CVP and SWP export operations have been made to improve survival of juvenile Chinook salmon migrating through the Delta. Improvements in fish passage facilities have also been made to improve migration and access to Butte Creek. These changes and management actions are thought to have contributed to the trend of increasing abundance of adult spring-run Chinook salmon returning to spawn in Butte Creek and other habitats within the upper Sacramento River system.

Coho Salmon General life history information and biological requirements of SONCC Coho salmon have been described in various documents (Shapovalov and Taft 1954, Hassler 1987, Sandercock 1991, CDFG 1994, Weitkamp et al. 1995), as well as the NMFS final rule listing SONCC Coho salmon (May 6, 1997; 62 FR 24588).

Adult Coho salmon typically enter rivers between September and February. Spawning occurs from November to January (Hassler 1987), but occasionally as late as February or March (Weitkamp et al. 1995). Coho salmon eggs incubate for 35-50 days between November and March. Successful incubation depends on several factors: DO levels, temperature, substrate size, amount of fine sediment, and water velocity. Fry start emerging from the gravel 2 to 3 weeks after hatching and move into shallow areas with vegetative or other cover. Peak emergence periods in the Trinity River are February through March (USFWS and Hoopa Valley Tribe 1999). As fry grow larger, they disperse upstream or downstream. In summer, Coho salmon fry prefer pools or other slower velocity areas such as alcoves, with woody debris or overhanging vegetation. Juvenile Coho salmon over-winter in slow-water habitat with cover. Juveniles may rear in freshwater for up to 15 months, then migrate to the ocean as smolts from

March to June (Weitkamp et al. 1995). Coho salmon adults typically spend 2 years in the ocean before returning to their natal streams to spawn as 3-year-olds.

Green Sturgeon North American green sturgeon have been separated into two DPSs: the northern DPS (all populations north of, and including, the Eel River) and the southern DPS (Coastal and Central Valley populations south of the Eel River). The southern DPS is currently listed as threatened under the ESA. On April 15, 2004, NMFS announced that the northern and southern DPSs of green sturgeon would change in listing status from a candidate species to a species of concern (69 FR 117, June 18, 2004). However, litigation challenged the NMFS determination that green sturgeon did not warrant listing as an endangered or threatened species under the ESA and asserted that the agency was arbitrary and capricious in failing to examine whether habitat loss constituted a significant portion of the species' range (70 FR 65, April 6, 2005). The court partially agreed with the plaintiff's motion, and remanded the determination back to NMFS for further analysis and decision as to whether green sturgeon are endangered or threatened in a significant portion of their range. Following this, NMFS listed green sturgeon as threatened (71 FR 17757, April 7, 2006). In April 2009, NMFS designated critical habitat for green sturgeon that includes the Sacramento, lower Feather, and lower Yuba rivers, Yolo and Sutter bypasses, the Delta, and Suisun, San Pablo, and San Francisco bays (74 FR 52300, April 9, 2009).

Not much is known about the life history of green sturgeon because of its low abundance, low sport fishing value, and limited spawning distribution, but spawning and larval ecology are assumed to be similar to that of white sturgeon (Moyle 2002, Beamesderfer and Webb 2002). Green sturgeon are mostly marine fish, spending limited time in estuaries or freshwater (State Water Board 1999). Green sturgeon also make extensive ocean migrations; consequently, most recoveries of individuals tagged in San Pablo Bay have come from the ocean and from rivers and estuaries in Oregon and Washington.

Within estuaries, green sturgeon reportedly tend to concentrate in deep areas with soft bottoms. In rivers, adult (and juvenile) green sturgeon have been observed primarily on clean sand (Environmental Protection Information Center et al. 2001). Adult green sturgeon are benthic, usually found in the Sacramento River in deep, off-channel areas with little current.

Indirect evidence indicates that green sturgeon spawn mainly in the Sacramento River; spawning has been reported in the mainstem as far north as Red Bluff. Migration of green sturgeon begins in late February and continues through July (for both upstream and downstream migration) and may cover as much as 200 miles (Beamesderfer and Webb 2002). Adults and juveniles are opportunistic carnivores, feeding on benthic invertebrates and may also take small fish (Adams et al. 2002). Adult green sturgeon are also known to feed on worms, clams, sand lances, callianassid shrimp, crabs, isopods, and anchovies

(Environmental Protection Information Center et al. 2001, Moyle 2002). Green sturgeon can withstand long periods of food deprivation during spawning migrations (Environmental Protection Information Center et al. 2001).

Most females reach sexual maturity at 20 to 25 years and 6 to 7 feet in length while males reach sexual maturity at 15 to 17 years and 5 to 6 feet in length (Beamesderfer and Webb 2002). Green sturgeon are thought to spawn every 3 to 5 years (70 FR 65, April 6, 2005). The green sturgeon spawning period is from February to July, with a peak in mid-April to mid-June (Kohlhorst 1976, Moyle 2002, Beamesderfer and Webb 2002). The reported range of preferred/optimal water temperatures for green sturgeon spawning is unclear, but spawning success is related to water temperature (Beamesderfer and Webb 2002). In the Sacramento River, sturgeon are seen in the river when water temperatures are between 46°F and 57°F (13.9°C) (Moyle 2002). Spawning occurs in deep pools in large, turbulent river mainstems (Moyle et al. 1992), and the preferred spawning substrate is likely large cobble-containing crevices in which eggs can become trapped and develop, but may range from clean sand to bedrock (Environmental Protection Information Center et al. 2001, Beamesderfer and Webb 2002).

Sturgeon eggs have been found in the Sacramento River from mid-February through July (Kohlhorst 1976, Moyle 2002, Beamesderfer and Webb 2002). Eggs are broadcast-spawned and externally fertilized in relatively high water velocities (1.5 to 3.0 meters per second) and probably at depths greater than 10 feet (USFWS 1996). The number of eggs green sturgeon females lay in a spawning season increases with body size, reportedly ranging from 60,000 to 140,000 eggs per female (Moyle et al. 1992) and are the largest egg of any sturgeon (Cech et al. 2000). Green sturgeon eggs are slightly adhesive, adhering to each other and to river substrates (CDFG 2002). The importance of water quality is uncertain, but silt is known to prevent green sturgeon eggs from adhering to each other (USFWS 1996)—sand and silt may suffocate the eggs (Environmental Protection Information Center et al. 2001). The comparatively large egg size, thin chorionic layer on the egg, and other characteristics suggest that green sturgeon probably requires colder, cleaner water for spawning than does the white sturgeon (USFWS 1996). Water temperatures above 68°F (20°C) are reportedly lethal to green sturgeon embryos (Beamesderfer and Webb 2002). Eggs hatch approximately 196 hours after spawning, and larvae are 8 to 19 millimeters long. Juveniles range in size from less than 1 inch to almost 5 feet.

Juvenile green sturgeon reportedly occur in shallow water (Radtke 1966) and probably move to deeper, more saline areas as they grow (Environmental Protection Information Center et al. 2001). Rearing juveniles remain in freshwater for 1 to 4 years before returning to their marine environment (Beamesderfer and Webb 2002, Environmental Protection Information Center et al. 2001). Juveniles in the Delta primarily feed on opossum shrimp and amphipods (Radtke 1966, Moyle 2002). The growth rate for green sturgeon

juveniles is roughly 3 inches per year until they reach maturity at 4 to 5 feet in length, around age 15 to 20, at which time the growth rate slows (Wang 1986).

The occurrence of green sturgeon in fishery sampling, and CVP/SWP (Jones Pumping Plant and Banks Pumping Plant) fish salvage is extremely low and therefore has not been used to represent the seasonal period of juvenile movement through the Delta. During 2007, for example, green sturgeon were collected in the Jones and Banks fish facilities during one day at each out of the year. Green sturgeon tend to remain near estuaries at first but may migrate considerable distances as they grow larger (State Water Board 1999).

There is no direct evidence of a decline in the numbers of green sturgeon in the Sacramento River. However, the population is so small that a collapse could occur, and it would hardly be noticed because of limited occurrence in conventional fishery sampling programs (State Water Board 1999). In the Delta, major factors that may negatively affect green sturgeon abundance are sport fisheries, modification of spawning habitat, entrainment, and toxic substances.

Delta Smelt Delta smelt is Federally listed as threatened (58 FR 12854, March 5, 1993); critical habitat was designated on December 19, 1994. Critical habitat includes the portion of the Sacramento River from Keswick Dam to Chipps Island, all waters westward from Chipps Island to the Carquinez Bridge, all waters of San Pablo Bay, and all waters of the Bay north of the San Francisco-Oakland Bay Bridge. Recent monitoring results indicate that the threatened delta smelt population continues to remain at or near all-time lows. In 2006, the USFWS was petitioned to upgrade the status of delta smelt to endangered (Center for Biological Diversity et al. 2006). In 2010, the USFWS conducted their 5-year review and found delta smelt warranted the upgrade in status, however, the listing was precluded by other higher priority-listing actions (75 FR 17667). The status of delta smelt under CESA was upgraded to endangered in January 2010 (CDFG 2011).

Delta smelt are endemic to the Delta. During the spawning season, adults move into the channels and sloughs of the Delta. When Delta outflows are high, delta smelt may occur in San Pablo Bay. Delta smelt have relatively low fecundity and most live for 1 year (Moyle 2002).

Estuarine rearing habitat for juvenile and adult delta smelt is typically found in the waters of the lower Delta and Suisun Bay where salinity is between 2 and 7 ppt. Delta smelt tolerate 0 to 19 ppt salinity. They typically occupy open shallow waters (less than 10 feet) but also occur in the main channel in the region where freshwater and brackish water mix. The zone may be hydraulically conducive to their ability to maintain position and metabolic efficiency.

Adult delta smelt begin a spawning migration, which may encompass several months, toward areas of the upper Delta and toward freshwater during

December or January. Spawning occurs between February and July, with peak spawning during April through mid-May. Spawning occurs in shallow edge-waters in the upper Delta channels, including the Sacramento River above Rio Vista, Cache Slough, Lindsey Slough, and Barker Slough. Spawning has not been documented in the Sacramento River upstream from the DCC. Eggs are broadcast over the bottom, where they attach to firm sediment, woody material, and vegetation. Hatching takes approximately 9 to 13 days and larvae begin feeding 4 to 5 days later. Newly hatched larvae contain a large oil globule that makes them semibuoyant and allows them to stay off the bottom. Larval smelt feed on rotifers and other zooplankton. As their fins and swim bladder develop, they move higher into the water column. Larvae and juveniles gradually move downstream toward rearing habitat in the estuarine mixing zone.

Longfin Smelt In April 2010, CDFW designated the longfin smelt as a threatened species (CDFG 2011). The USFWS were petitioned to list longfin smelt as endangered in 2007 (Center for Biological Diversity et al. 2007). The USFWS found that the Bay-Delta DPS does warrant listing, however, as with the delta smelt, the listing is precluded by other higher priority actions. Therefore, longfin smelt have been added to the candidate list (77 FR 19756, April 2, 2012). Historically, longfin smelt populations were found in the Klamath, Eel, and San Francisco estuaries, and in Humboldt Bay. From current sampling, populations reside at the mouth of the Klamath River and the Russian River estuary. In the Central Valley, longfin are rarely found upstream from Rio Vista or Medford Island in the Delta. Adults concentrate in Suisun, San Pablo, and North San Francisco bays (Moyle 2002).

Longfin smelt are anadromous, euryhaline, and nektonic. Adults and juveniles are found in estuaries and can tolerate salinities from 0 ppt to pure seawater. After the early juvenile stage, they prefer salinities in the 15 through 30 ppt range (Moyle 2002).

Longfin smelt are found in San Pablo Bay in April through June and disperse in late summer. In fall and winter, yearlings move upstream into freshwater to spawn. Spawning occurs below Medford Island in the San Joaquin River and below Rio Vista on the Sacramento River, as early as November, and larval surveys indicate spawning may extend into June (Moyle 2002).

While the eggs are adhesive, embryos, which hatch in 40 days at 45°F (7.2°C), are buoyant. They move into the upper part of the water column and are carried into the estuary. High outflows transport the larvae into Suisun and San Pablo Bays. In low-outflow years, larvae move into the western Delta and Suisun Bay. Higher outflows reflect positively in juvenile survival and adult abundance. Rearing habitat is better in Suisun and San Pablo bays because juveniles require brackish water in the 2 to 18 ppt range. If juveniles stay in the Delta, they become entrained and exposed to more adverse conditions (Moyle 2002). Seasonal occurrence of longfin smelt in CVP and SWP salvage is considered to

be representative of the seasonal periods when juvenile and adult longfin smelt would be in the Delta.

Consistently, a measurable portion of the longfin smelt population survives into a second year. During the second year of life, they inhabit the Bay and, occasionally, the Gulf of the Farallones (Wang 1986). This explains their common identification as anadromous (State Water Board 1999). Because longfin smelt seldom occur in freshwater except to spawn, but are widely dispersed in brackish waters of the Bay, it is likely that their range formerly extended as far up into the Delta as saltwater intruded. The easternmost catch of longfin smelt in fall mid-water trawl samples has been at Medford Island in the Central Delta. The depth of habitat is a pronounced difference between the two species in their region of overlap in Suisun Bay; longfin smelt are caught in greater quantities at deep stations (more than 32 feet), whereas delta smelt are more abundant at shallow stations (less than 10 feet) (State Water Board 1999).

The main food of longfin smelt is the opossum shrimp (*Neomysis mercedis*), although copepods and other crustaceans are important at times, especially to small fish. Longfin smelt, in turn, are eaten by a variety of predatory fishes, birds, and marine mammals (State Water Board 1999). Recent declines in the abundance of opossum shrimp and other zooplankton have been identified as a factor affecting the abundance of longfin smelt.

Longfin smelt were once one of the most common fish in the Delta. Their abundance has fluctuated widely in the past but, since 1982, abundance has declined significantly (Baxter 1996, The Bay Institute et al. 2007). The abundance of longfin smelt also has declined relative to other fishes, dropping from first or second in abundance in most trawl surveys during the 1960s and 1970s, to seventh or eighth in abundance. Abundance improved substantially in 1995 but was again relatively low in 1996 and 1997. Longfin abundance indices, although variable, were at very low levels. The causes of decline are thought to be multiple and synergistic, including reduction in outflows, entrainment losses to water diversions, climatic variation, toxic substances, predation, and introduced species (State Water Board 1999).

Sacramento Splittail In 1999, after 4 years of candidate status, the splittail was listed as threatened under the ESA (64 FR 25, March 10, 1999). Fall midwater trawl surveys indicate that juvenile splittail abundance has been highly variable from year to year, with peaks and declines coinciding with wet and dry periods, respectively, and correlated with the availability of flooded shallow water habitat. After the listing, the State Water Contractors, San Luis and Delta-Mendota Water Authority, and others challenged the listing, contending that it violated the ESA and the Administrative Procedures Act. On June 23, 2000, the U.S. District Court in Fresno ruled in favor of the plaintiffs and found the listing unlawful. On September 22, 2003, USFWS delisted splittail as a threatened species because habitat restoration actions under CALFED and the CVPIA were thought to likely keep the splittail from

becoming endangered in the foreseeable future (68 FR 55139, September 22, 2003). Splittail is identified as a species of special concern under CESA.

Splittail are found primarily in the Delta, Suisun Bay, Suisun Marsh, and Napa Marsh, but juveniles have been found in the Sacramento River as far upstream as its tributaries and Red Bluff (Sommer et al. 1997). Sommer et al. (1997, 2002) found that the Yolo and Sutter Bypasses provide important spawning habitat for splittail. Some adults spend the summer in the mainstem Sacramento River rather than return to the estuary.

The Sacramento splittail, which has a high reproductive capacity, can live 5 to 7 years, and generally begins spawning at 2 years of age. Spawning, which seems to be triggered by increasing water temperatures and day length, occurs over beds of submerged vegetation in slow-moving stretches of water (such as flooded terrestrial areas and dead-end sloughs). Adults spawn from February through May in the Delta, upstream tributaries, Napa Marsh, Napa and Petaluma rivers, Suisun Bay and Marsh, and the Sutter and Yolo bypasses (Baxter et al. 1996). Splittail prefer low water velocities for spawning and early rearing. However, some current is required to keep water temperature and clarity low, keep eggs free of silt, and facilitate suspension and attachment of eggs on vegetation (Jones & Stokes 2001). Adult splittail deposit adhesive eggs over flooded terrestrial or aquatic vegetation when water temperature is between 48 and 68°F (9°C to 20°C) (Moyle 2002, Wang 1986). Spawning occurs in depths less than 6 feet (Moyle et al. 2004). Splittail spawn in late April and May in Suisun Marsh and between early March and May in the upper Delta and lower reaches and flood bypasses of the Sacramento and San Joaquin Rivers (Moyle et al. 1995). Spawning has been observed to occur as early as January and may continue through early July (Wang 1986, Moyle 2002).

Larval splittail are commonly found in shallow, vegetated areas near spawning habitat. Larvae eventually move into deeper and more open water habitat as they grow and become juveniles. During late winter and spring, young-of-year juvenile splittail (i.e., production from spawning in the current year) are found in sloughs, rivers, and Delta channels near spawning habitat. Juvenile splittail gradually move from shallow, nearshore areas to the deeper, open water habitat of Suisun and San Pablo bays (Wang 1986). Young splittail may occur in shallow and open waters of the Delta and San Pablo Bay, but they are particularly abundant in the northern and western Delta (Sommer et al. 1997; State Water Board 1999). The seasonal occurrence of juvenile splittail in CVP and SWP fish salvage is representative of the periods when juvenile splittail inhabit the Delta. In areas upstream from the Delta, juvenile splittail can be expected to be present in the flood bypasses when these areas are inundated during winter and spring (Jones & Stokes Associates 1993, Sommer et al. 1997).

Although the Sacramento splittail is generally considered a freshwater species, the adults and subadults have an unusually high tolerance for saline waters (up

to 10 to 18 ppt) for a member of the minnow family (Young and Cech 1996). The salt tolerance of splittail larvae is unknown, but they have been observed in water with salinities of 10 to 18 ppt (State Water Board 1999).

Splittail are bottom foragers that feed extensively on opossum shrimp and opportunistically on earthworms, clams, insect larvae, and other invertebrates. They are preyed on by striped bass and other predatory fish in the estuary. In the past, anglers commonly used splittail as bait when fishing for striped bass (State Water Board 1999).

Hardhead Hardhead are widely distributed throughout the low- to mid-elevation streams in the main Sacramento-San Joaquin drainage as well as in the Russian River drainage. Undisturbed portions of larger streams at low to middle elevations are preferred by hardhead. They are able to withstand summer water temperatures above 68°F (20°C); however, hardhead will select lower temperatures when they are available. They are fairly intolerant of low-oxygenated waters, particularly at higher water temperatures. Pools with sand-gravel substrates and slow water velocities are the preferred habitat; adult fish inhabit the lower half of the water column, while the juvenile fish remain in shallow water closer to the stream edges. Hardhead tend not to do well in areas where introduced centrarchid fish (sunfish and bass) are abundant. Hardhead are relatively common in the Sacramento River from below Keswick Dam to the Tehama-Butte county line, where the river is less channelized, and in the low to mid-elevation reaches of most of its perennial tributaries (Moyle 2002). Although abundant in the Pit River above Shasta Lake, especially in Pacific Gas and Electric Company's run-of-the-river hydroelectric reservoirs (Moyle 2002; Pacific Gas and Electric Company 2007), hardhead have not been found in the Sacramento and McCloud rivers above the lake in recent surveys (Nevares and Liebig 2007; Weaver and Mehalik 2008). Hardhead typically feed on small invertebrates and aquatic plants at the bottom of quiet water (Moyle 2002). Hardhead is a State species of special concern and a USFS designated sensitive species.

Striped Bass Striped bass are anadromous fish that have been an important part of the sport fishing industry in the Delta. They were introduced into the Sacramento-San Joaquin estuary between 1879 and 1882 (Moyle 2002). Striped bass will not use fish ladders; therefore, their range in the Sacramento River is limited to the reach of the river below the RBPP. Striped bass may move into the lower reaches of the rivers year-round but probably most often between April and June, when they spawn. The species tends to remain in deep, slow-moving water, where it has access to prey without having to expend a great deal of energy.

Other Important Native Fish Species Present in Study Area

Upper Klamath-Trinity Chinook Salmon Upper Klamath-Trinity (UKT) Chinook salmon are found in the Trinity River within the extended study area (see biological requirements described above for Chinook salmon).

Klamath Mountains Province Steelhead KMP steelhead are found in the Trinity River within the extended study area and have similar biological requirements to those for steelhead described above.

California Roach California roach are distributed throughout the State; however, a specific subspecies is found in the Sacramento River drainage (excluding the Pit River), including tributaries to the Bay. California roach occupy small, warm streams with intermittent flow in mid-elevation foothills. Dense populations often occur in isolated pools. They are tolerant of high temperatures (86°F to 95°F (30°C to 35°C)) and low oxygen levels, although they also can be found in cold, well-oxygenated systems; human-modified habitats; and the main channels of larger rivers.

The California roach composes multiple subspecies, all of which are included as Federal Species of Concern, and all but one subspecies of which is identified by California as a Species of special concern.

White Sturgeon The white sturgeon, the largest freshwater or anadromous fish species in North America, can reach record sizes over 1,300 pounds. Historically, white sturgeon populations ranged from Alaska to central California (Moyle 2002); however, major spawning populations are now limited to the Fraser River (British Columbia, Canada), the Columbia River (Washington), and the Sacramento-San Joaquin River system.

Habitat use varies among populations. Portions of populations are considered anadromous, using fresh, brackish, and marine waters during different phases of their life history. White sturgeon are long-lived fish and can live as long as 100 years; however, fish that old are seldom found.

Upstream spawning migrations of white sturgeon in the Sacramento-San Joaquin river system occur between February and June (Miller 1972, Kohlhorst 1976, Wang 2006). Only a portion of the total adult sturgeon population migrates upstream from the Delta each year. Sturgeon that do move upstream are believed to be mature and ready to spawn.

Based on the recoveries of tagged adult sturgeon between 1974 and 1988, and collection of sturgeon eggs, larvae, and juveniles, most white sturgeon migrating up the Sacramento River congregate and spawn between Knights Landing and a point just above Colusa; however, juvenile sturgeon have been found by USFWS as far upstream as the RBPP.

The environmental cues that initiate upstream migration are not well understood. Mature fish could be stimulated to migrate upstream by cues triggering the final stages of gonadal development – such factors as flow, velocity, photoperiod (i.e., the number of daylight hours best suited to the growth and maturation of an organism), or temperature (Pacific States Marine Fisheries Commission 1992).

White sturgeon spawn in the Sacramento River between mid-February and late May, with a peak in spawning (93 percent) occurring between March and April (Kohlhorst 1976). Not all adults migrate upstream to spawn each year. Sexual cycles in sturgeon are complex because these fish mature at a late age and adults do not spawn every year. It is likely that only mature sturgeon migrate upriver to spawn and that most immature fish or fish in resting stages remain in the estuary. Few observations of wild sturgeon spawning have been reported. Apparently, sturgeon broadcast spawn in swift water. The current initially disperses the adhesive eggs, which sink and adhere to gravel and rock on the bottom. The adhesive properties of the eggs are adaptive to spawning and retention of eggs within swift current environments. Sediments can reduce this adhesiveness of eggs (Conte et al. 1988). Optimum temperatures for incubation and hatching are around 59°F (15°C); higher temperatures result in greater mortality and premature hatching (Adams et al. 2002).

Laboratory studies indicate that larval sturgeon demonstrate three behavioral phases after emergence: swim-up and dispersal, hiding, and feeding (Duke et al. 1990, Miller et al. 1991). After hatching, yolk sac larvae swim up into the water column. The currents act as a dispersal mechanism, transporting larvae downstream from the spawning area. Larvae swim toward or to the surface, then passively sink to the bottom (Brewer 1987). Either immediately or shortly after touching bottom, the larvae repeat the swimming activity.

When larvae enter the hiding phase, they are still nourished from the yolk sac. To hide, larvae place their heads within substrates (either rock or vegetation) and maintain a constant tail beat to retain their position. Substrate preference of hiding larvae is related to the degree of darkness the substrate provides, a negative phototactic (i.e., movement away from light) response. This hiding behavior may provide protection from predation as the larvae develop.

Larval sturgeon develop the mouth and olfactory morphology needed for feeding before the yolk sac is completely absorbed. Exogenous feeding occurs approximately 12 days after hatching at temperatures of 63°F (17.2°C) (Buddington and Doroshov 1984). During this phase, the larvae move out of hiding to forage actively for food. Young sturgeon appear to be opportunistic feeders (Moyle 2002). The senses of smell and touch appear to be more important than vision for locating prey. Larvae are territorial during this phase (Brannon et al. 1984).

The diet of sturgeon changes as the fish become larger. Young-of-year sturgeon (less than 8 inches long) feed on a number of prey, including small crustaceans and insect larvae, and can potentially consume small fish fry. As the fish grow, the diet becomes more diverse and includes several benthic invertebrates and seasonally abundant food items, such as fish eggs or fry. McKechnie and Fenner (1971) found that adult sturgeon caught in San Pablo and Suisun bays feed primarily on benthic invertebrates, including clams, barnacles, crab, and shrimp.

Seasonally, herring eggs and small fish, such as striped bass, flounder, goby, and herring, are important prey items.

Adult and subadult sturgeon inhabit estuarine areas year-round. Distribution in the Delta is thought to depend primarily on river flow and consequent salinity regimes. The center of the population is upriver during low-flow years and downriver during high-flow years.

Sacramento Sucker The Sacramento sucker is widely distributed throughout the Sacramento River system. Sacramento sucker occupy waters from cold, high-velocity streams to warm, nearly stagnant sloughs. They are common at moderate elevations (600 to 2,000 feet). Sacramento sucker feed on algae, detritus, and benthic invertebrates. They usually spawn for the first time in their fourth or fifth years. When they cannot move upstream, and instead spawn in lake habitat, they typically orient themselves near areas where spring freshets flow into the lake. They typically spawn in stream habitat on gravel riffles from late February to early June. The eggs hatch in 3 to 4 weeks, and the young typically live in the natal stream for a couple of years before moving downstream to a reservoir or large river (Moyle 2002).

Sacramento Pikeminnow Sacramento pikeminnow occupy rivers and streams throughout the Sacramento-San Joaquin river system, mainly at elevations between 300 and 2,000 feet. Sacramento pikeminnow spawn in April and May, with eggs hatching in less than a week. Within a week of hatching, the fry are free-swimming and schooling. Adult pikeminnow may feed on other fish, including juvenile pikeminnow, Chinook salmon, and steelhead, but, according to Moyle (2002), are overrated as predators on salmonid species in natural environments. They can, however, be major predators on juvenile salmon and steelhead in riverine environments modified by dams and fish ladders. Pikeminnow tend to remain in well-shaded, deep pools with sand or rock substrate and are less likely to be found in areas where there are higher numbers of introduced predator species, such as largemouth bass and other centrarchid species.

Pacific Lamprey Similar to Chinook salmon and steelhead, lamprey adults migrate upstream from the ocean during the winter and spring to spawn (Moyle 2002). Spawning occurs over gravel substrates. Larval lamprey rear in sand and mud substrates, gradually moving downstream over the rearing period. Little is known about water quality requirements and other habitat needs.

Important Nonnative Fish Species Present in Study Area

American Shad American shad are an anadromous fish that have been introduced into the Central Valley and have become established as a popular sport fish. American shad are present in the Sacramento River up to Red Bluff and in the lower reaches of the American and Feather rivers. American shad use the San Francisco Estuary after migrating from the ocean in the fall. They move into freshwater where they spawn from March to May. In the Sacramento River

basin, the main summer rearing areas are the lower Feather River, the Sacramento River from Colusa to the north Delta and, to some extent, the south Delta. Juvenile shad move to the ocean from September to November, although juvenile migration under high outflow conditions may begin in June.

Catfishes Four species in the catfish family are found in the study area – channel catfish, white catfish, black bullhead and brown bullhead. All were introduced into California. Channel catfish were established in the Sacramento-San Joaquin system in the 1940s. White catfish were brought into California in a small introduction to the San Joaquin River near Stockton in 1874. The earliest confirmed record of black bullhead in California was 1942. Brown bullhead were also among the earliest (1874) successful transplants to California.

Channel catfish are typically found in main channels of large rivers and streams, but inhabit a wide variety of water bodies, including farm ponds; reservoirs; turbid, muddy-bottom rivers; and large streams with ample riffle habitat. They can tolerate low oxygen levels (1 to 2 parts per million) and high water temperatures (97°F to 100°F (36°C to 37.8°C)). They tend to feed on detritus and plant material, but will ingest invertebrates and fish as well. These rapidly growing fish spawn anywhere from 2 to 8 years old, from April to June. They prefer cave-like sites for their spawning nests, such as undercut banks or log jams. Water temperatures between 70 and 84°F (21 and 28.9°C) are suitable for spawning. Eggs hatch in 6 to 10 days, and the young are actively swimming within 2 days of hatching (Moyle 2002, Wang 1986).

White catfish occupy slow-current habitat, avoiding areas with heavy beds of aquatic plants, or water less than 7 feet deep. They are often found in warm-water lakes, reservoirs, and farm ponds. Water temperatures must exceed 68°F in the summer and, if the lake they occupy stratifies, they will move to the level where the water temperatures exceed 70°F (21°C). White catfish are carnivorous bottom feeders, feeding primarily on smaller fish such as threadfin shad and silverside, and invertebrates and carrion. Spawning occurs from June to July, and eggs hatch about a week after spawning (Moyle 2002).

Black bullhead prefer ponds, small lake, river backwaters, and small stream pools with warm and turbid water, muddy bottoms, slow currents, and few other fish species. They are capable of surviving water temperatures up to 98°F (35°C) and salinities up to 13 ppt (Moyle 2002). Most foraging occurs at night, feeding mostly on aquatic insects, crustaceans, mollusks, and both live and dead fish. Spawning takes place after water temperatures exceed 68°F (20°C), during June and July, in a mud nest excavated by the female (Moyle 2002).

Brown bullhead are common throughout California, adapting to a large variety of water body types. They prefer water 7 to 16 feet deep with aquatic vegetation and sandy, muddy bottoms. They can survive a wide range of water temperatures (from 32°F to 99°F (0°C to 37.2°C)), although they prefer water

temperatures between 68°F and 95°F (20°C and 35°C). Brown bullhead feed on invertebrates, crustaceans, and fish, including silversides. Brown bullhead spawn for the first time during their third year, in May and June (Moyle 2002).

Sunfish Sunfish are a popular game fish in California, and almost every species has been introduced into California since the late 1800s. Typically, these fish prefer warm ponds and lakes, or slow moving streams, but can be found in the Sacramento River, including bluegill and green sunfish. A common trait among sunfish is the building of nests and the subsequent defending of the nest by the male of the species.

Bluegill are one of the most abundant sunfish in California. They prefer warm, shallow lakes, reservoirs, ponds, and sloughs at low altitudes. They can survive in waters with high turbidity and low oxygen levels. They are typically found around rooted aquatic vegetation, where they hide and feed. Substrate is typically silt, sand, or gravel, and they typically do not go deeper than 16 feet. Bluegill feed on whatever is most abundant, including aquatic insect larvae, planktonic crustaceans, terrestrial insects, snails, small fish, fish eggs, and even crayfish. Spawning occurs in the spring when water temperatures reach 64°F to 70°F (17.8°C to 21°C), and will continue through the summer. Eggs hatch within 2 to 3 days (Moyle 2002).

Green sunfish are aggressive, stout-bodied fish with large mouths that occupy small, warm intermittent streams, ponds, and lake edges. In lake conditions, they stay in shallow weedy areas, where there are few other species. Green sunfish are territorial and opportunistic predators, feeding on more active invertebrates and on small fish, including mosquitofish and other smaller sunfish. They begin spawning in their third year, and the spawning season is from May and June, but sometimes continues until August. Eggs hatch in 5 to 7 days, and the young are soon after free-swimming individuals (Moyle 2002.)

Black Bass Black bass, also in the sunfish family, is a generic name for several bass species, including largemouth and smallmouth bass. Both largemouth and smallmouth bass were introduced into California in 1874; they are some of the most valuable game fish in the state.

Largemouth bass are typically found in warm, quiet water with low turbidity, such as ponds, lakes, sloughs and river backwaters that contain beds of aquatic plants. Optimal growth occurs when water temperatures are between 68°F and 86°F. They typically occupy habitats 3 to 10 feet deep often near the edge of the water. Largemouth bass will feed on nearly everything around them, including crustaceans, frogs, and other fishes. Adults spawn after their second or third year, with the spawning season beginning when water temperatures reach 57°F to 61°F (13.9°C to 16°C), typically in April, and continuing until June. Males guard the nests. Eggs hatch within 2 to 5 days, and the sac fry remain near the nest for another 5 to 8 days (Moyle 2002).

Smallmouth bass prefer clearer, cooler water than largemouth bass, but can still be found in the same habitat as largemouth bass. Preferred summer water temperatures are from 68°F to 81°F (20°C to 27.2°C). The dominant food for these fish is crustaceans, aquatic insects, fish, and amphibians. They spawn after 3 or 4 years, in late spring when water temperatures reach 55°F to 61°F (12.8°C to 16.1°C). As with largemouth bass, males guard the nests. Eggs hatch in 3 to 10 days, and the young remain near the nest for another 3 to 4 days.

1.1.3 Aquatic Macroinvertebrates

Aquatic macroinvertebrates provide an important food base for many fish and wildlife species. In general, published information on the taxonomy, distribution, and abundance of macroinvertebrates in the Sacramento River drainage is limited. Current macroinvertebrate monitoring efforts on the Sacramento River have focused on large-basin scale patterns, and survey sites on the mainstem have been at various locations along the study reach. Under the Sacramento River Watershed Program, CDFW collected snag samples at two sites, one site near Colusa and one site near Hamilton City. Dominant taxa found in fall 1999 at the Hamilton City site include Orthocladiinae, Naididae, Ephemeroptera (*Baetis* and *Acentrella* sp.) and Trichoptera (*Hydropsyche* sp.) (Sacramento River Watershed Program 2002). Schaffter et al. (1983) found no significant difference in abundance of drifting invertebrates near riprapped and natural habitats on the Sacramento River. More than 50 percent of the drift was composed of chironomids, baetids, and aphids. Analysis of fish diets found the same three families in 72 percent of the guts sampled.

A large-scale monitoring effort in 2001 coordinated by DWR from Keswick Dam to Verona on the Sacramento River found that benthic macroinvertebrate diversity and richness decreased as the river moved downstream. Oligochaetes, chironomids, and mollusks became more prominent in this reach than in the reach from Keswick Dam to Red Bluff (Sacramento River Watershed Program 2002). More recently, the diurnal feeding habits of juvenile Chinook salmon in the upper Sacramento River (river mile (RM) 193 to RM 275) were examined in relation to drifting invertebrates by Petrusso and Hayes (2001). Chironomids and baetids dominated both the drift and stomach contents. Diets of 153 juvenile salmonids were examined; more than 63 percent of the diet was made up of chironomids of all life stages. Baetids comprised 14 percent of the total diet. It was concluded that based on measurements of mean stomach fullness and availability of drifting organisms, there was reasonable feeding opportunity during the sampling period in spring 1996. Mean drift densities ranged from 211 to 2,100 organisms per 100 cubic meters, with an overall mean of 617 organisms per 100 cubic meters (Petrusso and Hayes 2001). Daily mean drift density appeared to show no spatial patterns across the several sites sampled.

The constant flow of water in river systems provides an energetically convenient and economical way to disperse to new habitats; this movement downstream is known as drift. Some invertebrates passively enter the drift (e.g., benthic organisms may be entrained in the water column when a large current

sweeps through), and others exhibit active drift behavior (individuals actively enter the water column by voluntary actions) (Waters 1965, 1972; Müller 1974; Wiley and Kohler 1984). Macroinvertebrates drift to colonize new habitats (for dispersal of various life stages or to find suitable resources), or leave unsuitable habitats (in response to habitat quality or predation pressure). Drift is one of the most important downstream dispersal mechanisms for macroinvertebrates. Macroinvertebrates drift more commonly in the evening, usually at dusk (Waters 1972, Müller 1974, Wiley and Kohler 1984, Smock 1996).

Drifting invertebrates are the primary source of prey for juvenile fish, including salmonids (Chapman and Bjornn 1969). Juvenile Chinook salmon will often seek refuge in slow-velocity habitats where they can rest and drifting invertebrates will tend to be deposited.

In Shasta Lake, seasonal fluctuations in phytoplankton biomass regulate the abundance of the zooplankton, which form the base of the food chain for the lake's fisheries. Typically, the spring phytoplankton bloom peaks in late March and April at the on-set of thermal stratification, when nutrients are abundant in surface waters and available to the algae, and again in the fall coincident with the breakdown of the thermocline and mixing of the water column (Lieberman and Horn 1998). The zooplankton community of Shasta Lake is dominated by cladoceran and copepod species, with lower abundance of several rotifer species. Cladocera are most abundant during algae blooms and their abundance wanes, with a corresponding increase in copepod abundance, during the mid-summer (Lieberman and Horn 1998).

According to surveys conducted in 2011 and 2012 in tributaries to Shasta Lake, the benthic macroinvertebrates (BMI) communities, with a few exceptions, were generally indicative of good habitat and water quality conditions capable of supporting healthy functioning and productive ecosystems. The BMI community was largely dominated by cool/warm (eurythermal) taxa, which is expected as a function of the region's Mediterranean climate; taxa representing both pool and riffle specialists; and taxa representing the collector-filterer and collector-gatherer functional feeding guilds, which is also expected based on the relative position and trophic status of the tributary sampling sites within the watersheds (Reclamation 2014). Tributaries to the Sacramento River arm exhibited among the highest BMI abundances and taxa richness and diversity, although Pit River arm tributaries also exhibited relatively high taxa diversity. Tributaries in legacy mining districts immediately north of Shasta Dam and in portions of the Squaw Creek arm exhibited very depauperate BMI communities, with a high proportion of taxa tolerant of polluted conditions (Reclamation 2014).

A number of different aquatic mollusks (e.g., snails, limpets, mussels, and clams) are known to inhabit the principal tributaries and general vicinity of Shasta Lake, including several species of management importance (Frest and Johannes 1995, 1999; Howard 2010). Several species of hydrobiid "spring

snails” are known to inhabit the upper reaches of the Sacramento and McCloud rivers upstream from Shasta Lake (Frest and Johannes 1995, 1999) in spring complexes and associated headwater areas. These snails require clear, coldwater streams with cobbly gravel beds and tend to be associated with submergent vegetation; however, none of these species has been reported in the reaches of tributaries near Shasta Lake. A number of these spring snails and other stream-dwelling snails are ecologically important and used by the USFS for their survey and manage program (see Table 11-1 in Chapter 11 of the Environmental Impact Statement (EIS)).

The USFS sensitive freshwater mussel, the California floater (*Anodonta californiensis*), is also known historically to have occurred in Shasta Lake tributaries near the head of the lake (Howard 2010; J. Zustak, USFS, personal communication). However, surveys of historically occupied sites around Shasta Lake failed to find this species (Howard 2010), and it was not detected by casual surveys of the smaller perennial and intermittent tributaries to Shasta Lake surveyed in 2012 (Reclamation 2014). This species has experienced significant population declines throughout its range, primarily because of hydromodification of its habitat (Howard 2010). Its preferred habitat is unpolluted, slow-moving rivers and large streams, with beds composed of balanced mixtures of gravel, sand, and silt; however, California floaters are sometimes found in lake shore areas with stable water levels and suitable water currents and substrates (Pennak 1989). Other freshwater mollusks that are commonly observed in the tributaries of Shasta Lake include another freshwater mussel of the genus *Gonidea* and freshwater limpets of the genus *Lanx* (Howard 2010). The kneecap lanx (*Lanx Patelloides*) has been recently added to the USFS sensitive species list and is known to occur in the vicinity of the McCloud River Bridge. Another mollusk, Black juga (*Juga nigrina*) was recently added to the USFS sensitive species list. It was not detected during the 2014 field surveys but Shasta Lake and its tributaries are within the known range of this species (Cordeiro and Perez 2011). The western pearlshell (*Margaritifera falcata*) is also historically known from the McCloud River, but its close dependence on migratory salmonids for its life cycle has undoubtedly resulted in a decline in its abundance since construction of Shasta Dam blocked anadromous fish migrations (Howard 2010).

New Zealand Mudsnail

The New Zealand mudsnail (*Potamopyrgus antipodarum*), known to have been introduced to North America since about 1987 (Bowler 1991), was identified in Shasta Lake at the Bridge Bay Marina on September 10, 2007 (Benson and Kipp 2011). No New Zealand mudsnails were observed in the lake or in the vicinity of Shasta Lake tributaries during the 2011 and 2012 surveys (Reclamation 2014). New Zealand mudsnail have also been found lower in the Central Valley, including Sacramento River near Red Bluff, and the American, Mokelumne and Calaveras rivers (Benson 2011). This invasive aquatic mollusk is known from a number of other locations within California and can reach densities of more than 500,000 snails per square meter. Densities can fluctuate

seasonally, with lowest densities coinciding with the freezing winter months (Proctor et al. 2007). New Zealand mudsnails are highly effective competitors and predators of many native North American benthic macroinvertebrates, including other mollusks, crustaceans, and important aquatic insects. Predators of the New Zealand mudsnail include rainbow trout, brown trout, sculpins, and mountain whitefish (Proctor et al. 2007). Unfortunately, snails are capable of passing through the digestive system of fish alive and intact (Bondesen and Kaiser 1949).

Possible pathways of introduction into Shasta Lake include contaminated recreational watercraft and trailers and recreational water users (Proctor et al. 2007). Other vectors known to spread the snails, such as contaminated livestock, commercial ships, and dredging/mining equipment, are less likely in the case of Shasta Lake's recent invasion given the lack of commercial activities on the lake. If the particular clone detected in Shasta Lake is tolerant of the local conditions, a rapid colonization of the lake and its tributaries could occur through a variety of vectors.

The potential involvement of recreational watercraft and trailers and recreational water users in the translocation of New Zealand mudsnails between State waters is of immediate concern. Enlargement of Shasta Lake could provide a larger perimeter of shoreline accessibility for the snail, but not necessarily increase preferred lake habitats. In lakes in North America, New Zealand mudsnails do not commonly occupy shoreline habitats. Highest densities of New Zealand mudsnails occur between 20 to 25 meters in Lake Ontario (Proctor et al. 2007).

Quagga Mussel

Quagga mussels (*Dreissena bugensis*) and zebra mussels (*Dreissena polymorpha*) are invasive European aquatic mollusks introduced to North America in ship ballast water and first discovered in Lake Erie in 1989 (Spidle et al. 1994), have not been found in Shasta Lake, to date, but were discovered in California at Lake Havasu in 2007 (Cohen 2007). No quagga mussels were observed in the lake or in the vicinity of Shasta Lake tributaries during the 2011 and 2012 surveys (Reclamation 2014). CDFW has begun monitoring at Lake Shasta for adult mussels and veligers (S. Baumgartner, pers. comm., 2008). Possible pathways of introduction into Shasta Lake include contaminated recreational watercraft and trailers and recreational water users. The potential involvement of recreational watercraft and trailers and recreational water users in the translocation of dressenid mussels between State waters is of immediate concern. Enlargement of Shasta Lake could provide a greater area of deepwater and littoral habitat available for occupation by quagga and zebra mussels.

In a 2007 report produced for CDFW, Cohen (2007) described the temperature, calcium, pH, DO, and salinity tolerances of quagga mussels in an effort to assess the vulnerability of various California waters to invasion by quagga mussels and zebra mussels. Cohen identified calcium thresholds as the most

important environmental factor influencing distribution of zebra mussels in North America and applied similar thresholds for quagga mussels. In an investigation of the upper Sacramento River region, including Whiskeytown Reservoir and the watersheds above Shasta Dam, Cohen found that the McCloud River above Shasta Reservoir and the Pit River near Canby have the proper range of salinity, DO, temperature and calcium (at less than or equal to 12 milligrams per liter to be of low and moderate suitability to invasion by quagga mussels).

This page left blank intentionally.

Chapter 2 Impact Assessment

As described in Chapter 11 of the EIS, the SALMOD model was used to support technical analysis of Chinook salmon runs. Detailed modeling results are presented in Attachments 3 through 14 to the EIS Modeling Appendix as follows:

Modeling Appendix Attachment 3: Winter-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Future Conditions

Modeling Appendix Attachment 4: Winter-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Existing Conditions

Modeling Appendix Attachment 5: Winter-Run Chinook Salmon Production and Mortality from SALMOD AFRP Simulations Under Future Conditions

Modeling Appendix Attachment 6: Spring-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Future Conditions

Modeling Appendix Attachment 7: Spring-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Existing Conditions

Modeling Appendix Attachment 8: Spring-Run Chinook Salmon Production and Mortality from SALMOD AFRP Simulations Under Future Conditions

Modeling Appendix Attachment 9: Fall-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Future Conditions

Modeling Appendix Attachment 10: Fall-Run Chinook Salmon Production and Mortality from SALMOD 1999–2006 Average Simulations Under Existing Conditions

Modeling Appendix Attachment 11: Fall-Run Chinook Salmon Production and Mortality from SALMOD AFRP Simulations Under Future Conditions

Modeling Appendix Attachment 12: Late Fall-Run Chinook Salmon
Production and Mortality from SALMOD 1999–2006 Average
Simulations Under Future Conditions

Modeling Appendix Attachment 13: Late Fall-Run Chinook Salmon
Production and Mortality from SALMOD 1999–2006 Average
Simulations Under Existing Conditions

Modeling Appendix Attachment 14: Late Fall-Run Chinook Salmon
Production and Mortality from SALMOD AFRP Simulations Under
Future Conditions

Additional information on the fisheries, hydrology, and evaluation results for
Delta fisheries are presented in Attachment 1 to this Technical Report,
“Assessment of Fisheries Impacts within the Sacramento-San Joaquin
Delta.”

Chapter 3

References

3.1 Printed Sources

- Adams, P.B., C.B. Grimes, S.T. Lindley, and M.L. Moser. 2002. Status review for North American green sturgeon, *Acipenser medirostris*. NOAA, National Marine Fisheries Service, Southwest Fisheries Science Center, Santa Cruz, California. 49 p.
- Bachmann, Steve. 2000. Shasta Lake West Watershed Analysis: USDA Forest Service.
- Baker, P.F., and J.E. Morhardt. 2001. Survival of Chinook salmon smolts in the Sacramento-San Joaquin Delta and Pacific Ocean. Pages 163–182 in R. L. Brown, editor. Contributions to the biology of Central Valley salmonids. Fish Bulletin 179: Volume 2. California Department of Fish and Wildlife, Sacramento.
- Bartholow, J., R.B. Hanna, L. Saito, and M.J. Horn. 2001. Simulated limnological effects of the Shasta Lake temperature control device. *Environmental Management* 27 (4):609-626.
- Baxter, R., W. Harrell, and L. Grimaldo. 1996. 1995 Splittail Spawning Investigations. Interagency Ecological Program for the Sacramento-San Joaquin Estuary Newsletter 9(4):27–31.
- Beamesderfer, R.C., and M.A.H. Webb. 2002. Green Sturgeon Status Review Information.
- Beauchamp, D.A., M.F. Shepard, and G.B. Pauley. 1983. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Northwest)—Chinook Salmon. U.S. Fish and Wildlife Service Biological Report 82(11.6). U.S. Army Corps of Engineers, TR EL-82-4.
- Bell, M.C. 1990. Fisheries Handbook of Engineering Requirements and Biological Criteria. Fish Passage Development and Evaluation Program, U.S. Army Corps of Engineers, North Pacific Division. Portland, Oregon.
- Benson, A.J. 2011. New Zealand mudsnail sightings distribution. Retrieved 6/30/2011 from newzealandmudsnaildistribution.aspx.

- Benson, A.J., and R.M. Kipp. 2011. *Potamopyrgus antipodarum*. U.S. Geological Survey Nonindigenous Aquatic Species Database. Available: <<http://nas.er.usgs.gov/queries/factsheet.aspx?SpeciesID=1008>>. Revision Date: March 14, 2011.
- Bondesen, P., and E.W. Kaiser. 1949. Hydrobia (*Potamopyrgus*) jenkinsi (Smith) in Denmark illustrated by its ecology. *Oikos* 1 (252-281).
- Bowler, P.A. 1991. The rapid spread of the freshwater hydrobiid snail NZ mudsnail in the Middle Snake River, southern Idaho. *Proceedings of the Desert Fishes Council* 21:173-182.
- Brandes, P.L., and J.S. McLain. 2001. Juvenile Chinook salmon abundance, distribution, and survival in the Sacramento-San Joaquin Estuary. In: Brown, R.L., editor. *Contributions to the Biology of Central Valley Salmonids. Fish Bulletin* 179. Volume 2. Sacramento, California: California Department of Fish and Wildlife. Pp. 39-136. Available on website: http://www.delta.dfg.ca.gov/usfws/Report/SSJEFRO_Reports.html.
- Brannon, E.L., C.L. Melby, and S.D. Brewer. 1984. Columbia River White Sturgeon Enhancement. Final Report. (Contract No. DEAI79-84BP18952; Project No. 83 316.) Prepared for Bonneville Power Administration, Portland, Oregon.
- Buddington, R.K., and S.I. Doroshov. 1984. Feeding Trials with Hatchery Produced White Sturgeon Juveniles (*Acipenser transmontanus*). *Aquaculture* 36:237-243.
- Burau, J.R., J.W. Gartner, and M.T. Stacey. 1998. Results from the hydrodynamic element of the 1994 entrapment zone study in Suisun Bay. In: Report of the 1994 entrapment zone study, edited by Wim Kimmerer. Technical Report 56. Interagency Ecological Program for the San Francisco Bay-Delta Estuary. 55 pp.
- CALFED. *See* CALFED Bay-Delta Program.
- CALFED Bay-Delta Program. 2000. Multi-Species Conservation Strategy. Sacramento, California.
- California Department of Fish and Game. 1994. Petition to the California Board of Forestry to List Coho Salmon (*Oncorhynchus kisutch*) as a Sensitive Species. Sacramento, California.
- . 1998. Report to the Fish and Game Commission: A Status Review of the Spring-Run Chinook Salmon (*Oncorhynchus tshawytscha*) in the Sacramento River Drainage. Candidate Species Status Report 98-01. Inland Fisheries Division, Sacramento, California.

- . 2002 (March). Sacramento River Winter-Run Chinook Salmon Biennial Report, 2000–2001. Habitat Conservation Division.
- . 2007. Delta smelt.
<<http://www.delta.dfg.ca.gov/gallery/dsmelt.asp>> Accessed October 30, 2007.
- . 2011 (January). State and Federally Listed Endangered and Threatened Animals of California. Available: <<http://www.dfg.ca.gov/biogeodata/cnddb/pdfs/TEAnimals.pdf>> .
- California Department of Fish and Wildlife. 2014. GrandTab, California Central Valley Chinook Population Report. Fisheries Branch, Anadromous Resources Assessment. Sacramento, California. Available: <<http://www.calfish.org/tabid/213/Default.aspx>>.
- California Urban Water Agencies (CUWA). 1994 (March 7). Evaluation of potential effects of the proposed EPA salinity standard on the biological resources of the San Francisco Bay/Sacramento-San Joaquin Estuary (Draft). Prepared by R2 Resource Consultants, Inc. for The California Urban Water Agencies, Sacramento, California. Reference No. 5. 65 pp. plus appendices.
- Cantara Trustee Council. 2007. Final report on the recovery of the upper Sacramento River – subsequent to the 1991 Cantara spill. Cantara Trustee Council, Redding, California.
- Cech, J.J., S.I. Doroshov, G.P. Moberg, B.P. May, R.G. Schaffter, and D.M. Kohlhorst. 2000. Biological Assessment of Green Sturgeon in the Sacramento–San Joaquin Watershed (Phase 1). Final Report to CALFED Bay-Delta Program. Project No. 98-C-15, Contract No, B-81738.
- Center for Biological Diversity, The Bay Institute, The Natural Resources Defense Council. 2006. Emergency Petition to List the Delta Smelt (*Hypmesus transpacificus*) as an Endangered Species under the Endangered Species Act. Submitted to the U.S. Fish and Wildlife Service, Washington D.C. and Sacramento Field Office, California. Available: <http://www.biologicaldiversity.org/species/fish/Delta_smelt/pdfs/ds-endangered-petition-3-8-06.pdf>.
- . 2007. Petition to List the San Francisco Bay-Delta Population of Longfin Smelt (*Spirinchus thaleichtys*) as Endangered under the Endangered Species Act. Submitted to the U.S. Fish and Wildlife Service, Sacramento, California. Available: <http://www.biologicaldiversity.org/species/fish/longfin_smelt/pdfs/LF-S-petition-8-8-07.pdf>.

- Chapman, D.W., and T.C. Bjornn. 1969. Distribution of Salmonids in Streams, with Special Reference to Food and Feeding. Pages 153–176 in T.G. Northcote (ed.), Symposium on Salmon and Trout in Streams. H.R. MacMillan Lectures in Fisheries, University of British Columbia. Vancouver, British Columbia, Canada.
- Cohen, A.N. 2007. Potential distribution of zebra mussels (*Dreissena polymorpha*) and quagga mussels (*Dreissena bugensis*) in California, Phase 1 Report: Produced for the California Department of Fish and Wildlife by the San Francisco Estuary Report.
- Conte, F.S., S.I. Doroshov, P.B. Lutes, and E.M. Strange. 1988. Hatchery Manual for the White Sturgeon *Acipenser transmontanus* with Application to Other North American *Acipenseridae*. (Publication 3322.) University of California Cooperative Extension, Division of Agriculture and Natural Resources. Davis, California.
- Cramer S.P., and D.B. Demko. 1997. The Status of Late-Fall and Spring Chinook Salmon in the Sacramento River Basin Regarding the Endangered Species Act. Submitted to the National Marine Fisheries Service on behalf of the Association of California Water Agencies and California Urban Water Agencies.
- CUWA. *See* California Urban Water Agencies.
- Duke, S.D., T.J. Underwood, G.M. Asbridge, R.G. Griswold, M.J. Parsley, and L.G. Beckman. 1990. In A.A. Nigro (editor), Status and Habitat Requirements of White Sturgeon Populations in the Columbia River Downstream from McNary Dam. (Annual Progress Report.) Bonneville Power Administration. Portland, Oregon.
- Environmental Protection Information Center (EPIC), Center for Biological Diversity, and WaterKeepers Northern California. 2001. Petition to List the North American Green Sturgeon (*Acipenser medirostris*) as an Endangered or Threatened Species Under the Endangered Species Act.
- EPA. *See* U.S. Environmental Protection Agency.
- EPIC. *See* Environmental Protection Information Center.
- Finnell, L.M., and E.B. Reed. 1969. The diel vertical movements of Kokanee Salmon *Oncorhynchus nerka*, in Granby Reservoir, Colorado. Transactions of the American Fisheries Society 2:245-252.
- Fisher, F.W. 1994. Past and present status of Central Valley Chinook salmon. Conservation Biology 8(3):870-873.

- Frest, T.J., and E.J. Johannes. 1995. Freshwater mollusks of the upper Sacramento system, California, with particular reference to the Cantara Spill. 1994 Yearly report to California Department of Fish & Game. Deixis Consultants, Seattle, Washington.
- Frest, T.J., and E.J. Johannes. 1999. Field guide to survey and manage freshwater mollusk species. Bureau of Land Management, Oregon State Office and US Fish and Wildlife Service Regional Ecosystem Office, Portland, Oregon.
- Hallock, R.J., and F.W. Fisher. 1985. Status of the Winter-Run Chinook Salmon (*Oncorhynchus tshawytscha*) in the Sacramento River. Anadromous Fisheries Branch Office Report. California Department of Fish and Wildlife. Sacramento, California.
- Hassler, T.J. 1987. Species Profiles: Life Histories and Environmental Requirements of Coastal Fishes and Invertebrates (Pacific Southwest)—Coho Salmon. U.S. Fish and Wildlife Service Biological Report 82(11.70).
- Hieb, K., and R. Baxter. 1993. Delta outflow/San Francisco Bay. Pages 101—116 in P. L. Herrgesell (editor). 1991 Annual Report—Interagency Ecological Studies Program for the Sacramento-San Joaquin Estuary.
- Hill, K.A., and J.D. Webber. 1999. Butte Creek spring-run Chinook salmon, *Oncorhynchus tshawytscha*, Juvenile Outmigration and Life History, 1995–1998. California Department of Fish and Wildlife Inland Fisheries Administration. Report 99-5.
- Hinze, J.A., A.N. Culver, and G.V. Rice. 1956. Annual Report: Nimbus Salmon and Steelhead Hatchery, Fiscal Year 1955-56. California Department of Fish and Wildlife. Inland Fisheries Administrative Report Number 56-25. As cited in McCullough et al. 2001.
- Howard, J. 2010. Sensitive freshwater mussel surveys in the Pacific Southwest Region: assessment of conservation status. Prepared for the USDA Forest Service, Southwest Region. The Nature Conservancy, San Francisco, CA. 60 p.
- Jones & Stokes. 2001. Change in Potential Stranding Losses of Splittail and Juvenile Chinook Salmon in the Sacramento Bypass, 1999–2001. Final Report. Prepared for U.S. Army Corps of Engineers, Sacramento, California.
- Jones & Stokes Associates. 1993. Sutter Bypass Fisheries Technical Memorandum II: Potential Entrapment of Juvenile Chinook Salmon in the Proposed Gravel Mining Pond. Prepared for Teichert Aggregates, Sacramento, California.

- Kimmerer, W. 1992. An evaluation of existing data in the entrapment zone of the San Francisco Bay Estuary. Technical Report 3, Interagency Ecological Studies Program for the Sacramento-San Joaquin Estuary.
- Kimmerer W.J., J.R. Burau, and W.A. Bennett. 2002. Persistence of tidally-oriented vertical migration by zooplankton in a temperate estuary. *Estuaries* 23(3):359-371.
- Kjelson, M.A., P.F. Raquel, and F.W. Fisher. 1982. Life History of Fall-Run Juvenile Chinook Salmon, *Oncorhynchus tshawytscha*, in the Sacramento-San Joaquin Estuary, California. Pages 393–411 in V. S. Kennedy (editor), *Estuarine Comparisons*. Academic Press. New York, New York.
- Kohlhorst, D.W. 1976. Sturgeon Spawning in the Sacramento River in 1973, as Determined by Distribution of Larvae. *California Fish and Game* 62:32–40.
- Koski, M. and B.M. Johnson. 2002. Functional response of kokanee (*Oncorhynchus nerka*) feeding on *Daphnia* under a range of light intensities. *Canadian Journal of Fisheries and Aquatic Sciences* 59(4): 707-716.
- Lee, D. P. 1999. Water level fluctuation criteria for black bass in California reservoirs. California Department of Fish and Game, Fisheries Program Branch, Reservoir Research and Management Project – Information Leaflet No. 12.
- Lieberman D.M. and M.J. Horn. 1998. Pre- and post-operational effects of a temperature control device on physical, chemical, and biological attributes of Shasta Lake, California: phase 1, spring 1995 through fall 1997. Open file report 98-251. Denver (CO): US Geological Survey.
- Marine, K.R., and J.J. Cech, Jr. 2004. Effects of High Water Temperatures on Growth, Smoltification, and Predator Avoidance in Juvenile Sacramento River Chinook Salmon. *North American Journal of Fisheries Management* 24:198–210.
- McCullough, D.A. 1999. A Review and Synthesis of Effects of Alterations to the Water Temperature Regime on Freshwater Life Stages of Salmonids, with Special Reference to Chinook Salmon. Report No. EPA 910-R-99-010. U.S. Environmental Protection Agency, Region 10. Seattle, Washington.
- McCullough, D.A., S. Spalding, D. Sturdevant, and M. Hicks. 2001. Summary of Technical Literature Examining the Physiological Effects of Temperature on Salmonids—Issue Paper 5. Report No. EPA-910-D-01-005. U.S. Environmental Protection Agency.

- McKechnie, R.J., and R.B. Fenner. 1971. Food Habits of White Sturgeon, *Acipenser transmontanus*, in San Pablo and Suisun Bays, California. *California Fish and Game* 57:209–212.
- Miller, L.W. 1972. Migrations of Sturgeon Tagged in the Sacramento–San Joaquin Estuary. *California Fish and Game* 58(2):102–106.
- Miller, D.J., and R.N. Lea. 1972. Guide to the Coastal Marine Fishes of California. (Fish Bulletin 172.) California Department of Fish and Wildlife. Sacramento, California.
- Miller, A.I., P.J. Anders, M.J. Parsley, C.R. Sprague, J.J. Warren, and L.G. Beckman. 1991. Report C, in A.A. Nigro (editor), Status and Habitat Requirements of White Sturgeon Populations in the Columbia River downstream from McNary Dam. (Annual Progress Report.) Bonneville Power Administration. Portland, Oregon.
- Mills, T.J., and F. Fisher. 1994. Central Valley Anadromous Sport Fish Annual Run-Size, Harvest, and Population Estimates, 1967 through 1991. Inland Fisheries Technical Report. California Department of Fish and Wildlife.
- Moyle, P.B. 2002. *Inland Fishes of California*. University of California Press. Berkeley, California.
- Moyle, P.B., P.J. Foley, and R.M. Yoshiyama. 1992. Status of Green Sturgeon, *Acipenser medirostris*, in California. Final report submitted to National Marine Fisheries Service. University of California, Davis.
- Moyle, P.B., R.D. Baxter, T. Sommer, T.C. Foin, and S.A. Matern. 2004. Biology and population dynamics of the Sacramento splittail (*Pogonichthys macrolepidotus*) in the San Francisco Estuary: A Review. *San Francisco Estuary and Watershed Science* [online serial]. Vol 2, Issue 2 (May 2004), Article 3.
[Http://repositories.cdlib.org/jmie/sfews/vol2/iss2/art3](http://repositories.cdlib.org/jmie/sfews/vol2/iss2/art3).
- Moyle, P.B., R.M. Yoshiyama, J.E. Williams, and E.D. Wikramanayake. 1995. *Fish Species of Special Concern in California*. Second edition. California Department of Fish and Wildlife. Sacramento, California.
- Müller, K. 1974. Stream Drift as a Chronobiological Phenomenon in Running Water Ecosystems. *Annual Review of Ecology and Systematics* 5:309–323.
- Naiman, R.J. 1998. Biotic stream classification. Pages 97-119 in R.J. Naiman and R.E. Bilby (editors). *River ecology and management: lessons from the Pacific coastal region*. Springer, New York.

- National Marine Fisheries Service. 1993. Biological Opinion for the Operation of the Federal Central Valley Project and the California State Water Project. Southwest Region. Long Beach, California.
- National Marine Fisheries Service. 1996. Status Review of West Coast Steelhead from Washington, Idaho, Oregon, and California. NOAA Technical Memorandum NMFS-NWFSC-27. National Marine Fisheries Service, Northwest Fisheries Science Center. Seattle, Washington. August.
- . 1997. Status Review of West Coast Steelhead from Washington, Idaho, Oregon, and California. NOAA Technical Memorandum NMFS-NWFSC-27.
- . 2000. Biological Opinion for the Proposed Operation of the Federal Central Valley Project and the State Water Project for December 1, 1999 through March 31, 2000. Northwest and Southwest Regional Sustainable Fisheries Divisions.
- . 2002. Biological Opinion on Interim Operations of the Central Valley Project and State Water Project Between April 1, 2002 and March 31, 2004. Southwest Region. Long Beach, California.
- . 2014 (July). Recovery Plan for the Evolutionarily Significant Units of Sacramento River Winter-run Chinook Salmon and Central Valley Spring-run Chinook Salmon and the Distinct Population Segment of Central Valley Steelhead. California Central Valley Office. Sacramento, California.
- Nevarés, S., and R. Liebig. Hardhead surveys in the lower McCloud River, 2007. Technical memorandum 8. December 2007. McCloud-Pit Project (FERC Project No. 2106). Pacific Gas and Electric Company. 5 p.
- NMFS. See National Marine Fisheries Service.
- Orr, E.L., W.N. Orr, and E.M. Baldwin, 1992. Geology of Oregon, Fourth Edition
- Pacific Fishery Management Council. 2002. Review of 2001 Ocean Salmon Fisheries. Document prepared for the Council and its advisory entities. Portland, OR. Pacific States Marine Fisheries Commission. 1992. White Sturgeon Management Framework Plan. Portland, Oregon.
- Pacific Gas and Electric Company. 2007. Hardhead surveys. Technical memorandum.
- Pacific States Marine Fisheries Commission. 1992. 45th Annual Report of the Pacific States Marine Fisheries Commission for the Year 1992.

- Pennak, R.W. 1989. Freshwater invertebrates of the United States, 3rd edition. Protozoa to Mollusca. John Wiley and Sons, New York. 628 p.
- Petrusso, P.A., and D.B. Hayes. 2001. Invertebrate Drift and Feeding Habits of Juvenile Chinook Salmon in the Upper Sacramento River, California. *California Fish and Game* 87(1):1–18.
- Ploskey, G.R. 1986. Impacts of water-level changes on reservoir ecosystems, with implications for fisheries management. In *Reservoir Fisheries Management: Strategies for the 80s*, G. E. Hall and M. J. Van Den Avyle (editors). Bethesda, Maryland: American Fisheries Society, Southern Division.
- PFMC. See Pacific Fishery Management Council.
- Proctor, T., B. Kerans, and P. Clancey. 2007. National Management and Control Plan for the New Zealand mudsnail *Potamopyrgus antipodarum*: Prepared for the Aquatic Nuisance Species Task Force by the New Zealand Mudsnail Management and Control Plan Working Group.
- PSMFC. See Pacific States Marine Fisheries Commission.
- Quinn, T.P. 2005. *The Behavior and Ecology of Pacific Salmon and Trout*. University of Washington Press. Seattle, Washington.
- Radtke, L.D. 1966. Distribution of Smelt, Juvenile Sturgeon, and Starry Flounder in the Sacramento–San Joaquin Delta with Observations on Food of Sturgeon. In J. L. Turner and D. W. Kelly (editors), *Ecological Studies of the Sacramento–San Joaquin Estuary. Part II, Fishes of the Delta*. California Department of Fish and Wildlife Fish Bulletin 136:115-129.
- Raleigh, R.F., T. Hickman, R.C. Soloman, and P.C. Nelson. 1984. Habitat Suitability Information: Rainbow Trout. (Biological Report 82 (10.60)) U.S. Fish and Wildlife Service. Washington, D.C. (FWS/OBS-82/10.60.)
- Ratcliff, D.R. 2006. Evaluating the impactiveness of grass bed treatments as a habitat for juvenile bass in a drawdown reservoir. Master's thesis, Utah State University, Logan, Utah.
- Reclamation. See U.S. Department of the Interior, Bureau of Reclamation.
- Reeves, G.H., P.A. Bisson, and J.M Dambacher. 1998. Fish communities. Pages 200–234 in R.J. Naiman and R.E. Bilby (editors). *River ecology and management: lessons from the Pacific coastal region*. Springer, New York.

- Reiser, D.W., and T.C. Bjornn. 1979. 1. Habitat Requirements of Anadromous Salmonids. In W.R. Meehan (technical editor), Influence of Forest and Rangeland Management on Anadromous Fish Habitat in the Western United States and Canada. U.S. Forest Service GTR PNW-96.
- Rode, M. 1989. Administrative draft – California wild trout management program, McCloud River wild trout area management plan: California Department of Fish and Wildlife, Inland Fisheries.
- Rode, M., and M. Dean. 2004. Lower McCloud River wild trout area fishery management plan 2004 through 2009. Redding: California Department of Fish and Wildlife, Northern California – North Coast Region.
- Sacramento River Conservation Area Forum. 2003 (September). Sacramento River Conservation Area Forum Handbook. Red Bluff, CA. Prepared for the Resources Agency under Senate Bill 1086, authored by Senator Jim Nielsen.
- Sacramento River Watershed Program. 2002. 2000–2001 Annual Monitoring Report. June. Available: <http://www.sacriver.org/subcommittees/monitoring/documents/SRWP_AMR_00-01_FINAL.pdf>
- Saito, L., B.M. Johnson, J. Bartholow, and R. Blair Hanna. 2001. Assessing ecosystem effects of reservoir operations using food web-energy transfer and water quality models. *Ecosystems* 4:105-125.
- Sandercock, F.K. 1991. Life History of Coho Salmon (*Oncorhynchus kisutch*). Pages 395–446 in C. Groot and L. Margolis (eds.), *Pacific Salmon Life Histories*. University of British Columbia Press. Vancouver, BC, Canada.
- Schaffter, R.G., P.A. Jones, and J.G. Karlton. 1983. Sacramento River and Tributaries Bank Protection and Erosion Control Investigation—Evaluation of Impacts on Fisheries. California Department of Fish and Wildlife. Sacramento. Prepared for U.S. Army Corps of Engineers Sacramento District.
- Shapovalov, L., and A.C. Taft. 1954. The Life Histories of Steelhead Rainbow Trout (*Salmo Gairdneri gairdneri*) and Silver Salmon (*Oncorhynchus kisutch*) with Special Reference to Waddel Creek, California, and Recommendations Regarding Their Management. California Department of Fish and Wildlife, Fish Bulletin 98.
- Slater, D.W. 1963. Winter-Run Chinook Salmon in the Sacramento River, California with Notes on Water Temperature Requirements at Spawning. U.S. Fish and Wildlife Service. Washington, D.C.

- Smith, L.H. 1987. A review of circulation and mixing studies of San Francisco Bay, California. (Circular 1015). U.S. Geological Survey. Denver, Colorado.
- Smock, L.A. 1996. Macroinvertebrate Movements: Drift, Colonization and Emergence. Chapter 17 in F.R. Hauer and G.A. Lamberti (editors), *Methods in Stream Ecology*. Academic Press. San Diego, California.
- Sommer, T.R., R. Baxter, and B. Herbold. 1997. Resilience of Splittail in the Sacramento–San Joaquin Estuary. *Transactions of the American Fisheries Society* 126:961–976.
- Sommer, T.R., L. Conrad, G. O’Leary, F. Feyrer, and W.C. Harrell. 2002. Spawning and Rearing of Splittail in a Model Floodplain Wetland. *Transactions of the American Fisheries Society* 131:966–974.
- Sommer, T.R., W.C. Harrell, M.L. Nobriga and R. Kurth. 2003. Floodplain as habitat for native fish: Lessons from California’s Yolo Bypass. Pages 81–87 In P.M. Faber, editor. *California riparian systems: Processes and floodplain management, ecology, and restoration*. 2001 Riparian Habitat and Floodplains Conference Proceedings, Riparian Habitat Joint Venture, Sacramento, California.
- Spidle, A.P., J.E. Marsden, and B. May. 1994. Identification of the Great Lakes Quagga Mussel as *Dreissena bugensis* from the Dnieper River, Ukraine, on the basis of allozyme variation. *Canadian Journal of Fisheries and Aquatic Science* 51:1485-1489.
- SRCAF. *See* Sacramento River Conservation Area Forum.
- SRWP. *See* Sacramento River Watershed Program.
- Stables, T.B., G.L. Thomas, S.L. Thiesfeld, and B.G. Pauley. 1990. Effects of reservoir enlargement and other factors on the yield of wild rainbow and cutthroat trout in Spada Lake, Washington. *North American Journal of Fisheries Management* 10:305-314.
- State Water Board. *See* State Water Resources Control Board.
- State Water Resources Control Board. 1999. Final Environmental Impact Report for Implementation of the 1995 Bay/Delta Water Quality Control Plan.
- Stuber, R.J., G. Gebhart, and O.E. Maughan. 1982. Habitat suitability index models: Largemouth bass: U.S. Department of the Interior, Fish and Wildlife Service.

- The Bay Institute et al. 2007 (February 7). Petition to the State of California Fish and Game Commission and Supporting Information for Listing the Delta Smelt as an Endangered Species under the California Endangered Species Act.
- Unger, P.A. 1994. Quantifying salinity habitat of estuarine species. Newsletter. Interagency Ecological Program for the Sacramento-San Joaquin Estuary. Autumn 1994:7-10.
- USACE and The Reclamation Board. See U.S. Army Corps of Engineers and State of California Reclamation Board.
- USFS. *See* U.S. Forest Service.
- USFWS. *See* U.S. Fish and Wildlife Service
- U.S. Army Corps of Engineers and The California Reclamation Board. 2002 (December). Sacramento and San Joaquin River Basins California Comprehensive Study. Interim report. Sacramento District, Corps of Engineers and The Reclamation Board, Sacramento, California.
- U.S. Department of the Interior, Bureau of Reclamation. 1986. Central Valley Fish and Wildlife Management Study: Temperature and Flow Studies for Optimizing Chinook Salmon Production, Upper Sacramento River, California. Special report. Sacramento, California.
- . 2003. Shasta Lake Water Resources Investigation, Mission Statement Milestone Report. Mid-Pacific Region. Sacramento, California.
- . 2004. Shasta Lake Water Resources Investigation Initial Alternatives Information Report. Mid-Pacific Region. Sacramento, California.
- . 2007. Draft Shasta Lake Water Resources Investigation Plan Formulation Report. Mid-Pacific Region. Sacramento, California.
- . 2008 (August). Biological Assessment on the Continued Long-term Operation of the Central Valley Project and the State Water Project. U.S. Department of the Interior, Mid-Pacific Region, Sacramento, California.
- . 2014 (April). Tributary Fisheries Characterization Report. U.S. Department of the Interior, Mid-Pacific Region, Sacramento, California.
- U.S. Environmental Protection Agency. 1993. Draft Regulatory Impact Assessment of the Proposed Water Quality Standards for the San Francisco Bay/Delta and Critical Habitat Requirements for the Delta Smelt. December.

- U.S. Fish and Wildlife Service (USFWS). 1995. Working Paper: Habitat Restoration Actions to Double Natural Production of Anadromous Fish in the Central Valley of California. Vol. 1, 2, 3. Prepared by Anadromous Fish and Restoration Core Group. Stockton, California.
- . 1996. Recovery Plan for the Sacramento-San Joaquin Delta Native Fishes. U.S. Fish and Wildlife Service, Portland, Oregon.
- . 2008 (December). Formal Endangered Species Act Consultation on the Proposed Operations of the Central Valley Project (CVP) and State Water Project (SWP). Final. Sacramento, California.
- U.S. Fish and Wildlife Service and Hoopa Valley Tribe. 1999. Trinity River Flow Evaluation Final Report.
- Vogel, D.A., and K.R. Marine. 1991. Guide to Upper Sacramento River Chinook Salmon Life History. U.S. Department of the Interior, Bureau of Reclamation, Central Valley Project. CH2M Hill, Redding, California.
- Wang, J.C.S. 1986. Fishes of the Sacramento–San Joaquin Estuary and Adjacent Waters, California: A Guide to the Early Life Histories. IEP Technical Report No. 9. California Department of Water Resources, California Department of Fish and Wildlife, U.S. Department of the Interior, Bureau of Reclamation, and U.S. Fish and Wildlife Service.
- . 2006. Early life history comparison of the green sturgeon, *Acipenser medirostris*, and white sturgeon, *Acipenser transmontanus*, of the Sacramento-San Joaquin River Delta, California. Tracy Technical Bulliten 2006-1. 21p.
- Waters, T.F. 1965. Interpretation of Invertebrate Drift in Streams. *Ecology* 46:327–334.
- . 1972. The Drift of Stream Insects. *Annual Review of Entomology* 17:253–272.
- Weaver, J., and S. Mehalik. 2008. Upper Sacramento River summary report–August 4-5, 2008. California Department of Fish and Game, Heritage and Wild Trout Program, Sacramento, California. 12p.
- Weidlein, W.D. 1971. Summary progress report on the Shasta Lake Trout management investigations, 1967 through 1970. Redding, California: California Department of Fish and Wildlife, Inland Fisheries. Inland Fisheries Administrative Report 71-13.

- Weitkamp, L.A., T.C. Wainwright, G.J. Bryant, G.B. Milner, D.J. Teel, R.G. Kope, and R.S. Waples. 1995 (September). Status Review of Coho Salmon from Washington, Oregon, and California. NOAA Technical Memorandum NMFS-NWFSC-24.
- Wiley, M.J., and S.L. Kohler. 1984. Behavioral Adaptations of Aquatic Insects. In V. Resh and D. Rosenberg (editors), *The Ecology of Aquatic Insects*. Praeger Press, CBS Inc. New York, New York.
- Yoshiyama, R.M., F.W. Fisher, and P.B. Moyle. 1998. Historical Abundance and Decline of Chinook Salmon in the Central Valley Region of California. *North American Journal of Fisheries Management* 18:487–521.
- Young, P.S., and J.J. Cech, Jr. 1996. Environmental tolerances and requirements of splittail. *Transactions of the American Fisheries Society*. 125:664-678.
- Zedonis, P.A., and T.J. Newcomb. 1997. An Evaluation of Flow and Water Temperatures during the Spring for Protection of Salmon and Steelhead Smolts in the Trinity River, California. U.S. Fish and Wildlife Service, Coastal California Fish and Wildlife Office. Arcata, California.

3.2 Personal Communications

- Baumgartner, S. 2008.
- Baumgartner, S. and CDFG. 2006.
- Hanson, C. 2009 (January). Presentation at the U.S. Department of the Interior, Bureau of Reclamation Mid-Pacific Water Users Conference. Reno, Nevada.
- Zustak, Joe. 2007.