South Fork Clearwater: Summary of CHaMP Metrics and Modeled Products

- CHaMP metrics are directly measured at each CHaMP site according to probabilistic sampling design. Metric included in this summary are:
  - Sinuosity
  - Substrate < 2 mm
  - Substrate < 6 mm
- Modeled products are built from CHaMP data and additional mechanistic or empirical knowledge, assessed at each CHaMP Site
  - Habitat Suitability Index (HSI)
  - Net Rate of Energy Investment (NREI)

Summary by Steelhead Assessment Unit

South Fork Clearwater: Summary of CHaMP Metrics and Modeled Products

Steelhead Intrinsic Potential



Steelhead Juvenile Weighted Usable Area

- Source: Habitat Suitability Index (HSI) Model
- Definition: Suitable habitat area (m^2) per meter of stream length for Juvenile Steelhead



Steelhead Spawner Weighted Usable Area

- Source: Habitat Suitability Index (HSI) Model
- Definition: Suitable Spawning habitat area (m^2) per meter of stream length for Steelhead



Juvenile Salmonid Capacity

- Source: Net Rate of Energy Intake (NREI) Model
- Definition: Carrying capacity (fish) per meter of stream length



Sinuosity

- Source: CHaMP Metrics (www.champmonitoring.org)
- Definition: Ratio of the thalweg length to the straight line distance between the start and end points of the thalweg.



Substrate < 2 mm

- Source: CHaMP Metrics (www.champmonitoring.org)
- Definition: Average percentage of pool tail substrates comprised of fine sediment <2 mm



Substrate < 6 mm

- Source: CHaMP Metrics (www.champmonitoring.org)
- Definition: Average percentage of pool tail substrates comprised of fine sediment <6 mm




QRF Capacity Estimates
Juvenile Chinook
Fish per meter
< 1.5
1.5 to 3.0
3.0 to 4.5
4.5 to 6.0
> 6.0
N
0
5
10 Km

FLOODPLAIN CONDITION ASSESSMENT

Considering riparian conditon, land use intensity & floodplain accessibility



FLOODPLAIN CONDITION ASSESSMENT

Considering riparian conditon, land use intensity & floodplain accessibility



SCS 1:



SCS 2:



SCS 3:



SCS 4:



SCS 5:



Floodplain Condition

Status

- Poor
- Moderate
- Good
- Intact

FLOODPLAIN CONDITION ASSESSMENT

Considering riparian conditon, land use intensity & floodplain accessibility



SCS 6:

SCS 7:



SCS 8:



SCS 9:



SCS 10:



Floodplain Condition

Status

- Poor
- Moderate
- Good
- Intact

LARGE WOODY DEBRIS RECRUITMENT POTENTIAL



LWD Recruitment Potential

Probability of LWD Recruitment

- Minimal < 33%
- Moderate 33% to 66%
- High 66% to 90%
- Very High > 90%

Sources: Esri, DeLorme, USGS, NPS

LARGE WOODY DEBRIS RECRUITMENT POTENTIAL



SCS 1:



SCS 2:



SCS 3:

SCS 4:



SCS 5:



LWD Recruitment Potential

Probability of LWD Recruitment

- Minimal < 33%
- Moderate 33% to 66%
- High 66% to 90%
- Very High > 90%

LARGE WOODY DEBRIS RECRUITMENT POTENTIAL



SCS 6:

SCS 7:



SCS 8:



SCS 9:

SCS 10:

LWD Recruitment Potential

Probability of LWD Recruitment

- Minimal < 33%
- Moderate 33% to 66%
- High 66% to 90%
- Very High > 90%

RIPARIAN CONDITION ASSESSMENT

Estimate of current departure from historic within valley bottom



RVCA - Riparian Vegetation Condition Assessment

Current Departure from Historic

- Large Departure: > 66%
- Significant Departure: 33% to 66%
- Minor Departure: 10% to 33%
- Negligable Departure: < 10%

Sources: Esri, DeLorme, USGS, NPS

RIPARIAN CONDITION ASSESSMENT

Estimate of current departure from historic within valley bottom



SCS 1:



SCS 2:



SCS 3:



SCS 4:



SCS 5:



RVCA - Riparian Vegetation Condition Assessment

Current Departure from Historic

- Large Departure: > 66%
- Significant Departure: 33% to 66%
- Minor Departure: 10% to 33%
- Negligable Departure: < 10%

RIPARIAN CONDITION ASSESSMENT

Estimate of current departure from historic within valley bottom



SCS 6:



SCS 7:



SCS 8:



SCS 9:



SCS 10:



RVCA - Riparian Vegetation Condition Assessment

Current Departure from Historic

- Large Departure: > 66%
- Significant Departure: 33% to 66%
- Minor Departure: 10% to 33%
- Negligable Departure: < 10%

RIPARIAN CONVERSION

Causes for departure from historic



RVCA - Riparian Conversion

Historic to Existing Riparian Conversion Type

- Conifer Encroachment
- Conversion to Barren
- Conversion to Developed
- Riparian to Riparian
- Upland Encroachment

Sources: Esri, DeLorme, USGS, NPS