

Yakima River Reaches: Instream Flow Improvement Matrix

River Reach	Problem	Flow Objective	Priority	Potential Projects	Other Notes
Keechelus Dam to Lake Easton	Flow too high in July, Aug & 1 st week of-Sept; over 800 cfs	Improve summer rearing by reducing flows down to 450-550 cfs. Increase winter flow to 120 cfs (connection to side channels at that flow). Provide pulses in winter.	High	K to K Pipeline Wymer storage downstream of Keechelus Aquifer storage	Spring is probably okay
Kachess River	No change proposed – lesser priority for improving river flow because of other objectives				
Easton Reach	Spring – need outmigration flow for spring Chinook	1000 cfs for 48 hours during dry years, augment spring Q for channel maintenance occasionally (5-yr for riparian recruitment – bank full)	Medium	Wymer Aquifer storage	Uncertainties: Don't know fish usage May be fish in future? Look at pit-tag relationship to determine pulse size/duration
	Fall/Winter – need additional flow for spawning and rearing	Currently 180 cfs, start spawning flow at 220 cfs, increase to 250-300 cfs in winter, 250 cfs provides connection to side channels. Spawning flows at 220 cfs.	High		
Cle Elum River	Summer flows (July and August) are too high	Reduce flow, modify flip flop to give more gentle change in hydrograph. In wet years, hold water back in August and reduce flow (maybe to 1000 cfs)	High	Bumping Wymer Flip / flop modification/relax Aquifer storage K to K Cle Elum pool raise	This reach is ripe for restoration as floodplain ownership is held in conservation easements. One-third of spring Chinook population spawns here.
	Fall/Winter Flows (September 10 through March): no flow variation (sp. Chinook, steelhead)	Increase to 500 cfs September through March, side channels are thought to be activated around 500 cfs, two were recently modified to activate at 200 cfs, provide pulse flows.			
Cle Elum to Teanaway River	Summer flows are too high	Reduce flows from 4000 cfs to 1000 cfs by late August. Ok to have high flow in July, as mimics unregulated hydrograph.	High	See Cle Elum list	Spring flows support cottonwood regeneration

River Reach	Problem	Flow Objective	Priority	Potential Projects	Other Notes
	Need Channel shaping flows in spring occasionally for riparian restoration	Provide channel shaping flows every 5 years or so.	Medium		
	In winter, a flat hydrograph exists	Provide flow variability, see Cle Elum River.	Medium		
Teanaway to Roza Dam	Summer flows are too high	Reduce flows	High	See Cle Elum list	There are tributaries in this reach which reduces effects in spring and winter
	Need channel shaping flows in spring occasionally for riparian restoration Not as a big as issue as upstream reach because of tributary inflow	Provide channel shaping flows every 5 years or so	Medium		
	Some flow variability exists because of tributaries but magnitude could be increased	Provide flow variability, time pulses to match natural events.	Medium		
Roza-Naches	In Spring from beginning of March to end of May, need additional outmigration flow	Increase flow to about 1400 cfs from March through May.	High	See Cle Elum list Roza hydropower subordination Roza dam removal	Predation Issues: Late fall to early winter At 400 cfs to support power production Uncertainties: Opportunity to support reintroduced sockeye and
	In Fall/winter there needs to be sufficient flow to support movement of fish to lower river, rearing and spawning	Increase to 1000-1400 cfs (use IFTAG flows). Link flows to habitat needs. Compare to 2-D habitat model for reach above Roza Dam.	High		

River Reach	Problem	Flow Objective	Priority	Potential Projects	Other Notes
	In Summer flow is not an issue because of Roza diversions but ability to modify flow in this reach may be useful – talk to habitat subcommittee to determine	Variability (?)	Low to medium		summer Chinook Consider three potential scenarios: 1) Current operations 2) Without operating power plant 3) With subordination Try to more closely mimic unregulated flow during subordination period(s)
Bumping Dam— Lower Naches	Fall flows after flip-flop are too high, then get reduced in winter	Reduce flows by 70-100 cfs from August through October	Medium		Bumping Expansion, change operations in Naches arm
Tieton River	Low winter flows and no variation (November to March)	Increase to 275 cfs to improve winter bull trout conditions. (NMFS had recommended 125 cfs, unregulated = 400-450 cfs)	High	Change in operations (minor improvements on shoulders of flip-flop)	Current winter flows (75 – 120 cfs)
	High flows in September due to flip-flop (and shoulders)	Reduce flows as much as possible.	Medium	Contingent on South Fork Fish Passage Project to allow more flexibility in reservoir operations	Key project is South Fork Fish Passage Project, needs to be constructed to allow flexibility in reservoir operations
Lower Naches River	Summer flows are low, ramping rate from high spring flows to summer flows is abrupt, affects rearing for steelhead, coho, spring Chinook. Up to ½ of flow in river is lost to groundwater in part of this reach.	Change ramping rate from spring to summer, increase summer low flow, check habitat needs vs flow.	High	Change operations to improve Water Conservation for Glead, Naches Selah, YU Canal (non-YRBWEP projects) to improve summer flow	Uncertainties: Gaining/losing reaches Complexity Limited recharge Try to stay below flow level that affects cottonwood regeneration

River Reach	Problem	Flow Objective	Priority	Potential Projects	Other Notes
	High flows in flip-flop operations	Reduce flows as much as possible, look at releasing more in summer and reducing flip flop.	High		
Yakima River from Naches River to Parker	High summer flow	Reduce as much as possible	Low	Habitat projects	Habitat improvements will be primary enhancement strategy
Yakima River from Parker to Toppenish Creek (Wapato reach)	In spring, flow is low and outmigration flow needed, mostly during dry years	15,000 – 20,000 acre-feet to use specifically for smolt outmigration. See SOAC recommendations for pulse flows. Maybe early and late pulse? Sockeye passage also?	High	Water Conservation primarily. Secondly Water Storage in Wymer, Cle Elum, Kachess Dead Storage, Bumping, Modify operations	Temperature issues with shoulders of spring and fall Fit channel to river
	Low Summer flow	link to habitat needs	TBD		
Yakima River between Toppenish Creek to Prosser Dam	In spring, flow is low (similar to Wapato Reach)	see Wapato Reach	TBD	See Wapato Reach	Summer and Winter flow—OK
Yakima River—Chandler Reach	Need more flow in July (shoulder period) and September	Need greater than 1000 cfs in September	Low	Chandler Power Plant subordination, KID projects See Wapato Reach for more projects	Biggest issue: mortality at canal Winter flow—OK
	Need more flow in Spring	Although some subordination occurs to provide 1000 cfs, need more flow	Low		
Lower Yakima River (Chandler Powerplant to mouth)	In spring, flow is low (similar to Wapato Reach) but more emphasis in June needed to push fish out	see Wapato Reach	Low	See Wapato Reach KID	Winter—OK
	in Summer flow can be low, cover is an issue	link to habitat needs	Low		

Tributaries					
Big, Little, Manastash			TBD	Deliver water directly to tribs if supply replacement not feasible	See IFTAG for Manastash Creek Discuss with habitat group
Ahtanum	Urbanization, irrigation conservation are issues		TBD		Water rights concerns will limit ability to implement any projects in Ahtanum Creek Basin
Wenas	Need to redo irrigation diversions—will there be water for streamflow		TBD		
North Side Kittitas Valley Tributaries	Fish barriers are big issue		TBD	Thorp to Wymer using KRD North Branch Canal could serve water users	Need to figure out creek systems—too many distributaries. May need to simplify the systems to keep enough water in stream