

**Glen Canyon Dam Adaptive Management Work Group**  
**Agenda Item Information**  
**August 24-25, 2011**

---

---

Agenda Item

Species of Concern Ad Hoc Committee Report

---

---

Action Requested

- ✓ Motion requested. The following motion is recommended by TWG. However, no motion is officially made unless and until an AMWG member makes the motion in accordance with the AMWG Operating Procedures.

AMWG accepts the report of the Species of Concern Ad Hoc Committee entitled, “Assessment of Taxa of Management Concern in the Colorado River Ecosystem, Glen and Grand Canyons, Arizona, USA: Habitat Needs, Availability, and Ecosystem Roles” dated 15 June 2011 (revised July 8, 2011), which meets the criteria set forth in the AMWG motion of April 2009, calling for “a review of information about an assessment of the status of habitat needs and availability, and ecosystem roles” of native species of management concern.

---

---

Presenters

Shane Capron, Technical Work Group Chair (Western Area Power Administration)

Larry Stevens, Species of Concern Ad Hoc Committee Chair; AMWG member, Grand Canyon  
Wildlands Council

---

---

Previous Action Taken

- ✓ By AMWG: At its April 2009 meeting, AMWG passed the following motion by consensus. In recognition of GCDAMP goals toward management of the Colorado River through an ecosystem approach, AMWG directs the TWG to establish the Species of Concern Ad Hoc Committee (SCAHC) and requests the participation of GCMRC in that ad hoc committee, to produce a draft report to be presented to AMWG on or before by May 1, 2011, that contains the following with regard to species of management concern in the CRE: A review of information about an assessment of the status of habitat needs and availability, and ecosystem roles of the species.
- ✓ By TWG: At its June 2011 meeting, TWG passed the following motion by consensus. TWG recommends that AMWG consider the report of the Species of Concern Ad Hoc Committee entitled, “Assessment of Taxa of Management Concern in the Colorado River Ecosystem, Glen and Grand Canyons, Arizona, USA: Habitat Needs, Availability, and Ecosystem Roles” dated 15 June 2011, which meets the criteria set forth in the AMWG motion of April 2009, calling for “a review of information about an assessment of the status of habitat needs and availability, and ecosystem roles” of native species of management concern.
- ✓ By an Ad Hoc Group: See description below, under “Background Information.”

---

---

Relevant Science

See attached report.

---

---

Background Information

The attached report, *Assessment of Taxa of Management Concern in the Colorado River Ecosystem, Glen and Grand Canyons, Arizona, USA: Habitat Needs, Availability, and Ecosystem Roles*, was developed by the Species of Concern Ad Hoc Committee in response to the motion passed at the April 2009 AMWG meeting, as documented above. Changes were made in response to comments from TWG.

In this document, the Committee provides previously uncompiled background information on the status of plant and animal taxa of management concern (TMC) in the CRE, identifies previously unrecognized dam-related influences on CRE biological diversity, and summarizes TMC restoration prioritization approaches. TMC were defined primarily as native CRE species that were:

- presently or recently federally listed or were recent candidate species;
- federally or state protected;
- believed by the Committee to be declining, extirpated, or extinct; or
- rare endemic taxa or highly isolated populations, as well as those rare taxa for which too few data are available to assess present status (Appendix A).

TMC were selected by the Species of Concern Ad Hoc Committee (the Committee) based on peer-reviewed published and agency literature, and recommendations from staff and independent experts. The Committee met six times, mostly through teleconferencing, and held numerous informal discussions. We reviewed and documented the background literature, including the conservation status of each taxon.

At least 14, and perhaps more than 23, animal taxa that existed or regularly bred in the CRE have been extirpated, with only two of those extirpations occurring prior to the closure of Glen Canyon Dam in 1963. A total of 18 taxa (21 percent of the TMC list) are federally or state-protected or are of specific importance to Tribes, either through the ESA, through separate federal legislation, or through state or Tribal recognition. Among these 18 taxa are at least 10 extant endangered species. Extirpated or at-risk species that are not federally listed make up the majority of the TMC in the CRE (including all of the above taxa). A total of 46 (53.5 percent) of all TMC had restoration potential scores  $\geq 50$ , suggesting that they may warrant management attention, including research, monitoring, or restoration. Although the focus of this report is on native TMC, we note the presence of several hundred non-native taxa (particularly plants, fish, several invertebrates, and several birds), which interact with native species in complex fashions and play important ecological roles in the CRE. A more comprehensive ecosystem-based understanding of the role of non-native species is warranted, but is beyond the scope of this report.

To better understand restoration prioritization, we investigated five approaches by considering:

- 1) federal endangered status under the Endangered Species Act (1973),
- 2) NatureServe's regional, national, and global conservation status,
- 3) multi-agency policy status,
- 4) a detailed ranking of restoration issues, and
- 5) trophic status and ecological roles in relation to habitat requirements.

Lastly, the Committee reviewed restoration prioritization by undertaking a post-process “reality check” to confirm whether information was sufficiently robust to recommend a taxon to the AMWG for further consideration.

There is at present no commitment for the AMP to fund TMC restoration efforts, and such activities may be the specific purview of the resource managing agencies. Restoration of TMC must be consistent with NPS and other managing agencies’ reintroduction policies, but heretofore has been regarded as tertiary to improving the population health of one federally listed fish species (humpback chub) and sediment management. Nonetheless, the AMWG can play an important multi-agency advisory role to the Secretary in evaluating TMC restoration priorities and potential in the CRE. In addition, the AMP may benefit from monitoring information about reintroduced or translocated TMC, as well as in public outreach. We offer this report as information to the AMWG, recognizing that the diverse views among AMWG stakeholders as to the AMP’s responsibilities require further discussion about roles and responsibilities for CRE management.

**ASSESSMENT OF TAXA OF MANAGEMENT CONCERN IN THE COLORADO RIVER ECOSYSTEM, GLEN AND GRAND CANYONS, ARIZONA, USA: HABITAT NEEDS, AVAILABILITY AND ECOSYSTEM ROLES  
DRAFT FINAL REPORT 15 JUNE 2011**

**EXECUTIVE SUMMARY**

**Introduction**

The loss of native species is one of the three largest human impacts on natural ecosystems, next to habitat destruction and the introduction of non-native species (Groom et al. 2006; Kalyani 2007). Removal of ecologically important foundation taxa (e.g., dominant trees, abundant prey taxa) or keystone species (e.g., species that influence trophic structure and composition), and the introduction of non-native taxa into those roles in altered habitats, affect the structure, composition, function, and resilience, as well as the goods and services provided by ecosystems. Understanding the distribution and ecological roles of native species that are declining or no longer present in altered environments can provide long-term guidance on the stewardship of human-dominated ecosystems, such as large, regulated rivers.

Here we describe the status of species known or suspected to be missing or in decline in the Colorado River ecosystem (CRE) in lower Glen Canyon and throughout Grand Canyon on the southern Colorado Plateau (Fig. 1). We provide approaches to understanding restoration potential to the Adaptive Management Work Group (AMWG) in response to its April 2009 motion:

“In recognition of GCDAMP [Glen Canyon Dam Adaptive Management Program] goals toward management of the Colorado River through an ecosystem approach, AMWG directs the TWG [Technical Work Group] to establish the Species of Concern Ad Hoc Committee [SMCAHC, (the Committee)] and requests the participation of GCMRC [U.S. Geological Survey Glen Canyon Monitoring and Research Center] in that ad hoc committee, to produce a draft report to be presented to AMWG on or before by May 1, 2011, that contains the following with regard to species of management concern in the CRE: A review of information about an assessment of the status of habitat needs and availability, and ecosystem roles of the species.”

The focus of this report is on the native plant and animal taxa of management concern (TMC) in the CRE, and assessment of their restoration potential. TMC were defined primarily as native CRE species that were: presently or recently federally listed or were recent candidate species; federally or state protected; believed by the Committee to be declining, extirpated, or extinct; or rare endemic taxa or highly isolated populations, as well as those rare taxa for which too few data are available to assess present status (Appendix A). TMC were selected by the Goal 3 Ad Hoc Committee (the Committee) based on peer-reviewed published and agency literature, and recommendations from staff and independent experts.

**Methods**

The Committee developed the TMC list from diverse information sources, including published and unpublished literature, and we conducted interviews and discussions with experts

and Committee members in 6 formal and numerous informal meetings. We reviewed and documented the background literature, including the conservation status of each taxon.

To better understand restoration prioritization we investigated five approaches by considering: 1) federal endangered status under the Endangered Species Act (1973), 2) NatureServe's regional, national, and global conservation status, 3) multi-agency policy status, 4) a detailed ranking of restoration issues, and 5) trophic status and ecological roles in relation to habitat requirements. The status of (1) and (2) were determined from the literature. In process (3) we qualitatively evaluated policy issues for a given taxon among the AMP participating agencies, including: a) extirpation status; b) federal listing status or prior inclusion in conservation plans; and c) taxa of interest because: i) distribution and status information was insufficient; ii) species of individual agency concern, including Tribally important taxa; iii) taxa occurred incidentally or were rare; iv) taxa were known or suspected to be affected by dam operations, including populations that may be increasing; v) species were suspected to be declining directly or indirectly due to dam operations; and vi) the special case of federally protected non-native taxa (feral burro; *Equus asinus*).

Process (4) involved consideration by the Committee of a suite of 18 issues that influence management decisions about population restoration (Table 1). These questions involved the role of the dam in the population decline, conceptual and practical restoration issues (e.g., stock availability), the ease with which restoration could be accomplished, the conservation significance of the action, and the level of stakeholder agreement for restoration. Scoring was performed by Committee members, expert staff from TWG participating agencies, and independent academic experts, each of whom scored each issue for each taxon with which they were familiar. We used a 0 (no value, low restoration potential) to 4 (high importance, high restoration potential) scoring scale. A total score was calculated for each species based on the total percent possible of non-blank, unweighted criteria scored by independent experts and knowledgeable Committee members. This process involved considerable discussion within the Committee. A total score of 50 percent or higher was considered to indicate potential conservation concern.

Process (5) was evaluated by examining the trophic position of each TMC in relation to a generalized ecosystem model involving habitat use, ecological role, ecosystem energy flow, and degree of ecological interaction with other taxa.

Lastly, the Committee reviewed restoration prioritization by undertaking a post-process "reality check" to confirm whether information was sufficiently robust to recommend a taxon to the AMWG for further consideration. We discuss our results in relation to agency restoration practices and long-term ecosystem management implications of the findings.

## **Results**

***Extirpated Taxa:*** At least 14, to more than 23 animal species that existed or regularly bred in the CRE have been extirpated, with only two of those extirpations occurring prior to the closure of Glen Canyon Dam in 1963 (California condor - *Gymnogyps californianus*, and gray wolf - *Canis lupus youngi*; Table 2; Appendix A). TMC known to have been extirpated include: 4 fish (Colorado pikeminnow – *Pychocheilus lucius*, razorback sucker – *Xyrauchen texanus*, and bonytail and roundtail chub – *Gila elegans* and *G. robusta*, respectively), 2 amphibians (*Lithobates pipiens* and *L. nr. yavapaiensis*), zebra-tailed lizard, 4 birds (breeding southwestern willow flycatcher – *Empidonax trailii extimus*; western yellow-billed cuckoo - *Coccyzus americanus occidentalis*, Gambel's quail - *Callipepla gambelii*, and California condor, which

was reintroduced outside the CRE in 1996); and 3 mammals (badger – *Taxodea taxa*; the likely extinct southwestern river otter – *Lutra canadensis sonora*; and gray wolf, which was extirpated from northern Arizona prior to 1946). The status of several apparently rare herpetofauna (e.g., regal ring-necked snake – *Diadophis punctatus regalis*, western threadsnake – *Leptotyphlops humilis*, other snakes), as well as numerous rare plant and invertebrate taxa, remains unknown.

***Endangered or Previously Listed TMC:*** A total of 18 taxa (21 percent of the TMC) are presently, or recently have been federally or state-protected, or are of specific importance to Tribes, either through the ESA, through separate federal legislation, or through state or Tribal recognition. Among these 18 taxa are at least 10 extant endangered species exist in the CRE, including: 1 terrestrial gastropod, 1 fish species, 4 bird species, and several incidentally-occurring listed bird taxa (Appendix A).

***Non-listed Extirpated, Declining, and At-risk TMC:*** Extirpated or at-risk species that are not federally listed make up the majority of the TMC in the CRE. These species include all of the above taxa. A total of 46 (53.5 percent) of all TMC had restoration potential scores  $\geq 50$ , suggesting that they may warrant management attention, including research, monitoring, or restoration. For example, Goodding's willow (*Salix gooddingii*) has disappeared from numerous localities at which it occurred at and before 1963, with loss related to beaver foraging and post-dam recruitment failure driven by coarsened grain size (Stevens 1989, Grand Canyon Wildlands Council, Inc. 2009). However, the quality of status and distribution data on at least 52 (61 percent) of the TMC was inadequate to determine management needs or options for many rare taxa, including plant, invertebrate, herpetofaunal (especially amphibians and snakes), mammal, and migratory bird and bat taxa.

***Non-Native Species:*** Although the focus of this report is on native TMC, we note the presence of several hundred non-native taxa (particularly plants, fish, several invertebrates, and several birds), which interact with native species in complex fashions and play important ecological roles in the CRE. A more comprehensive ecosystem-based understanding of the role of non-native species is warranted, but is beyond the scope of this report.

***Dispersal Limitations:*** Glen Canyon Dam may play a long-term role in CRE diversity through interruption of migration and passive propagule distribution. Obviously, Hoover and Glen Canyon dams present barrier to the movement of highly migratory aquatic vertebrates (e.g., several native fish and southwestern river otter); however, prior to 1963, "waif" plant and animal species also were transported into the CRE by via hydrochory or gravity. These plant and perhaps some riparian faunal taxa established, but often were too rare or physiologically stressed to reproduce successfully. Many unlisted but rare taxa identified in this analysis were downstream-dispersed waif species, known only from a few collections in the post-dam CRE. Upstream colonization via animal or wind transport is generally slower than downstream or gravity-based waif recruitment, but both dispersal processes are likely important to long-term biological diversity in the CRE (Stevens and Waring 1987; Shafroth et al. 2002, Stevens and Polhemus 2008; Stevens and Bailowitz 2009).

***Trophic Status of TMC:*** We analyzed the trophic status of extirpated and at-risk TMC among the array of narrow and wide reaches of the CRE (Schmidt and Graf 1990). The present

monitoring focus on 3 TMC (i.e., humpback chub – *Gila cypha*, Kanab ambersnail – *Oxyloma haydeni kanabensis*, and southwestern willow flycatcher) occupy lower and middle trophic positions. While those focal species are administratively important, they may not provide adequate “umbrella” protection for most other CRE TMC and important ecological processes. Therefore, the usefulness of these 3 endangered taxa for understanding and managing the full array of ecosystem dynamics may be limited (Stevens et al. 2001). Three groups of terrestrial CRE species in particular have not received sufficient monitoring or conceptual attention: rare and declining riparian cornerstone plant taxa (e.g., Goodding’s willow); mid-trophic and top predators; and rare or endemic taxa, including insects. Additional monitoring data are needed on such species to develop a more thoroughly linked aquatic-terrestrial conceptual ecosystem model for the CRE, and such an effort is likely to better serve long-term CRE management and the AMP.

### **Comparative Evaluation of Restoration Potential**

The approaches to restoration prioritization we investigated progressively encompass additional CRE taxa, habitats, ecological processes, and trophic strata, the understanding of which may contribute to enhanced ecosystem stewardship. The AMP is presently focused on endangered HBC and its habitat needs, with minor attention on KAS and SWWF. Emphasis on HBC focuses attention on the aquatic domain, primarily in lower Marble Canyon and the lower Little Colorado River. Consideration of KAS directs management attention to the wetland habitat of a single riverside spring in upper Marble Canyon, and expanding focus to SWWF brings management attention to specific locations on riparian terraces in Marble Canyon and upper Lake Mead. Thus, the conservation umbrella provided by these 3 listed species focuses management attention on Marble Canyon and upper Lake Mead, and on primary and secondary consumer trophic levels (Fig. 2). Addition of extirpated species to the stewardship list, such as the missing native fish, expands management attention to the full river corridor and lower reaches of tributaries; addition of amphibians and muskrats adds emphasis to wetland habitats; and addition of extirpated carnivores (e.g., river otter, badger, spotted skunk) and scavengers (e.g., California condor) adds attention to the top predator and decomposer trophic levels, as well as to aquatic-riparian linkage throughout the river corridor. Consideration of non-endangered TMC provides an even more robust stewardship framework. For example, the addition of Goodding’s willow increases management focus on cornerstone riparian primary producers and important herbivores (i.e., beaver, *Castor canadensis*), while focus on endemic taxa covers numerous niches and microhabitats that are not well-conserved by management of the above species.

### **TMC Restoration**

Restoration of TMC must be consistent with NPS and other managing agencies reintroduction policies, but heretofore has been regarded as a tertiary priority, behind improving the population health of one federally listed fish species (humpback chub), and sediment management. Thus, questions of whether, how and when to restore other at-risk or extirpated endangered or non-listed species remain unaddressed. Major information gaps exist with regards to understanding the distribution and present status of many TMC in the CRE. Such information and discussion may play an important role in the planning and implementation of population reintroduction and restoration actions. We describe and discuss issues associated with reintroduction of extirpated CRE TMC, including: a) development of the administrative context;

b) compiling relevant information, stock assessment, and prioritization, c) plan development, d) multi-agency compliance, e) implementation, potentially with multiple stocking efforts, and f) integration of monitoring and feedback information.

## **Discussion**

Effective ecosystem stewardship requires adequate understanding of the ecological roles and potential for restoration of extant as well as extirpated TMC and the processes to which they contribute. Understanding the feasibility and advisability of restoration of extirpated species is coordinated through Goal 3 of the AMWG mission, and also by the wildlife managing agency stakeholders, most notably the NPS, FWS, and AGFD, and several Native American Tribes. Filling information gaps about missing or declining species is needed to more fully inform the Secretary of the Interior about Glen Canyon Dam impacts, as well as the compliance requirements for reintroduction efforts. Through its multi-stakeholder structure, AMWG is uniquely positioned to advise the Secretary of the Interior on the technical, policy, and monitoring information needs and requirements through which to achieve the purposes of AMWG's Goal 3. There is at present no commitment for the AMP to fund TMC restoration efforts, and such activities may be the specific purview of the resource managing agencies. Nonetheless, the AMWG can play an important multi-agency advisory role to the Secretary in evaluating TMC restoration priorities and potential in the CRE. In addition, the AMP may benefit from monitoring information about reintroduced or translocated TMC, as well as in public outreach. In this document we provide previously uncompiled background information on missing and at-risk species, identify previously unrecognized dam-related influences on CRE biological diversity, and summarize TMC restoration prioritization approaches. We offer this information as guidance to the AMWG, recognizing that the diverse views among AMWG stakeholders as to the AMP's responsibilities require further discussion about roles and responsibilities for CRE management.

AMWG monitoring currently focuses on HBC, and habitat issues related to flow, sediment transport, and camping beaches, and minor attention is paid to endangered KAS and SWWF. However, the Committee found at least 5 species (1 plant, 1 lizard, and 3 non-listed fish) that had total ranked scores higher than that of HBC. Concern for Goodding's willow arose due to its CRE-wide demise in the post-dam river corridor. Concern for the 3 non-listed CRE fish arose from the desire of the Committee to prevent those species from becoming federally endangered. The zebra-tailed lizard scored above HBC because its population restoration would be inexpensive and easily accomplished.

In addition, many TMC were recognized that may warrant stewardship actions, or at least periodic monitoring, either by the AMWG and/or by other managing agencies. A total of 47 (55 percent) of the 86 TMC considered here had potential restoration scores  $\geq 50$  percent (Table 2; Appendix A); however, information on distribution, habitat use, and population status was not available for many of those species, as well as species with scores  $< 50$  percent. Opportunities exist to improve the quality of basic information on TMC, and periodic evaluation and assessment of these rare species may be useful to AMP deliberations on ecosystem management. We hope the information provided here contributes to those discussions and to improved stewardship of the CRE.

## INTRODUCTION

### Impacts of Native Species Loss on Ecosystems

The loss of native species is one of the three primary human impacts on natural ecosystems, which also include environmental change and introduction of non-native taxa (Groom et al. 2006; Kalyani 2007). Removal of ecologically important foundation taxa (e.g., dominant trees, abundant prey taxa) or keystone species (e.g., taxa that strongly influence trophic structure and composition), and the substitution of non-native taxa into those roles in altered habitats affect the structure, composition, function, and resilience, as well as the goods and services provided by those ecosystems. Understanding the distribution and ecological roles of native species that are declining or no longer present in regulated rivers, such as the Colorado River ecosystem (CRE) in lower Glen Canyon and throughout Grand Canyon on the southern Colorado Plateau, is fraught with uncertainties because: 1) information on the natural condition of those populations and the ecosystem in general often is limited, and 2) novel arrays of environmental gradients and trophic linkages arise in anthropogenically altered ecosystems, affecting native species population dynamics, as well as ecosystem processes and functions. Accounting for missing or declining species and the ecological processes that characterized the natural ecosystem is fundamental to adaptive management (Walters et al. 2000). Failure to account for missing species, coupled with continued species loss, means that iterative planning and stewardship efforts result in stepwise degradation of ecosystem integrity. Understanding the extent of species loss and restoration




Figure 1: Map of the Colorado River ecosystem between Glen Canyon Dam and Lake Mead, Arizona. Turbidity segments: VT – variably turbid, UT – usually turbid.

potential allows ecosystem stewards to define and manage for desired future ecosystem conditions (Schmidt et al. 1998, Myers et al. 2007).

### **Species Loss in the Grand Canyon Region**

As has occurred in many of the nation's great landscape parks, a substantial number of native taxa have been extirpated or may be at risk in the CRE (Minckley 1991, Newmark 1995, Stevens et al. 2001, Stevens and Gold 2003). Much ecological research has been devoted to the study of reduced or dwindling populations of currently endangered species in the CRE, particularly the big river fishes, and specifically humpback chub (HBC; *Gila cypha*; Minckley 1991; Valdez and Ryel, 1995, 1997; Marsh and Douglas 1997; Stone and Gorman 1999; Gloss et al. 2005; Coggins 2008; Coggins and Walters 2009; Andersen et al. 2010; Yard et al. 2011), as well as one landsnail (Kanab ambersnail – KAS, *Oxyloma haydeni kanabensis*; Spamer and Bogan 1994; Meretsky et al. 2000) and one bird species (southwestern willow flycatcher – SWFL, *Empidonax trailii extimus*; summarized in Paxton et al. 2007). Federal listing prompted ecological research on other formerly listed CRE taxa, including bald eagle (*Haliaeetus leucocephalus*; Brown et al. 1989, 1998; Brown and Stevens 1997) and peregrine falcon (*Falco peregrinus anatum*; Brown et al. 1992; Stevens et al. 2009). However, the ecological roles and distributions of other endangered species that occur incidentally in or near the river corridor (e.g., California condor - *Gymnogyps californicus*; Mexican spotted owl - *Strix occidentalis lucida*), as well as non-listed endemic, declining, or extirpated species, is less well known.

Umbrella species are those that, when conserved, provide protection to co-occurring taxa, habitats, or ecosystems. Federally endangered (“listed”) taxa often are assumed to serve an umbrella function; however, meta-analyses of this conservation concept indicate that focal species selection requires careful assessment of the ecological roles of the assemblage of concern (Branton and Richardson 2010). Importantly, they report that listed taxa may not necessarily serve as adequate umbrella taxa (Roberge and Angelstam 2004), particularly in complex landscapes and human-dominated ecosystems. For example, Ozaki et al. (2006) reported that northern goshawk (*Accipiter gentilis*), a wide-ranging predator that facultatively adapts to anthropogenically altered landscapes, did not provide adequate protection for associated species and habitats.

Although HBC is used as a focal species for CRE management, but adults are relatively low-vagility fish that and almost exclusively spawn in the lower Little Colorado River, a spring-fed tributary in Grand Canyon. Therefore, HBC may not well represent the ecosystem requirements of more highly migratory fish species, such as razorback sucker (RBS; *Xyrauchen texanus*) or Colorado pikeminnow (CPM; *Pychocheilus lucius*), both of which have been extirpated, or terrestrial species (e.g., KAS or SWFL; Stevens et al. 2001). Interestingly, most listed taxa in the CRE display adaptations to the altered riverine environment: HBC, Kanab ambersnail, bald eagle, peregrine falcon, and southwestern willow flycatcher all preferentially use non-native or highly altered food resources (e.g., Valdez and Ryel 1995, Stevens et al. 2001, Brown et al. 1998, Stevens et al. 2009).

A broader examination of the status of taxa of management concern (TMC) in the CRE can provide insight into improved stream-riparian ecosystem restoration options, consideration by the Glen Canyon Dam Adaptive Management Program (AMP), and provide support to the National Park Service (NPS) and other resource managers working to fulfill responsibilities, such as that of the NPS to “... strive to understand, maintain, restore, and protect the inherent

integrity of the natural resources, processes, systems, and values of parks ... (U.S. National Park Service 2006:36).

### **Ecosystem Roles of Extirpated or Declining Taxa**

Extirpated and declining taxa, federally listed or not, may have had important ecosystem roles in the past, and the loss of those species may greatly compromise ecosystem function (Soulé et al. 2005; Myers et al. 2007). Because of the important role of keystone (ecologically strongly interacting species) on ecosystem function, and because the legal framework may not have kept up with advances in ecosystem and conservation ecology, Soulé et al. (2005) recommend that natural resource managers... "adhere to a doctrine of 'best conservation practices based on the best science,' applying a more rigorous standard for the management of relatively interactive species than may be mandated by older statutes and effected by current practice and convention." In addition, little scientific attention has been paid to the distribution and conservation of rare and endemic taxa, in Grand Canyon and elsewhere, particularly invertebrates, for which few population status or life history data exist. The continued health of those populations may be jeopardized by uninformed or insensitive land management practices. Stevens and Polhemus (2008) reported that 53% of 89 aquatic Heteroptera species in the Grand Canyon region were found at 3 or fewer sites (primarily springs), and that more than 25% were found only at single localities. Springs in the vicinity of Grand Canyon are widely recognized as being threatened by regional groundwater pumping. External threats to the regional aquifers around Grand Canyon may affect discharge of springs and therefore the habitats of many springs-requiring species. For example, groundwater pumping may threaten the flow of Blue Springs in the lower Little Colorado River, a springs complex that provides essential habitat for spawning humpback chub. Therefore, management of both species and habitats in a regional context may be needed to maintain and restore CRE species and ecosystem health.

### **Species Translocations and Reintroductions in the Grand Canyon Region**

The establishment and translocation of native and non-native species has a lengthy history in the CRE. Federal management of the North Rim early in the 20<sup>th</sup> Century involved widespread elimination of upland predators, actions that may have had large impacts on ecosystem structure (Rasmussen 1941). Exotic plants in the CRE include tamarisk (*Tamarix* spp.), Russian olive (*Elaeagnus angustifolia*), and more than 75 other species (Stevens and Ayers 2002). Numerous aquatic invertebrate taxa were released by the AGFD into the tailwaters of Glen Canyon Dam in the mid-late 1960's to improve the aquatic foodbase (Stone and Queenan, 1967; Stone and Rathbun 1968, 1969). Endangered Kanab ambersnail were successfully introduced into upper Royal Arch Creek in Grand Canyon in 1998 by AGFD and the NPS to develop a refuge population if dam operations negatively affect natural snail habitat at Vaseys Paradise Spring in the CRE (Spamer and Bogan 1994; Sorensen 2005). Introduced fish include common carp (*Cyprinus carpio*) and catfish (Ictaluridae), which already were the most abundant fish in the lower Colorado River by the beginning of the Twentieth Century (Grinnell 1914). Rainbow (*Oncorhynchus mykiss*) and brown trout (*Salmo trutta*) were released by the NPS shortly after the establishment of Grand Canyon National Park, and 17 other non-native fish species also have become established. Rainbow trout became an important recreational resource in Glen Canyon following closure of the dam in 1963, but trout piscivory may threaten native fish (Yard et al. 2011). The NPS, FWS, and AGFD have translocated HBC into previously unoccupied habitat in the Little Colorado River, upstream from a barrier falls (U.S. Fish and

Wildlife Service 2010), and into Shinumo Creek. Non-native burros (*Equus asinus*) were widely introduced by early prospectors, but were removed from Grand Canyon National Park and most of the CRE in the early 1980s (a small population still exists on the Hualapai Indian Reservation in western Grand Canyon). Semi-native terrestrial vertebrate introductions include: wild turkey (*Meleagris gallopavo*; Brown et al. 1987), and bighorn sheep (*Ovis canadensis nelsoni*) in upper Grand Canyon. Restoration of endangered California condor (*Gymnogyps californianus*) since 1995 generally has been successful (Southwest Condor Release Team 2007); but Meretsky et al. (2000a) question whether reintroduction can be successful because of condor susceptibility to lead poisoning. That issue has prompted proposal of a regional ban on lead shot. Grand Canyon supports the largest breeding population of previously federally listed peregrine falcons (*Falco peregrinus anatum*) of any land management area in the lower 48 states as a result of pesticide control and reintroduction efforts elsewhere in the Colorado River basin in the early 1980's (Brown et al. 1992; Stevens et al. 2009). Louisiana river otters (*Lontra canadensis lataxina*) were translocated into the upper Colorado River and Verde River drainage in Arizona, restoring their contribution to ecosystem function (GCWC 2005). Clearly, much of the CRE has been affected by species introductions and translocations.

### **Adaptive Management Goal 3**

The Adaptive Management Work Group (AMWG) approved the following motion at its April 2009 meeting:

*In recognition of GCDAMP [Glen Canyon Dam Adaptive Management Program] goals toward management of the Colorado River through an ecosystem approach, AMWG directs the TWG [Technical Work Group] to establish the Species of Concern Ad Hoc Committee (SMCAHC) and requests the participation of GCMRC [U.S. Geological Survey Glen Canyon Monitoring and Research Center] in that ad hoc committee, to produce a draft report to be presented to AMWG on or before by May 1, 2011, that contains the following with regard to species of management concern in the CRE: A review of information about an assessment of the status of habitat needs and availability, and ecosystem roles of the species.*

This directive requires a more complete review of information on the distribution, status, ecological role(s) than has heretofore been conducted, as well as development and comparison of potential methods to prioritize the reintroduction potential of extirpated and at-risk TMC to better understand and manage the CRE as an ecosystem.

Here we present a list of the native CRE TMC, including extirpated, endangered, declining, or those for which insufficient data exist to determine current population status. We describe the distribution, habitat requirements, ecological role(s), and status of those taxa, as well as the quality of available information. We integrate selected taxa into a conceptual trophic diagram of the CRE, discuss the advantages and challenges associated with developing a more complete understanding of their status, and we present approaches for prioritizing the reintroduction potential. We briefly review the compliance requirements of participating agencies in relation to reintroduction efforts. We provide this information to clarify dam impacts on the CRE and assist the AMWG achieve its mission of adaptive ecosystem management.

## **METHODS**

### **Study Area**

The highly regulated CRE under the purview of the AMWG extends from the forebay of Lake Powell and Glen Canyon Dam in Glen Canyon National Recreation Area (GLCA) and for a total distance of 447 river km downstream through Grand Canyon National Park (GRCA) to the boundary of Lake Mead National Recreation Area at about Colorado River Mile (RM) 278 (Fig. 1). The CRE includes the river and shorelines within the 100-yr flood stage, the lower portions of selected tributaries, and the lower 15 km of the Little Colorado River. The linked aquatic-terrestrial domains of the CRE are structured into 13 geologically defined geomorphic reaches, varying by width and depth, with upstream clearwater and variably turbid reaches sustaining higher aquatic productivity, lower temperature, and coarser grain size than the downstream, usually turbid reaches (Schmidt and Graf 1990; Stevens et al. 1997b). The CRE is structured at a fine scale into a repeated microhabitat array at the mouths of more than 500 tributaries (Schmidt and Graf 1990). Turbidity swamps the ecosystem influences of geomorphology in the aquatic domain, while geomorphology overrides turbidity impacts in the terrestrial domain (Stevens et al. 1995, 1997a,b). River mile (RM) from Lees Ferry is traditionally used to designate location in the CRE, with mileage measured for both upstream (negative) and downstream (positive) distance. The side of the river is designated looking downstream (right=R or left=L).

The CRE is managed under several administrative processes. The National Park Service (NPS) manages natural and cultural resources of 2 NPS unit (Grand Canyon National Park and Glen Canyon National Recreation Area), pursuant to the NPS Organic Act (1916), Grand Canyon National Park enabling legislation (1919), Glen Canyon National Recreation Area enabling legislation (1972), and subsequent legislation including the 1992 Grand Canyon Protection Act, and in accord with: the U.S. Fish and Wildlife Service (FWS) and the Endangered Species Act of 1973 (ESA) and the Fish and Wildlife Coordination Act (FWCA). The Arizona Department of Water Resources and the Arizona Game and Fish Department (AGFD) administer water, fish and wildlife resources management through state policies. Five actively participating Native America Tribes have sovereign nation rights over selected CRE resources and sites. Glen Canyon Dam management is achieved under the 1997 Record of Decision on dam operations through Secretarial advisement by the AMWG -- a Federal Advisory Committee composed of representatives from 27 major stakeholder groups (Stevens and Gold 2003, Melis and Lovich 2005), and by the Colorado River Annual Operating Plan (AOP) group. The Secretary of the Interior is the Water Master of the Colorado River and integrates recommendations from both AMWG and the AOP group. Goal 3 of the Glen Canyon Dam Adaptive Management Program (AMP) is to "Restore populations of extirpated species, as feasible and advisable." The AMWG developed the above-quoted motion to provide relevant information as a comprehensive first step towards achieving Goal 3.

### **Information Sources**

The focus of this document is the array of native plant and animal taxa of management concern (TMC) in the CRE. TMC were defined as native CRE species (but including federally protected non-native feral ass, *Equus asinus*) that were: presently or recently federally listed or were recent candidate species; Arizona state listed; federally protected; believed by the Committee to be declining, extirpated, or extinct; or rare endemic taxa or highly isolated populations, as well as those for which too few data are available to assess present status (Appendix A). TMC were selected by the Goal 3 Ad Hoc Committee (the Committee) based on

peer-review published and government literature, and recommendations from staff and independent experts, as well as the published and unpublished literature, including a regional conservation area plan developed by Grand Canyon Wildlands Council, Inc. (2009).

### **Restoration Scoring and Prioritization**

The Committee developed the TMC list from diverse information sources, and investigated five restoration prioritization approaches (Table 1): 1) endangered status, 2) NatureServe conservation status, 3) multi-agency policy status, 4) a detailed ranking of restoration issues, and 5) consideration of trophic status. Process (3) involved qualitative consideration of policy among participating agencies, including: a) extirpation status; b) TMC that were endangered, declining, rare, or already included in existing conservation plans; c) taxa of interest because: i) distribution and status data are insufficient; ii) species of individual agency concern, including Tribally important taxa; iii) incidentally occurring rare species; iv) species known or suspected to be affected by dam operations, which may be increasing; v) species suspected to be declining directly or indirectly due to dam operations; and vi) federally protected non-native taxa (feral burro; *Equus asinus*).

Process (4) involved consideration of a suite of 18 issues that influence decisions about population restoration (Table 1). These questions involved the role of the dam in the population decline, conceptual and practical restoration issues (e.g., stock availability), the ease with which restoration could be accomplished, the conservation significance of the action, and the level of stakeholder agreement for restoration. Scoring was performed by Committee members, expert staff from TWG participating agencies, and independent academic experts, each of whom scored each issue for each taxon with which they were familiar. We used a 0 (no value, low restoration potential) to 4 (high importance, high restoration potential) scoring scale, and variables were not weighted. A total score was calculated for each species based on the total percent possible of non-blank issues scores. Total scores were generated from unweighted criteria, scoring by independent experts and knowledgeable TWG staff, and considerable discussion within the Committee. A total score of 50 percent or higher was considered by the Committee to indicate potential conservation concern.

Process (5) was evaluated by examining the position of each TMC in relation to a generalized ecosystem model involving trophic function, ecosystem energy flow, and degree of ecological interaction.

Lastly, the Committee reviewed the process by conducting a post-process “reality check” to confirm whether information was sufficient to recommend a taxon to the AMWG for further consideration. We discuss our results in relation to agency restoration practices and long-term ecosystem management implications of the findings.

Table 1: Issues related to restoration potential of species of management concern in the CRE. Each issue was scored on a 0 to 4 scale by each expert within a taxon, where information was deemed sufficient. The total score was the percent of possible scores excluding missing or blank values. Asterisk (\*) denotes an issue not included in analyses.

<b>Potential Restoration Issues</b>
Influence of Glen Canyon Dam on population
Conservation relevance
Quality of background information
Stock availability for restoration
Legal mandates, requirements
Habitat availability under present dam operations
Habitat availability under realistic future dam operations
Technical feasibility
* Agreement among stakeholders
Inexpensiveness (low cost=high score)
Time-frame for success (short=high, long=low)
Logistical ease
Ease of compliance
Are genetics issues easily resolved?
Lack of disease transmission potential
Urgency in relation to climate change impacts
Beneficial interactions with other species

**RESULTS**

**Overview**

We assessed the status and restoration potential of a list of 86 TMC from the CRE (Appendix A; Table 2). At least 14 (and perhaps as many as 23 TMC have been extirpated from the CRE. total of 18 taxa (21 percent) are presently or recently have been federally or state-protected. Four federally endangered fish species have been extirpated, as have 2 candidate leopard frog species, and 3 bird species. Formerly listed peregrine falcon and bald eagle are extant. All presently or formerly ESA-listed and candidate species also are recognized as Species of Greatest Conservation Need in Arizona’s State Wildlife Action Plan (SWAP; AGFD 2006). Several TMC not federally listed have been extirpated, including one lizard and the likely extinct southwestern river otter, with loss of river ecosystem function (GCWC 2005). Only 2 species (HBC, KAS; 2.5 percent of the total TMC) are regularly monitored by the AMWG program, while the NPS monitors SWFL.

At least 46 (53 percent) of CRE TMC had restoration potential scores  $\geq 50$ , suggesting that they may warrant management attention or at least monitoring. All native migratory bird species are federally protected under the Lacey Act of 1900 and the Migratory Bird Protection Act of 1918, and their habitats whenever flows in the Colorado River are modified due to operations of Glen Canyon Dam under the Fish and Wildlife Coordination Act (FWCA) of 1934, as amended. Additionally, the habitat requirements of fish species that are not listed and protected by the Endangered Species Act of 1973, as amended, are similarly protected under the FWCA. However, the quality of status and distribution data on at least 52 (60 percent) of the TMC was inadequate to determine management needs or options, including rare invertebrate,

herpetofaunal (especially amphibians and snakes), rodent, and migratory bird and bat taxa. In the following section, we describe extirpated, federally or state listed or former candidate taxa, and rare, endemic and at-risk TMC.

## **SPECIES ACCOUNTS**

### **Extirpated Taxa**

**Overview:** At least 14, and as many as 23 animal species that formerly regularly bred in the CRE have been extirpated, with only two of those prior extirpations occurring prior to the closure of Glen Canyon Dam in 1963 (California condor - *Gymnogyps californianus*, and gray wolf - *Canis lupus youngi*; Table 2; Appendix A). TMC known to have been extirpated include: 4 fish (Colorado pikeminnow – *Ptychocheilus lucius*, razorback sucker – *Xyrauchen texanus*, and bonytail and roundtail chub – *Gila elegans* and *G. robusta*, respectively), 2 amphibians (*Lithobates pipiens* and *L. nr. yavapaiensis*), zebra-tailed lizard, 4 birds (breeding southwestern willow flycatcher – *Empidonax trailii extimus*; western yellow-billed cuckoo - *Coccyzus americanus occidentalis*, Gambel’s quail - *Callipepla gambelii*, and California condor, which was reintroduced outside the CRE in 1996); and 3 mammals (badger – *Taxodea taxa*; the likely extinct southwestern river otter – *Lutra canadensis sonora*; and gray wolf, which was extirpated from northern Arizona prior to 1946). The status of several apparently rare herpetofauna (e.g., regal ring-necked snake – *Diadophis punctatus regalis*, western threadsnake – *Leptotyphlops humilis*, other snakes), as well as numerous rare plant and invertebrate taxa, remains unknown. Below we describe these extirpated taxa.

**Razorback Sucker (RBS; *Xyrauchen texanus*):** RBS were previously widely distributed and highly migratory in the Colorado River, but have been extirpated from the CRE (Dowling et al. 1996, Mueller et al. 2000). RBS presently are being raised and restocked into lower Colorado River reservoirs, including golf course ponds. RBS restoration downstream from Glen Canyon Dam may require continued stocking if recruitment fails and non-native predator control efforts are not successful. Discussion of RBS reintroduction into Grand Canyon has been on-going since 1990. RBS in spawning condition and early stage larvae have recently been collected near the CRE inflow to Lake Mead, and the species is recruiting successfully in other parts of Lake Mead (<http://www.usbr.gov/uc/feature/Razorback/index.html>, accessed February 7, 2011).

**Colorado Pikeminnow (CPM; *Ptychocheilus lucius*):** CPM were the largest native fish in the pre-dam Colorado River. Called “white salmon” or “Colorado River salmon” by early settlers, this species apparently undertook long annual migrations through much of the Colorado River basin (Sigler and Sigler 1996). Breeding populations are now restricted to the upper Colorado River basin. CPM is a large, piscivorous species, and restoration of CPM in the CRE likely will require analysis of reintroduction success in this regulated reach, as well as potential predation impacts on HBC.

**Bonytail and Roundtail Chub (BTC, RTC; *Gila elegans* and *G. robusta*, respectively):** BTC were previously known from near Glen Canyon Dam and Lake Mead (Minckley 1973), but at this time there are no known wild populations of *Gila* in the mainstream lower Colorado River downstream from Grand Canyon. Augmented populations are sustained by stocking in both Lake Havasu and Lake Mohave. The pre-dam distribution of RTC is not well documented, but is reported as primarily occurring in tributaries of the Colorado basin, not in the mainstem

Table 2: Summary of prioritization approaches for taxa of management concern (TMC) in the Colorado River ecosystem downstream from Glen Canyon Dam. Cells totals do not sum because some taxa qualify in multiple categories. Appendix A contains detailed accounts of the scoring of each TMC.

Criterion	Number of CRE TMC									
	Plants	Trematode	Insects	Snails	Fish	Amphibians	Reptiles	Birds	Mammals	Total
Extirpated	2?	0	2?	0	4	2	2	3 (4)	8	14- 23
Federally ES or recent ES	0	0	0	1	5	2	0	9	1	18
Composite Score $\geq 50$	0	1	3	1	8	3	3	9	19	47
3a. TMC about which little is known	1	0	9	0	3	2	5	6	26	52
3b. TMC of individual agency concern	6	0	3	1	8	4	3	16	21	62
3c. Rare or endemic TMC	1	1	3	1	0	1	0	5	2	14
3d-e. TMC affected by dam, (+ increasing or - decreasing)	6	1	4	1	8	4	4	13	11	52
Reality Check ("yes")	4	0	0	1	6	0	0	2	0	13
Reality Check ("mixed")	3	1	9	0	2	4	6	17	31	73
<b>Total number TMC *</b>	<b>7</b>	<b>1</b>	<b>9</b>	<b>1</b>	<b>8</b>	<b>4</b>	<b>6</b>	<b>19</b>	<b>31</b>	<b>86</b>

Colorado River downstream of Glen Canyon Dam (Holden and Stalnaker, 1975). Douglas and Douglas (2006) analyzed *Gila* genetics, concluding that the Grand Canyon population was distinct, but indicating some intergradation had occurred with other *Gila species*. Therefore, introduction of other *Gila* species into Grand Canyon is contraindicated, at least at present, as it might result in hybridization, threatening the genetic integrity of the Grand Canyon HBC population.

**Lowland Leopard Frog (*Lithobates nr. yavapaiensis*; LLF):** A single population of leopard frog was detected in lower Grand Canyon at Surprise Canyon (Colorado River Mile 248R; Brennan and Holycross 2005). This population has been documented along the Colorado River (L.E. Stevens, unpublished photograph) but not in recent decades, and primarily occurs several km upstream in Surprise Canyon. This leopard frog population is highly isolated and may be a unique form of *L. yavapaiensis* (Oláh-Hemmings et al. 2010).

**Northern Leopard Frog (NLF, *Lithobates pipiens*):** Northern leopard frog was detected at Cardenas Creek (RM 71L; Tompko 1976; Miller et al. 1981) and at -9 Mile Marsh upstream from Lees Ferry (Brennan and Holycross 2005). The Cardenas Marsh backwater has filled with sediment and both historical sites have been heavily colonized by *Phragmites australis*, rendering them unsuitable as NLF habitat. NLF have not been detected at Cardenas Marsh since 1975 or at -9 Mile Marsh since the late 1990s, and this species is considered to have been extirpated from the CRE. A small population also was detected at Hidden Sloughs (-6.5R) in the mid-1990's but was likely introduced there and has not persisted (J. Spence, NPS Biologist, personal communication). Historical records of northern leopard frogs exist from Bright Angel Creek

and lower Kanab Creek, but recent, intensive searches by the U.S. Geological Survey and Northern Arizona University have failed to reveal any extant populations (C. Drost, U.S. Geological Survey biologist, personal communication).

***Southwestern Willow Flycatcher:*** Migrant and breeding SWFL have been documented in the CRE at Lees Ferry and in lower Marble Canyon since the 1930s (Brown et al. 1987). However, breeding SWFL disappeared from their historic range in eastern Grand Canyon, where they regularly nested until the late 1990s (Paxton et al. 2007). Loss of this nesting population has been attributed to external factors, such as winter habitat alteration, as well as to changes in the CRE, such as increased brown-headed cowbird (*Molothrus ater*) brood parasitism (Brown et al. 1992; review in Paxton et al. 2007). Single male SWFL are still occasionally detected singing along the river in upper Grand Canyon, and limited breeding occasionally occurs along the Colorado River in Lake Mead, but such activities are erratic. Although SWFL occupies earlier successional stages of riparian habitat, effective conservation of SWFL could umbrella coverage of some of the more than two dozen other neotropical migrant birds that nest in the CRE riparian zone (Brown et al. 1987; Stevens et al. 2001). Many other neotropical breeding species also are negatively affected by cowbird brood parasitism.

***Western Yellow-billed Cuckoo (YBCU; *Coccyzus americanus occidentalis*):*** Listed as a federally threatened species, YBCU requires gallery riparian forest, and therefore is not well-protected by conservation of endangered SWFL, which requires relatively early seral successional riparian vegetation (Layman and Halterman 1987, Brown et al. 1987). In contrast, YBCU requires more-or-less continuous tall-canopy riparian cottonwood and willow cover. YBCU is known in the CRE from a single report in Marble Canyon by Brown et al. (1987), at RM -14.2 in 1995, and at Lee's Ferry (LaRue et al. 2001), but it has not been detected in the CRE in more a decade. YBCU is identified as a Species of Greatest Conservation Need in Arizona's SWAP (AGFD 2006). Depending on how the NPS manages tamarisk in the wake of the arrival of tamarisk leaf beetles (*Diorhabda* spp.), suitable riparian habitat can be promoted that may support this and many other neotropical migrant birds in the CRE.

***Plains Gray Wolf (*Canis lupus youngi*):*** Wolves were extirpated from the region by 1945 (Rasmussen 1941, Hoffmeister 1986); however, John D. Lee reported seeing wolves on the south (left) bank of the river at Lees Ferry early in his tenure as ferryman there in the early 1960s (Brooks and Cleland 2004). Wolf restoration is being tested to the south of the Grand Canyon region in Arizona; however, the subspecies most often suggested for restoration is the Mexican wolf (*C. l. baileyi*). Hoffmeister (1986) explicitly states that *C. l. baileyi* did not occur on either the south or north rims of Grand Canyon. Introduction of Mexican wolf north of the Colorado River could threaten the genetic integrity of the plains gray wolf farther north, and therefore is not advised. Consideration of wolf population restoration in the CRE was rejected by the Committee, as there is little to no relationship of dam operations on wolf recovery.

### **Extant Endangered and Recently Listed Taxa**

***Overview:*** A total of 18 taxa (21 percent of the TMC) are presently, or recently have been federally or state-protected, or are of specific importance to Tribes, either through the ESA, through separate federal legislation, or through state or Tribal recognition. Among these 18 taxa are at least 10 extant endangered species exist in the CRE, including: 1 terrestrial gastropod, 1 fish species, 4 bird species, and several incidentally-occurring listed bird taxa (Appendix A).

***Kanab Ambersnail:*** KAS occurs naturally in Arizona only at Vaseys Paradise Springs in the Colorado River in Grand Canyon, where it remains the focus of considerable taxonomic, ecological, and management debate (Spamer and Bogan 1993, Stevens et al. 2001). Although this taxon likely will be synonymized with *O. h. haydeni* on the basis of genetic analyses, until those analyses are published in the peer-reviewed scientific literature, KAS remains an endangered species and a highly restricted population, and therefore of concern in CRE management. The AGFD, NPS, FWS, Department of Interior Central Utah Project Completion Act Office, and Upper Colorado River Basin Office of the Bureau of Reclamation partnered beginning in 1996 to translocate populations of KAS into remote, off-river tributaries and springs in Grand Canyon, and the population in Royal Arch Creek has persisted since the 1998 translocation, with protection under the ESA. This effort helped mitigate planned flooding impacts at Vaseys Paradise under a FWS Reasonable and Prudent Measure in 1996 and 1997 to find or create at least one population of KAS not influenced by Glen Canyon Dam (summarized by Sorensen 2005). KAS largely supports the only known population of the trematode, *Leuchochloridium cyanocittae* known in Arizona.

***Humpback Chub:*** HBC has received most of the species management attention in the AMWG program because Grand Canyon supports the largest naturally reproducing population of this endangered minnow species in the vicinity of the Little Colorado River (LCR). Intensive monitoring and non-native trout control programs (Grand Canyon Monitoring and Research Center 2008; Coggins et al. 2011; Yard et al. 2011) provide improved insight into long-term population trends, and *in situ* HBC population augmentation in the Little Colorado River upstream from the previously known range of HBC has been successfully undertaken by FWS, adding to the in-LCR population (Sponholtz and Stone 2004). Translocation also has been conducted by the NPS, FWS, AGFD, and collaborators into Shinumo and Havasu Creeks in central Grand Canyon, tributary populations that may help ensure in-Canyon persistence of the species (GCWC 2008; NPS 2011). The NPS and FWS also have discussed HBC translocation into Bright Angel Creek.

***California Condor (Gynogyps californianus):*** California condors were extirpated from Grand Canyon early in the 20<sup>th</sup> Century. External to the AMWG program and the National Park Service, reintroduction of an experimental, non-essential population of this endangered species into the Grand Canyon region has been successful, and pairs of birds nesting in Grand Canyon caves have successfully fledged young (Snyder and Snyder 2000; Alagona 2004). Occupation and use of the CRE by condors is opportunistic, but external restoration efforts have returned this important top scavenger into the CRE. Management of condor-human interactions has proven challenging because condors are naturally attracted to human activities.

***Mexican Spotted Owl (MSOW); Strix occidentalis lucida:*** MSOW occasionally occur in the CRE, coming down from their normal ranges at the back of nearly every large tributary in central Grand Canyon (REF). MSOW have been observed and photographed at the mouths of tributaries such as Shinumo Creek and National Canyon (LES unpublished data, LaRue et al. 2001). Although apparently incidental in the CRE, the high density of riparian rodents may attract this species; however, MSOW occurrence in the CRE is fairly rare and the population certainly does not depend on riparian or dam-related resources.

**Bald Eagle (BAEA; *Haliaeetus leucocephalus*):** Although delisted since 2007, BAEA are still protected under the 1940 Bald Eagle Protection Act, and therefore continue to be TMC in the CRE. Wintering BAEA colonized Grand Canyon beginning in 1982, and were regularly monitored until the mid-1990s (Brown et al. 1989). Their feeding behavior and distribution are strongly affected by dam operations and human recreational disturbance (Brown and Stevens 1997, Brown et al. 1998). However, CRE monitoring data on CRE bald eagles over the past decade are not known to us.

**Peregrine Falcon (PEFA; *Falco peregrinus*):** Delisted from endangered species status in 1999, PEFA have become a common top predator in the CRE (Brown et al. 1992, Stevens et al. 2009). Grand Canyon PEFA are presently monitored by NPS, but population trends and ecological impacts on the CRE avian assemblage, if any, are unclear.

**Incidentally-Occurring Listed Species:** Several other listed species occur or may occur on an incidental basis in the CRE, including: mountain plover (*Charadrius montanus*), which is proposed for listing as threatened (proposed rule June 29, 2010, FR 75 37353), and the recently delisted brown pelican (BRPE; *Pelicanus occidentalis*). The occurrence of these species is incidental in the CRE (Brown et al. 1987; Stevens et al. 1997a), and CRE management attention is unlikely to influence their populations.

#### **Non-listed Extirpated or At-risk Taxa**

**Overview:** Extirpated or at-risk species that are not federally listed make up the majority of the TMC in the CRE. These species include all of the above taxa. A total of 46 (53.5 percent) of all TMC had restoration potential scores  $\geq 50$ , suggesting that they may warrant management attention, including research or monitoring. However, the quality of status and distribution data on at least 52 (61 percent) of the TMC was inadequate to determine management needs or options, including rare invertebrate, herpetofaunal (especially amphibians and snakes), rodent, and migratory bird and bat taxa.

**Plants:** Although no plant species to our knowledge have been verified as extirpated from the CRE, the status of rare and declining wetland and riparian taxa is of concern to long-term ecological processes. Although extirpated from Grand Canyon, yerba-mansa (*Anemopsis californica*) only occurred in GRCA at Indian Gardens on Garden Creek, not in the CRE. Nonetheless, the diminished presence and importance of Goodding's willow, Fremont cottonwood, coyote willow, honey mesquite, netleaf hackberry, and other plant species in the CRE is of sufficient magnitude that loss of their respective ecological functions may limit associated faunal species and habitats.

The persistence of Goodding's willow trees (*Salix gooddingii*) in the CRE is unlikely because bank-dwelling beavers (*Castor canadensis*) in the mainstream have removed most of the pre-dam trees (Mast and Waring 1997, Mortenson et al. 2008), and no recruitment is occurring. A recent survey by Grand Canyon Wildlands Council, Inc. (2009) found that Goodding's willow has been eliminated from 8 of 17 (47.1%) sites between Lees Ferry and Diamond Creek where the tree occurred prior to the closure of Glen Canyon Dam. At most sites, trees had been felled by beaver and continued beaver attack removed resprouting stems until the trees perished.

Successful recruitment was not observed at any of those 17 sites, or elsewhere along the river during that and other vegetation surveys over the past three decades.

Fremont cottonwood (*Populus fremontii*) was relatively rare in the pre-dam CRE, but a large stand existed at Lees Ferry and the mouth of Bright Angel Creek, and a large individual tree existed near RM 193L in pre-dam time (Clover and Jotter 1944, Turner and Karpiscak 1980). This species shares a similar fate with Goodding's willow and cottonwood recruitment requirements are similar to those of the willow. Although cottonwood seedlings occasionally are detected along the river, establishment fails because, like Goodding's willow, virtually all individuals are attacked by beaver. These Salicaceae are renowned as ecologically important species in southwestern rivers, diversifying canopy and subcanopy structure, and supporting high concentrations of neotropical migrant birds (e.g., Carothers et al. 1974).

Although non-native tamarisk extensively colonized the lower riparian zone of the CRE immediately after dam closure, clonal coyote willow (*Salix exigua*) had become a co-dominant species by 1980 (Stevens 1985, 1989). In recent decades CRE coyote willow, like the other Salicaceae, has declined in response to intensive beaver attack, with every clone undergoing repeated harvest by beaver. Like Fremont cottonwood and Goodding's willow, coyote willow supports more than 100 native herbivore and primary consumer taxa. Insect taxa, such as larval viceroy butterfly (*Limenitis archippus*) are host specific on *Salix exigua* in the Southwest. The disappearance of viceroy butterflies from the CRE since 1982 reflects on-going reduction and disappearance of host plants along the river.

Honey mesquite (*Prosopis glandulosa* var. *torreyana*) is abundant on pre-dam terraces in from Mile 39 to 140 and from Mile 165 to Lake Mead, and makes up a large portion of the relict pre-dam riparian vegetation zone (Carothers et al. 1979, Anderson and Ruffner 1989; Waring et al. 2010). While this species is actively colonizing the lower riparian zone in post-dam time, no recruitment is occurring on on upper riparian terraces, where historically it was the dominant species (Waring et al. 2010). Two long-term issues about this pre-dam mesquite belt are the change in potential fire frequency driven by invasion of non-native *Bromus* grasses, and the potential for gradual replacement of mesquite by facultatively riparian catclaw (Fabaceae: *Acacia greggii*).

Netleaf hackberry (*Celtis laevigata* var. *reticulata*) also does not appear to be recruiting successfully in Grand Canyon. Individual stems of this clonal species in the CRE date back to 1776 (Salzer et al. 1996); however, most existing clones appear to be decadent and few seedlings have been reported in post-dam time. Hackberry is remarkably tightly restricted to the pre-dam 2-year flood return stage (ca. 2,830 m<sup>3</sup>/s) along the Colorado River, but unlike honey mesquite, it does not appear to be colonizing the lower riparian zone. Therefore, hackberry may warrant monitoring and determination of flow and habitat requirements for sustainable management.

Several other non-listed native plants are endemic in the CRE riparian zone, and more than 150 plant taxa are known from fewer than 4 records there, including many first-records for the state of Arizona (e.g., Ayers and Stevens 1994, Ayers et al. 1994). Among these rare CRE plants are *Leersia oryzoides*, *Lycopus americanus*, and *Parthenocissus vitacea* at -9L Spring, *Polygonum amphibium* at Vaseys Paradise (32R), *Erigeron utahensis* at 123L, and *Reverchonia arenaria* at 142L, to mention just a few that have more upstream and more northerly ranges. Common riparian taxa in the upper Colorado River in Utah and Glen Canyon often are rare in Grand Canyon, and the absence of these plant taxa may reflect natural physiological or biogeographic limitations, or susceptibility to river flow and sediment management.

**Invertebrates:** Several hundred of invertebrate taxa are rare but of unknown status in the CRE. Among these, several previously documented or reported insect species now are either extremely rare or may have been extirpated from the CRE, including: viceroy butterfly (*Limenitis archippus*; Garth 1950, L.E. Stevens unpublished data), dark buckeye (*Junonia genoveva nigrosuffusa*; L.E. Stevens unpublished data), and the recently described noctuid moth (*Schinia immaculata* Pogue 2004). Several endemic chironomid midges were collected along the Colorado River in the mid-1970s but subsequently have not been detected (Sublette et al. 1998), and several black fly and other nematoceran Diptera collected in the 1970s have not been found in the river corridor recent decades (Stevens, unpublished data). A single population of the nepomorph waterbug, *Belostoma* nr. *flumineum* (Belostomatidae) exists in Vulcan's Well Spring (179L; Stevens and Polhemus 2008). Niobrara ambersnail (*Oxyloma h. haydeni*) occurs in the CRE only at Leopard Frog Marsh (CR -9 L) Springs, as well as at Indian Gardens on the Tonto Platform in Grand Canyon (Sorensen and Nelson 2003), and are the only documented populations of this genus in Arizona. The Niobrara ambersnail is identified as a Species of Greatest Conservation Need in Arizona's SWAP (AGFD 2006). The only known population of the trematode snail parasite, *Leucochloridium cyanocittae* McIntosh 1932 in Arizona occurs in endangered Kanab ambersnail and in *Catinella* nr. *vermeta* snails at Vaseys Paradise (Sorensen et al., unpublished data).

Numerous endemic or extremely isolated insect populations are found adjacent to the CRE and particularly in the lower reaches of perennial springfed streams in Grand Canyon, but their status in the CRE remains unknown. Such species include masked clubskimmer (Odonata – dragonflies, *Brechmorhoga pertinax*) and Grand Canyon wetsalts tiger beetle (*Cicindela haemorrhagica arizonae*), both of which occur in tributaries segments near the Colorado River from Nankoweap Creek (Mile 52R) downstream to Stone Creek (Mile 132R). The distribution and status of the Little Colorado River hairy-necked tiger beetle (*Cicindela hirsuta coloradula*), which is endemic to the riparian zone of lower Little Colorado River, have not been determined (Stevens and Huber 2004; Pearson et al. 2005). The only known U.S. population of the sallow waterbug, *Micracanthia quadrimaculata* occurs at and below the pre-dam 10 year flood stage at “Upper Bass Cable Springs” near Mile 108R (Stevens and Polhemus 2008). Similarly, endemic or isolated nepomorph water bugs (Hemiptera: Belostomatidae, *Abedus breviceps* in Boucher Creek and Ochteridae, *Ochterus rotundus* at low elevation desert springs in the middle reaches of Grand Canyon), and several chironomid midge taxa described by (Sublette et al. 1998), and the empidid *Wiedemannia* (Sinclair 2007) occur at other springs or springfed desert streams along the Colorado River. A newly described asilid robber fly (*Efferia tapeats* Aubrey et al. in prep.) occurs on CRE sandbars and in adjacent tributary riparian zones in the middle reaches of Grand Canyon. Numerous other rare or undescribed invertebrate species exist in the CRE, but their distribution, status, and the potential impacts of dam operations on them remain unstudied.

**Vertebrates:** Non-listed extirpated, seriously declining, or rare and apparently at-risk vertebrate species in the CRE include many species, of which several ecologically important taxa are described below.

1) The only population of CRE zebra-tailed lizard (ZTL; *Callisaurus draconoides*) was extirpated in 1983 when river-runner trucks conducting take-outs were redirected onto the upper sand dunes at Diamond Creek (Tompko 1976; Miller et al. 1981; Stevens et al. 2001, Brennan

and Holycross 2005). ZTL still occur upstream in the Peach Springs Wash drainage, and a population could easily be reintroduced to the mouth of Diamond Creek.

2) The western threadsnake (*Leptotyphlops humilis*) is known in the CRE from three records in the CRE and lower Little Colorado River (Stevens 1983, Brennan and Holycross 2005, D. Stone, FWS, unpublished data). All records of this species are from sand-dominated habitats in the riparian zone of the Colorado River and the lower Little Colorado River; however, its present status is unknown. Likewise, a single record of regal ring-necked snake (*Diadophis punctatus regalis*) exists from RM 194L by LES in 1978. The large, sandbar grassland in which it was detected was eliminated by mid-1980's high flows. The status of several other rare CRE snake taxa also remains unknown.

3) Muskrat (*Ondatra zibithicus*; Hoffmeister 1986; Stevens 1983; Stevens et al. 2001; NPS files) have been reported several times from the post-dam Colorado River corridor in Grand Canyon, and a skull was reported from Tapeats Creek in the 1990's. Additional evidence of this species' presence was detected after 2000 near Mile 77R. This is presently a rare species in the CRE, but muskrats have been detected recently at Lees Ferry and by GCMRC fisheries biologists as occasionally occurring in the lower Little Colorado River (Stone et al. 2008, 2010).

4) Southwestern river otter (SRO, *Lontra canadensis sonora*; Hoffmeister 1986; GCWC 2005; NPS files) is likely extinct in Grand Canyon. Although several reports of otter in the CRE exist from prior to 2000, repeated searches over 30 years have failed to document live individuals. If a resident population of SRO still exists, the introduction of extra-regional otter subspecies in upper Colorado River drainages and the Verde River is likely to eliminate the pure otter genotype through hybridization (Stevens et al. 2001). GCWC (2005) conducted an analysis of the potential for otter reintroduction into Grand Canyon to restore the missing ecological role of this top mammalian fish predator in the CRE; however, interagency concerns over potential impacts of otter on HBC have been raised and need to be resolved for restoration of the otter's ecological role (GCWC 2005)..

5) Bank-dwelling American beaver (Castoridae: *Castor canadensis*) play an important role consuming and disturbing the CRE riparian zone vegetation (Mortenson et al. 2008). Beaver strongly affect establishment and survivorship of Salicaceae (e.g., Goodding's willow and Fremont cottonwood) and other plant species along the river. Improved understanding of long-term beaver habitat availability, population dynamics, and distribution require additional research and more consistent monitoring.

6) Badger (*Taxidea taxa*) were last reported in the CRE in mid-winter 1990-1991 by National Park Service Biological Technician N. Kline at Hidden Sloughs (Colorado River Mile -6.5R; Stevens unpublished data). Jacob Hamblin reportedly shot a badger near the mouth of Badger Creek (Colorado River Mile 8R) in the 1850's (Wixom 1997). Similarly, the distribution and status of other carnivore populations is poorly known and not regularly monitored along the river, including: Procyonidae - ringtail (*Basariscus astutus*) and raccoon (*Procyon lotor*); Canidae - coyote (*Canis latrans*) and gray fox (*Urocyon cinereoargaeus*); Mustellidae - spotted skunk (*Spilogale gracilis*), striped skunk (*Mephitis mephitis*), and hog-nosed skunk (*Conepatus*

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*

*leuconotus*); Felidae - bobcat (*Lynx rufus*) and mountain lion (*Puma concolor*); and Ursidae - black bear (*Ursus americanus*).

7) The nativity of painted turtle (*Chrysemys picta*) is uncertain in Arizona, but there are multiple observations of this species in the pre-dam Colorado River in Glen Canyon and near Lees Ferry. S. Johnson recalled collecting turtles from the mouth of the Paria River in the late 1920's (oral communication, LES files). Observations and movie footage of this species were made by T. Nichols in Glen Canyon in the 1950's. Woodbury et al. (1959) reported several records of *C. picta* in Glen Canyon prior to the construction of Glen Canyon Dam. The species is now extremely rare in Arizona, and has been extirpated from the CRE.

8) Gambel's quail (*Callipepla gambelii*) were detected up until the early 1990's along the lower CRE (Brown et al. 1987, L.E. Stevens unpublished field notes), but have not been reported recently. This species may have been introduced into the CRE, but more information is needed, as it was previously abundant along the western South Rim (Brown et al. 1987). Although common on the North Rim, wild turkey (*Meleagris gallopavo*) occasionally colonize the river corridor, but populations rarely persist more than 1-2 yr.

9) Numerous vagrant and accidentally occurring bird species have been detected in the CRE. At least 61 waterbird species in five guilds have been detected in the CRE: shorebirds, waders, raptors, waterfowl, and kingfishers (Brown et al. 1987, Stevens et al. 1997a, LaRue et al. 2001). Dam operations may affect waterbird populations directly (e.g., by forcing waterfowl and BAEA to shift their feeding strategies) or indirectly (e.g., stable flows may permit expansion of shoreline vegetation, increasing breeding habitat for mallards (*Anas platyrhynchos*), but eliminating spotted sandpipers (*Actitis macularia*) breeding habitat – no breeding sandpipers have been reported in recent decades in the CRE. Many waterbird species occur in migration and use the river corridor as wintering habitat. Protected as a migratory bird species, but not otherwise federally listed, white pelicans (*Pelicanus erythrorhynchos*) regularly pass through Grand Canyon during the migratory period (April and October; Stevens et al. 1997a). Single individuals or small flocks are regularly reported, and flocks in excess of 200 birds have been sighted on the river (LaRue et al. 2001). Concern about range connectivity for this species may result in increased conservation attention in the future (e.g., Anderson and King 2005). Several waterbird species occur as vagrants or accidental species, including: 2 records of magnificent frigatebird (*Fregata magnificens*), several records of trumpeter swan (*Cygnus buccinator*), a lone harlequin duck (*Histrionicus histrionicus*, seen between miles 49-52 in 2006-2007), and a single record of a snow goose (*Chen caerulescens*) observed at Lees Ferry in April 2011. Similarly, many rare terrestrial bird species occur as accidental or vagrant species in the CRE (e.g., painted bunting - *Passerina ciris*; groove-billed ani - *Crotophaga sulcirostris*; scissor-tailed flycatcher – *Tyrannus forficatus*). We excluded these vagrant or accidental bird taxa from the analysis unless they were ESA- or specifically federally protected.

10) Gila monster (*Heloderma suspectum*) are known from a very few detections in the lower CRE, primarily between Mile 202 to Diamond Creek on the left side of the river.

11) In the absence of monitoring, the impacts of dam operations on riverine amphibians may go undetected. For example, Rocky Mountain toad (*Anaxyrus w. woodhousei*) diet, movement,

reproduction success, and potential hybridization with other congeneric taxa (i.e., red-spotted toad, *A. punctatus*) may be affected by season, flow, and nearshore habitat availability, but the ecology of this species in relation to flow regulation has yet to be examined in the CRE.

### **CRE Native Species Rarity**

Three general patterns of rarity appear to exist among rare but unlisted species in the CRE. Relatively few species are unique to the CRE (Stevens and Polhemus 2008), although Grand Canyon endemic *Flaveria macdougalii*, *Euphorbia arron-rossii*, other endemic plant taxa, and several insect taxa occasionally occur along the river. Most endemic taxa in Grand Canyon appear to exist in springs or in harsh but evolutionarily persistent habitats (Phillips et al. 1987).

A more pronounced pattern of rarity involves propagules of headwaters species transported via hydrochory or gravity into the CRE, become established, but are too rare or physiologically stressed to reproduce. These “waif” species include the following examples. Only 3 individuals of poison ivy (Anacardiaceae: *Toxicodendron rydbergii*) are known at CRE springs (32R, 136R, and 142R). A single individual of Asteraceae: *Aster utahensis* was detected at 123L in the middle riparian zone by M. Yeatts (Hopi Tribe, personal communication). Only a few individual of common dunebroom (Fabaceae: *Pareyella fillifolia*) exist in the CRE (Lees Ferry, RM 68R, RM 91R). Four individuals of Euphorbiaceae: *Reverchonia arenaria* were detected at Mile 142R by Ayers and Stevens (1994). Many other unlisted waif plant species are rare and known only from  $\leq 3$  collections in the CRE. The presence of Glen Canyon Dam likely limits downstream dispersal of waif propagules (e.g., Shafroth et al. 2002), and likely has interrupted long-term distributional processes that may reduce CRE riparian diversity.

A third process is the upstream immigration into the CRE through anemochory or zoochory (wind or animal dispersal), and is more likely to occur in lower reaches of the river corridor. For example, the range of desert mistletoe (*Phoradendron californicum*) extends upriver as far as Mile 159R in catclaw. Scale-dependent dispersal (Aukema 2004) and the laxative properties of mistletoe berries (Willson and Travaset 1992) prevent Phainopepla (*Phainopepla nitens*), western bluebirds (*Sialia mexicana*) and other mistletoe seed dispersers from reaching the next *Acacia* upstream in the steep, narrow, sparsely vegetated Muav gorge. Similarly, whipple yucca (Liliaceae: *Yucca whipplei*), creosotebush (*Larrea tridentata*), ocotillo (*Fouquieria splendens*) and other Mohave Desert plant species have extended their ranges upstream as far the Muav Gorge, but have been blocked by the steep topography of that reach (Stevens and Polhemus 2008). In contrast, desert-broom (*Baccharis sarothroides*) likely colonized upriver via anemochory and has extended its range upstream through the CRE to the upper Muav Gorge. Its germination requirements involve unvegetated silty fine sand and moist soils after flooding (Stevens and Waring 1985), conditions which rarely exist in the contemporary CRE. Its continued expansion into the eastern basin of Grand Canyon is likely reduced by the coarser grainsize of the post-dam river corridor (Stevens 1989). Upstream colonization is generally slower than downstream or gravity-based waifing recruitment and non-native plant distribution, although both processes are important long-term biogeographic processes in the CRE riparian zone (Stevens and Waring 1985).

### **Non-native Species**

While the focus of this report is on native TMC, the diversity and ecological roles of non-native taxa deserves mention. The total number of non-native taxa in the CRE likely exceeds 300

species (Stevens and Ayers 2002). The list of non-native species is strongly dominated by more than 200 species of plants, several of which strongly preempt colonization space, increase fire frequency, and affect food resources and pollinator populations (e.g., *Tamarix* spp., *Bromus* spp.). Several dozen non-native invertebrate taxa have been detected in the CRE, including: important fish parasites (e.g., Asian tapeworm, *Bothriocephalus acheilognathi*; and anchorworm, *Lernaea cyprinacea*; Hoffnagle and Cole 1999); important food species for fish (e.g., *Gammarus lacustris*) and birds (e.g., tamarisk leafhopper, *Opsiurus stactogalus*); nursery escapees (e.g., *Anatis lecontei* ladybird beetles); nuisance species (e.g., New Zealand mudsnail, *Potomopyrgus antipodarum*; rock dove – *Columba livia*, European starling – *Sternus vulgaris*, and house sparrow- *Passer domesticus*); and non-native predators (e.g., *Procambarus* crayfish and softshelled turtles in lowermost Grand Canyon. Pending invasions by other invertebrates, such as (*Dreissena rostriformis bugensis*) guagga mussel and *Halyomorpha halys* stinkbugs, may substantially alter CRE structure. Among CRE vertebrates, approximately 19 non-native fish species have been detected in the Colorado River in recent decades, with salmonids implicated in declines in native fish (Marsh and Douglas 1997; Yard et al. 2011). Feral burros were intentionally removed from the CRE by the NPS in 1982, but burros remain federally protected.

Non-native species, as well as several native populations that have proliferated in post-dam time, strongly affect ecosystem form and function in the CRE. However, interactions among these species are complicated and require further study before restoration of native populations. Among several examples are the following:

- 1) Although non-native rainbow trout (RBT) removal around the Little Colorado River confluence coincided with stabilization of a declining HBC population, field data and modeling indicates that HBC recovery began prior to trout removal. In addition, the role of relatively warmer water during the same time period confound conclusions regarding the benefit of nonnative fish removal (Coggins 2008; Yard et al. 2011).
- 2) Translocation of HBC into Shinumo Creek (RM 109R) is being undertaken following attempts to remove non-native rainbow trout by electrofishing (GCWC 2008, NPS 2011); however, robust populations of native speckled dace (*Rhinichthys osculus*) and bluehead sucker (*Catostomus discobolus*) co-exist with RBT in that stream, indicating that in some settings, native and nonnative fish species coexist over long time periods. This topic merits additional research.
- 3) Large Goodding's willow trees commonly co-occur with non-native tamarisk, and the willow has largely been lost to beaver foraging (the beaver population in Grand Canyon is likely much larger than in pre-dam time; Mortenson et al. 2008). Therefore, successful restoration of Goodding's willow stands requires both the elimination of space-preempting nonnative plant species as well as protection of plantings from beaver attack.
- 4) SWFL and other native neotropical migrant birds in the CRE preferentially nest in non-native tamarisk (Brown and Trossett 1989), a common phenomenon throughout Arizona, and SWFL decline has been related to native brown-headed cowbird brood parasitism. To what extent does SWFL restoration require maintenance of suitable vegetation stands from which cowbirds are excluded?
- 5) The upstream movement of non-native crayfish (*Procambarus clarkii*) from Lake Mead, the potential for downstream dispersal from Glen Canyon and headwater tributaries south of Grand Canyon, and the potential for accidental or deliberate introduction into the CRE poses a serious threat to Grand Canyon's 50 or so perennial tributaries. These are some of

the only remaining largely pristine streams in western North America, and are among the only streams in Arizona not invaded by crayfish. As important natural laboratories and sentinel sites for understanding stream-riparian ecosystem ecology and restoration, these streams should be rigorously protected from crayfish invasion.

- 6) On-going invasion of *Dreissena* mussels into the lower Colorado River basin almost certainly will be followed by invasion into upper basin streams and reservoirs. The impacts of this mussel on the CRE and Grand Canyon tributaries are unknown, but may strongly affect river ecosystem structure and function (Nalepa 2010).
- 7) Conflicting reports exist over the ecological significance of New Zealand mudsnail (NZM) in the CRE. Cross et al. (2010) reported no impact of NZM on the CRE foodbase, while Melis et al. (2010) and Rosi-Marshall et al. (2010) simultaneously reported extensive impacts of NZM on the CRE foodbase.
- 8) Tamarisk leaf-beetles (*Diorhabda* spp.) are presently defoliating tamarisk stands throughout the CRE. Despite the recent closure of the beetle release program, this non-native beetle species is well established upstream of Colorado River, and is presently erupting in the CRE. *Diorhabda* invasion may greatly negatively influence river corridor bird habitat and populations, as most of the two dozen neotropical migrant bird species in Grand Canyon nest commonly or preferentially in tamarisk. The NPS is actively monitoring this developing situation which, next to the completion of Glen Canyon Dam, is likely to be the largest anthropogenic change to the CRE in recent history.


### **Trophic Status of TMC**

TMC are variously distributed from the clear tailwater Glen Canyon reach downstream to through the usually turbid reaches of lower Grand Canyon (Schmidt and Graf 1990; Walters et al. 2000; Fig. 1). Analysis of the trophic organization of the CRE indicates that AMP focal TMC (i.e., HBC, KAS, and SWFL) occupy lower and middle trophic positions (Fig. 2). While those focal species occupy selected aquatic, wetland, and riparian habitats, respectively, they may not provide adequate protection for the full array of TMC and ecological processes. In particular, 2 groups of CRE species have not received sufficient monitoring or conceptual modeling attention: top riparian predators, and rare or endemic taxa, including insects. Additional inventory and monitoring data may be needed on such species to develop a more robust ecosystem stewardship approach that embraces aquatic-terrestrial ecosystem linkage in the CRE.

### **Comparative Evaluation of Restoration Potential**

The approaches to restoration prioritization we investigated progressively encompass additional CRE taxa, habitats, ecological processes, and trophic strata, the understanding of which may contribute to enhanced ecosystem stewardship. The AMP is presently focused on endangered HBC and its habitat needs, with minor attention on KAS and SWWF. Emphasis on HBC focuses attention on the aquatic domain, primarily in lower Marble Canyon and the lower Little Colorado River. Consideration of KAS directs management attention to the wetland habitat of a single riverside spring in upper Marble Canyon, and expanding focus to SWWF brings management attention to specific locations on riparian terraces in Marble Canyon and upper Lake Mead. Thus, the conservation umbrella provided by these 3 listed species focuses management attention on Marble Canyon and upper Lake Mead, and on primary and secondary consumer trophic levels (Fig. 2). Addition of extirpated species to the stewardship list, such as;

## SIMPLIFIED CONCEPTUAL MODEL OF THE COLORADO RIVER ECOSYSTEM, SHOWING THE TROPIC POSITION OF EXTIRPATED AND LISTED FAUNA



numerous niches and microhabitats that are not well-conserved by management of the above species.

## **THE REINTRODUCTION PROCESS**

### **Reintroduction Justification, Planning, Compliance, and Implementation**

Restoration of TMC consistent with NPS and other stewardship agency reintroduction and translocation policies heretofore has been regarded as a tertiary AMP priority, behind improving the population health of federally listed species, and sediment management. Thus, questions of whether, how and when to restore at-risk or extirpated endangered or non-listed species remain unaddressed, but require prioritization. Unfortunately, major information gaps exist with regards to the distribution and present status of many TMC in the CRE. For example, the importance and phenological timing of imprinting on spawning habitat stream waters among young native fishes remains an important but largely unaddressed issue that is highly relevant to fish translocation and reintroduction efforts. Also, the issue of range restoration for native fish remains out-standing. Furthermore, the second largest concentration of HBC occurred at the mouth of the Paria River prior to and at the time of dam construction (Minckley 1973), but HBC have since disappeared from that reach, and little attention to range restoration of HBC has been entertained by the AMWG to date. Such information and discussion will play an important role in the planning and implementation of population reintroduction and restoration actions.

Based on Committee experience with translocation of KAS into Royal Arch Creek and HBC into Shinumo Creek, reintroduction of extirpated CRE species likely will require several phases of effort, including: a) development of the administrative context; b) compiling relevant information, stock assessment, and prioritization, c) plan development, d) multi-agency compliance, e) implementation, potentially with multiple stocking efforts, and f) monitoring and feedback.

a) Administrative oversight for reintroduction can be complex. The NPS has direct oversight of natural and cultural resource management of the CRE, as it carries out NPS statutorily-based actions in behalf of the Secretary of the Interior (U.S. Department of the Interior 2006:45). However, several agencies also have oversight for CRE fish and wildlife management. . Other agencies/entities that advise and assist the NPS on management of Grand Canyon plant and wildlife species include: individual NPS units; the FWS, AGFD; several Native American Indian tribes; the Bureau of Reclamation; the U.S. Geological Survey; the State of Arizona; and various other cooperating stakeholders in the Glen Canyon Dam AMWG. Each of these entities has its own mission, and differences in management focus are the norm, making reintroduction of some extirpated species more difficult. The first step in a reintroduction program is to have clearly defined interagency agreement on the effort and clear identification of funding; however, while the administrative context for reintroduction of extirpated species is explicit in the policies of the NPS and in AMWG Goal 3, reintroduction actions remain administratively challenging. For example, reintroduction of razorback sucker into the CRE is regarded as the easiest and most justifiable action, but discussions of this topic began before 1989 and continue to the present day without any action. Administratively successful examples include translocation of KAS into Royal Arch Creek and HBC into Shinumo Creek (NPS 2011).

b) The second step in CRE reintroduction projects involves compilation of sufficient information on the ecology, life history, diseases and parasites, status of remaining populations as potential stock, reasons for extirpation, trophic role(s), relationships to other species, and other information to support reintroduction efforts. This information needs to be further evaluated to allow the CRE managing agencies to clearly justify and prioritize reintroduction efforts, including where and when reintroduction efforts should be conducted. The prioritization process we used includes consideration of all aspects of reintroduction, including: legal and managerial justification, funding, stock availability and genetic issues, potential socio-ecosystem impacts, , reintroduction logistics, treatment for diseases and parasites, risks and contingency planning, monitoring protocols and schedules, reporting, feedback related to improvement of techniques, public relations, and administrative accounting. Clear definition of such issues is critical to resource management agency's decisions regarding reintroduction or rehabilitation of TMC.

c) Development of a reintroduction, translocation, or population augmentation plan is based on the findings in (b) above. The plan should include all aspects of the reintroduction process, including detailed stock assessment to guarantee, to the greatest extent possible, that the reintroduced stock is the most fit and genetically the most capable of surviving the translocation process and proliferating in the renewed landscape. Parasite treatment may be required, as the stock to be translocated should be parasite-free. Also, it is important to determine the level of population take that an existing native species can withstand and still remain functional. The plan must be regarded as a living document, as it will require continued updating as challenges are discovered in the initial phases of the effort. The plan will require agreement by the cooperating agencies.

d) Compliance requirements for species translocation, augmentation, or reintroduction differ and overlap to various extents among the several stakeholders with wildlife management responsibilities in the CRE. NPS compliance requirements for such activities include attention to autecological (stock source), synecological, cultural, other tribal, recreational, natural quiet, and other resource impacts, in accordance with National Environmental Policy Act and the Code of Federal Regulations (24.4.5i.i, 24.6.a.2). FWS reintroduction requirements involve consideration of Section 7 of the 1973 ESA and other obligations. Wildlife reintroduction actions by the AGFD involve a 12-step reintroduction process (AGFD 1987) and Arizona Game and Fish Commission approval. All wildlife stewardship agencies generally have interacted in proposing and accomplishing previous translocation, augmentation, or reintroduction projects in the region (e.g., KAS, HBC, California condor).

## **DISCUSSION**

Ecosystem management requires adequate understanding of the ecological role and potential for restoration of extant as well as extirpated species and processes. Understanding the feasibility and advisability of restoration of extirpated species is mandated through Goal 3 of the AMWG mission, and also is mandated by the wildlife managing agency stakeholders, most notably the NPS, FWS, and AGFD, and several Native American Tribes. Filling information gaps about missing or declining species is needed to more fully inform the Secretary about Glen Canyon Dam impacts, as well as providing information for the compliance requirements of AMWG cooperating agencies needed to foster reintroduction efforts. Through its multi-stakeholder structure, AMWG is uniquely poised to recommend to the Secretary of the Interior

the technical, interagency, and societal information needs and processes through which to achieve the purposes of AMWG's Goal 3. However, there is at present no commitment that the AMP will fund species restoration efforts, as such activities are the specific purview of the resource managing agencies. Nonetheless, the AMWG can play an important and beneficial multi-agency advisory role to the Secretary in evaluating TMC restoration priorities and potential in the CRE, and AMWG may benefit from monitoring information on reintroduced or translocated TMC.

The responsibility of the AMWG is to advise the Secretary of the Interior on the environmental and economic impacts of Glen Canyon Dam operations, and the potential to mitigate impacts within the context of existing law. Our analysis here addresses a motion from AMWG in relation to Goal 3 of the AMP to compile information on the restoration needs and potential for missing and at-risk native species in the CRE. We provide previously uncompiled background information, and an assessment approaches to prioritize TMC reintroduction potential. We offer this information as guidance to the AMWG, recognizing the diverse views among AMWG stakeholders as to the AMP's responsibilities regarding CRE management. Our primary objective here is to present information on the status and priorities for management of rare and declining CRE species to better help AMWG manage the river corridor as an ecosystem.

Many TMC in addition to HBC, KAS, and SWWF were recognized from the CRE that may warrant management attention, if not by the AMWG, then by other managing agencies. A total of 47 (55 percent) of the 86 TMC considered here had potential restoration scores  $\geq 50$  percent (Table 2; Appendix A), but many lacked critical information on status, distribution, and habitat use. There is considerable opportunity for improving the quality of basic information on TMC in the CRE. In particular, at least 5 species (1 plant, 1 lizard, and 3 non-listed fish) scored higher than HBC in our analysis. Concern for the 3 non-listed CRE fish arose from the desire of the Committee to prevent those species from becoming federally endangered. The zebra-tailed lizard scored above HBC because it would be such an inexpensive and easily-accomplished restoration action.

AMWG may choose to engage in further discussion of the status and restoration potential of the TMC list presented here. Periodic updates on TMC populations would help alleviate conservation concerns over the status of those taxa, and may enhance basic understanding of ecosystem condition and linkages. We hope that the information provided here contributes to those discussions and to improved stewardship of the CRE.

## **ACKNOWLEDGEMENTS**

The members of the Goal 3 Ad Hoc Committee included: L.E. Stevens (GCWC, Chair); Jeff Sorensen and Bill Steward (AGFD); Shaula Hedwall, Sam Spiller, and Pam Sponholtz (FWS); Ted Melis, Bill Persons, and David Ward (GCMRC); Mike Yeatts (Hopi Tribe); Todd Chaudhry, Brian Healy, Martha Hahn, and Melissa Trammell (NPS); Marianne Crawford (Reclamation); and Shane Capron (Western). We thank the ad hoc committee members and the following taxon experts for their efforts scoring the species: L. Makarik and R.V. Ward (NPS-Grand Canyon); Carol Chambers, Neil Cobb, Tad Theimer (Northern Arizona University, Flagstaff); and Richard Bailowitz and R. Roy Johnson of Tucson. Bill Persons and Melissa Trammell provided valuable comments on early versions of this manuscript. We kindly thank Ms. Linda Whetton (Reclamation) for her tireless administrative support of these proceedings. L.E. Stevens efforts on this project were partially supported by the Annenburg Foundation's Explore Program.

**REFERENCES CITED**

- Alagona, P.S. 2004. Biography of a "Feathered Pig": The California Condor conservation controversy. *Journal of the History of Biology* 37:557-583.
- Anderson, D.W. and D.T. King. 2005. Introduction: Biology and conservation of the American white pelican. *Waterbirds* 28 (Special Publication 1):1-8.
- Anderson, S. and G.A. Ruffner. 1988. Effects of post-Glen Canyon Dam flow regime on the old high water line plant community along the Colorado River in Grand Canyon. U.S. Bureau of Reclamation Glen Canyon Environmental Studies Report 22. NTIS No. PB88-183504/AS.
- Andersen, M.E., M.W. Ackerman, K.D. Hilwig, A.E. Fuller, and P.D. Alley. 2010. Evidence of young humpback chub overwintering in the mainstem Colorado River, Marble Canyon, Arizona, U.S.A. *The Open Fish Science Journal* 3:42–50 (<http://www.bentham.org/open/tofishsj/openaccess2.htm>).
- Roberge, J-M. and P. Angelstam. 2004. Usefulness of the umbrella species concept as a conservation tool. *Conservation Biology* 18:76-85.
- Arizona Game and Fish Department (AGFD). 1987. Procedures for nongame wildlife and endangered species establishment projects in Arizona. Nongame and Endangered Wildlife Program, Arizona Game and Fish Department, Phoenix.
- Arizona Game and Fish Department (AGFD). 2006. Arizona's Comprehensive Wildlife Conservation Strategy 2005-2015. State Wildlife Action Plan for Arizona. Arizona Game and Fish Department, Phoenix.
- Aukema, J.E. 2004. Distribution and dispersal of desert mistletoe is scale-dependent, hierarchically nested. *Ecography* 27:137-144.
- Ayers, T.J. and L.E. Stevens. 1994. Noteworthy collections – Arizona. *Madroño* 41:228-229.
- Ayers, T.J., R.W. Scott, L.E. Stevens, K. Warren, A.M. Philips, III, and M.D. Yard. 1994. Additions to the flora of Grand Canyon National Park – I. *Journal of the Arizona-Nevada Academy of Science* 1:70-75.
- Branton, M. and J.S. Richardson. 2010. Assessing the value of the umbrella-species concept for conservation planning with meta-analysis. *Conservation Biology* 25:9-20.
- Brennan, T.C. and A.T. Holycross. 2007. A field guide to amphibians and reptiles in Arizona. Arizona Game and Fish Department, Phoenix. 150 pp.
- Brooks, J. and R.G. Cleland. 2004. A Mormon chronicle: the diaries of John D. Lee (reissue). Huntington Library Press, Los Angeles.
- Brown, B.T. and L.E. Stevens. 1992. Winter abundance, age structure, and distribution of bald eagles along the Colorado River, Arizona. *Southwestern Naturalist* 37: 404-435.
- Brown, B.T. and L.E. Stevens. 1997. Winter bald eagle distribution is inversely correlated with human activity along the Colorado River in Grand Canyon. *Journal of Raptor Research* 31:7-10.
- Brown, B.T. and M.W. Trossett. 1989. Nesting-habitat relationships of riparian birds along the Colorado River in Grand Canyon, Arizona. *Southwestern Naturalist* 34:260-270.
- Brown, B.T., R. Mesta, L.E. Stevens, and J. Weisheit. 1989. Changes in winter distribution of bald eagles along the Colorado River in Grand Canyon, Arizona. *Journal of Raptor Research* 23:110-113.
- Brown, B. T., G. S. Mills, R. L. Glinski, and S. W. Hoffman. 1992. Density of nesting peregrine falcons in Grand Canyon National Park, Arizona. *Southwestern Naturalist* 37:188-193.

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*

- Brown, B.T., L.E. Stevens and T.A. Yates. 1998. Influences of fluctuating river flows on bald eagle foraging behavior. *The Condor* 100:745-748.
- Brown, B.T., S.W. Carothers, and R.R. Johnson. 1987. *Grand Canyon birds*. University of Arizona Press, Tucson.
- Carothers, S.W., R.R. Johnson, and S.W. Atchinson. 1974. Population structure and social organization of Southwestern riparian birds. *American Zoologist* 14: 97-108.
- Carothers, S. W., S. W. Aitchison, and R. R. Johnson. 1979. Natural resources, white water recreation, and river management alternatives on the Colorado River, Grand Canyon National Park, Arizona. *Proceedings of the First Conference on Scientific Research in the National Parks* I:253-259.
- Clover, E.U. and Jotter, L. 1944. Floristic studies in the canyon of the Colorado and tributaries. *American Midland Naturalist* 32: 591-642.
- Coggins, L.G. Jr. 2008. Active adaptive management for native fish conservation in the Grand Canyon—Implementation and evaluation. University of Florida PhD dissertation, Gainesville.
- Coggins, L.G. Jr., and C. Walters. 2009. Abundance trends and status of the Little Colorado River population of humpback chub—An update considering data from 1989–2008. U.S. Geological Survey Open-File Report 2009-1075.
- Coggins, L.G. Jr, W.E. Pine III, D.R. Van Haverbeke, D. Ward, and H.C. Johnstone. 2006. Abundance trends and status of the Little Colorado River population of humpback chub. *North American Journal of Fisheries Management* 26:233-245.
- Cross, W.F., E.J. Rosi-Marshall, K.E. Behn, T.A. Kennedy, R.O. Hall Jr., A.E. Fuller, and C.V. Baxter. 2010. *Biological Invasions*, published on-line at: <http://www.springerlink.com/content/bv834031865h2077/>.
- Dowling, T.E., W.L. Minckley, and P.C. Marsh. 1996. Mitochondrial DNA diversity within and among populations of razorback sucker (*Xyrauchen texanus*) as determined by restriction endonuclease analysis. *Copeia* 1996:542-550.
- Gloss, S., J.E. Lovich, and T.S. Melis, editors. 2005. The state of the Colorado River ecosystem in Grand Canyon. U.S. Geological Survey Circular 1282 (<http://pubs.usgs.gov/circ/1282/>).
- Grand Canyon Monitoring and Research Center (GCMRC). 2008. USGS workshop on scientific aspects of a long-term experimental plan for Glen Canyon Dam, April 10-11, 2007, Flagstaff, Arizona. U.S. Geological Survey Open-file Report 2008-1153, Washington, DC.
- Grand Canyon Wildlands Council, Inc. 2005. (GCWC). Background and feasibility of restoring river otter (*Lontra canadensis*) into the Colorado River in Grand Canyon National Park, Arizona. Grand Canyon Wildlands Council, Inc., Flagstaff.
- Grand Canyon Wildlands Council, Inc. 2008. (GCWC). A National Park Service plan to translocate humpback chub into Shinumo Creek, Grand Canyon: October 1, 2008. Grand Canyon Wildlands Council, Inc., Flagstaff.
- Grand Canyon Wildlands Council, Inc. (GCWC). 2009. Recommendations on riparian habitat restoration in Grand Canyon National Park, Arizona. Grand Canyon Wildlands Council, Inc., Flagstaff.
- Groom, M.J., G.K. Meffe, and C. R. Carroll. 2006. *Principles of Conservation Biology*, 3rd edition. Sinauer Associates, Inc., Sunderland.

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*

- Hinman, K.E. and T.K. Snow, editors. 2003. Arizona Bat Conservation Strategic Plan. Nongame and Endangered Wildlife Program Technical Report 213. Arizona Game and Fish Department, Phoenix.
- Hoffmeister, D. 1986. Mammals of Arizona. University of Arizona Press, Tucson.
- Hoffnagle, T. L. and R. A. Cole. 1999. Distribution and prevalence of *Lernaea cyprinacea* in fishes of the Colorado River and tributaries in Grand Canyon, Arizona. Proceedings of the Desert Fishes Council 29:45-46.
- LaRue, C.T., L.L. Dickson, N.L. Brown, J.R. Spence, and L.E. Stevens. 2001. Recent bird records from the Grand Canyon region, 1974-2000. Western Birds.
- Lavender, D. 1976. In the house of stone and light. Grand Canyon Natural History Association, Grand Canyon.
- Layman, S.A. and M.D. Halterman. 1987. Can the western subspecies of the yellow-billed cuckoo be saved from extinction? Western Birds 18:19-25.
- Lovich, J.E. and T.S. Melis. 2005. Lessons from 10 years of adaptive management in Grand Canyon. Pp. 207-220 in Gloss, S., J.E. Lovich, and T.S. Melis, editors. The state of the Colorado River ecosystem in Grand Canyon: a report of the Grand Canyon Monitoring and Research Center 1991-2004. U.S. Geological Survey Circular 1282, Washington, DC.
- Marsh, P. C. and M. E. Douglas. 1997. Predation by introduced fishes on endangered humpback chub and other native species in the Little Colorado River, Arizona. Transactions of the American Fisheries Society 126:343-346.
- Mast, J.N. and G.L. Waring. 1997. Dendrochronological analysis of Goodding Willow in Grand Canyon National Park. Pp. 101-114 in van Riper, C.III and E.T. Deschler, editors. Proceedings of the Third Biennial Conference of Research on the Colorado Plateau. National Park Service Transactions and Proceedings Series NPS/NRNAU/NRTP-97/12.
- McIntosh, A. 1932. Some new species of nematode worms of the genus *Leucochloridium* Carus, parasitic in birds from northern Michigan, with a key and notes on other species in the genus. Journal of Parasitology 19:1-\_\_\_.
- Melis, T.S., D.J. Topping, P.E. Grams, D.M. Rubin, S.A. Wright, A.E. Draut, J.E. Hazel, Jr., B.E. Ralston, T.A. Kennedy, E. Rosi-Marshall, J. Korman, K.D. Hilwig, and L.M. Schmit. 2008. High-flow experiment at Glen Canyon Dam benefits Colorado River resources in Grand Canyon National Park. U.S. Geological Survey Fact Sheet 2010-3009. Available online at: <http://pubs.usgs.gov/fs/2010/3009/>.
- Meretsky, V.J., D.L. Wegner, and L.E. Stevens. 2000. Balancing endangered species and ecosystems: a case study of adaptive management in Grand Canyon. Environmental Management 25:579-586.
- Miller, D.M., R.A. Young, T.W. Gatlin, and J.A. Richardson. 1982. Amphibians and reptiles of the Grand Canyon. Grand Canyon Natural History Association Monograph 4, Grand Canyon.
- Minckley, W. 1991. Native fishes of the Grand Canyon region: an obituary? Pages 124-177 in National Research Council. Colorado River Ecology and Dam Management. National Academy Press, Washington.
- Mortenson, S.G., P.J. Weisberg, and B.E. Ralston. 2008. Do beaver promote the invasion of non-native *Tamarix* in the Grand Canyon riparian zone. Wetlands 28:666-675.
- Mueller, G.A., C.P. Marsh, G. Knowles, and T. Wolters. 2000. Distribution, movements, and habitat use of razorback sucker (*Xyrauchen texanus*) in a lower Colorado River reservoir. Great Basin Naturalist 60:180-187.

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*

- Myers, R.A., J.K. Baum, T.D. Shepherd, S.P. Powers, and C.H. Peterson. 2007. Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science* 315: 1846-1850.
- Nalepa, T.F., 2010, An overview of the spread, distribution, and ecological impacts of the Quagga Mussel, *Dreissena rostriformis bugensis*, with possible implications to the Colorado River system, in Melis, T.S., Hamill, J.F., Bennett, G.E., Coggins, L.G., Jr., Grams, P.E., Kennedy, T.A., Kubly, D.M., and Ralston, B.E., editors. Proceedings of the Colorado River Basin Science and Resource Management Symposium, November 18-20, 2008, Scottsdale, Arizona: U.S. Geological Survey Scientific Investigations Report 2010-5135, 113-121 p., accessed on July 15, 2010, at <http://pubs.usgs.gov/sir/2010/5135/>.
- Newmark, W.D. 1995. Extinction of mammal populations in western North American national parks. *Conservation Biology* 9:512-526.
- Oláh-Hemmings, V, JR Jaeger, MJ Sredl, MA Schlaepfer, RD Jennings, CA Drost, DF Bradford, and BR Riddle. 2010. Phylogeography of declining relict and lowland leopard frogs in the desert Southwest of North America. *Journal of Zoology* 280:343-354.
- Ozaki, K, M. Isono, T. Kawahara, S. Iida, T. Kudo, and K. Fukuyama. 2006. A mechanistic approach to evaluation of umbrella species as conservation surrogates. *Conservation Biology* 20:1507-1515.
- Paxton, E.H., M.K. Sogge, S.L. Durst, T.C. Theimer, and J.R. Hatten. 2007. The ecology of the southwestern willow flycatcher in central Arizona—a 10-year Synthesis Report. U.S. Geological Survey Open-File Report 2007-1381, version 1.0.
- Phillips, B.G., A.M. Phillips, and M.A.Schmidt Bernzott. 1987. Annotated Checklist of Vascular Plants of Grand Canyon National Park. Grand Canyon Natural History Association, Grand Canyon.
- Pogue, M.G. 2004: A new species of *Schinia* Hübner from riparian habitats in the Grand Canyon (Lepidoptera: Noctuidae: Heliethinae). *Zootaxa*, 788:1-4.
- Rasmussen, D. I. 1941. Biotic communities of Kaibab Plateau, Arizona. *Ecological Monographs* 11:229-275.
- Robbins, K. 2007. Missing the link: the importance of keeping ecosystems intact and what the Endangered Species Act suggests we do about it. *Environmental Law* 37:573-594.
- Rosi-Marshall, E.J., T.A. Kennedy, D.W. Kincaid, W.F. Cross, H.A.W. Kelly, K.A. Behn, T. White, R.O. Hall Jr., and C.V. Baxter, 2010. Short-term effects of the 2008 high-flow experiment on macroinvertebrates in Colorado River below Glen Canyon Dam, Arizona. U.S. Geological Survey Open-File Report 2010-1031 (<http://pubs.usgs.gov/of/2010/1031/>).
- Salzer, M.W., V.A.S. McCord, L.E. Stevens and R.H. Webb. 1996. The dendrochronology of *Celtis reticulata* in the Grand Canyon: assessing the impact of regulated river flow on tree growth. Pp. 273-281 in J.S. Dean, D.M. Meko and T.W. Swetnam, editors. *Tree Rings, Environment and Humanity*. Radiocarbon.
- Schmidt, J.C., R.H. Webb, R.G. Marzolf, R.A. Valdez and L.E. Stevens. 1998. Science and values in river restoration in the Grand Canyon. *BioScience* 48:735-747.
- Shafroth, P.B., J.M. Friedman, G.T. Auble, M.L. Scott, and J.H. Braatne. 2002. Potential responses of riparian vegetation to dam removal. *BioScience* 52:703-712.
- Sigler, W.F. and J.W. Sigler. 1996. *Fishes of Utah: a natural history*. University of Utah Press, Salt Lake City.

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*

- Simberloff, D. 1998. Flagships, umbrellas, and keystones: is single-species management passé in the landscape era? *Biological Conservation* 83:247-257.
- Sinclair, B. J. 2007. A new species of *Wiedemannia* Zetterstedt from Grand Canyon National Park, with notes on additional Nearctic species (Diptera: Empididae). *Entomological Society of Ontario, Journal*, 137:25-30.
- Snyder, N.F.R. and H. Snyder. 2005. Introduction to the California Condor. California Natural History Guides, Ewing, NJ.
- Sorensen, J.A. 2005. Kanab Ambersnail 2005 Progress Report: Status of Translocated Populations and Initial Results from the November 2004 Habitat Mitigation Experiment. Nongame and Endangered Wildlife Program Technical Report 243. Arizona Game and Fish Department, Phoenix, Arizona.
- Sorensen, J.A. and D.M. Kubly. 1997. Investigations of the endangered Kanab ambersnail: monitoring, genetic studies, and habitat evaluation in Grand Canyon and northern Arizona. Nongame and Endangered Wildlife Program Technical Report 122. Arizona Game and Fish Department, Phoenix, Arizona.
- Sorensen, J.A. and C.B. Nelson. 2003. 2003 Status Report on Niobrara Ambersnails in Northern Arizona. Nongame and Endangered Wildlife Program Technical Report 225. Arizona Game and Fish Department, Phoenix, Arizona.
- Soulé, M.E., J.A. Estes, B. Miller, and D.L. Honnold. 2005. Strongly interacting species: Conservation policy, management, and ethics. *BioScience* 55:168-176.
- Southwest Condor Release Team. 2007. A Review of the Second Five Years of the California Condor Reintroduction Program in the Southwest. U.S. Fish and Wildlife Service, Phoenix.
- Spamer, E.E. and A.E. Bogan. 1993. Mollusca of the Grand Canyon and vicinity, Arizona: new and revised data on biodiversity and distributions, with notes on Pleistocene-Holocene mollusks of the Grand Canyon. *Proceedings of the Academy of Natural Sciences of Philadelphia* 144:21-68.
- Sponholtz, P. and D. Stone. 2004. Monitoring efforts for humpback chub (*Gila cypha*) above chute falls, Little Colorado River. Presentation to Desert Fishes Council 2004:37.
- Stevens, L.E. 1989. Mechanisms of riparian plant community organization and succession in the Grand Canyon, Arizona. Northern Arizona University Ph.D. Dissertation, Flagstaff.
- Stevens, L.E. and T.J. Ayers. 2002. The biodiversity and distribution of alien vascular plant and animals in the Grand Canyon region. . Pp. 241-265 in Tellman, B., editor. *Exotic species in the Sonoran Desert*. University of Arizona Press, Tucson.
- Stevens, L.E. and R.L. Huber. 2004. Biogeography of tiger beetles (Cicindelidae) in the Grand Canyon Ecoregion, Arizona and Utah. *Cicindela* 35:41-64.
- Stevens, L.E. and J.T. Polhemus. 2008. Biogeography of aquatic and semi-aquatic Heteroptera in the Grand Canyon ecoregion, southwestern USA. *Monographs of the Western North American Naturalist* 4:38-76.
- Stevens, L.E. and R.A. Bailowitz. 2009. Odonata biogeography in the Grand Canyon ecoregion, southwestern U.S.A. *Annals of the Entomological Society of America* 102(2):261-274.
- Stevens, L.E. and G.L. Waring. 1985. Effects of prolonged flooding on riparian vegetation in Grand Canyon. Pp. 81-86 in Johnson, R.R., C.D. Ziebell, D.F. Patten, P.F. Ffolliott, and R.H. Hamre, editors. *Riparian ecosystems and their management: reconciling conflicting uses*. U.S. Forest Service General Technical Report RM-120. Ft. Collins.

*GCWC Draft White Paper: DRAFT 7/08/2011 AMP Goal 3 White Paper*


- Stevens, L.E., K.A. Buck, B.T. Brown and N. Kline. 1997a. Dam and geomorphic influences on Colorado River waterbird distribution, Grand Canyon, Arizona. *Regulated Rivers: Research & Management* 13:151-169.
- Stevens, L.E., J.P. Shannon and D.W. Blinn. 1997b. Benthic ecology of the Colorado River in Grand Canyon: dam and geomorphic influences. *Regulated Rivers: Research & Management* 13:129-149. Abstract available online at: <http://www3.interscience.wiley.com/journal/11832/abstract?CRETRY=1&SRETRY=0>.
- Stevens, L.E., T.J. Ayers, J.B. Bennett, K.Christensen, M.J.C. Kearsley, V.J. Meretsky, A.M. Phillips III, R.A. Parnell, J.Spence, M.K. Sogge, A.E. Springer, and D.L. Wegner. 2001. Planned flooding and Colorado River riparian trade-offs downstream from Glen Canyon Dam, Arizona. *Ecological Applications* 11:701-710.
- Stone, D.M. and O.T. Gorman. 1999. Ontogenesis of endangered humpback chub (*Gila cypha*) in the Little Colorado River, Arizona. *American Midland Naturalist* 155:123-135.
- Stone, J.L. and A.B. Queenan. 1967. Tailwater fisheries investigations: creel census and limnological study of the Colorado River below Glen Canyon Dam, July 1, 1966-June 30, 1967. Arizona Game and Fish Department.
- Stone, J.L. and N.L. Rathbun. 1968. Tailwater fisheries investigations: creel census and limnological study of the Colorado River below Glen Canyon Dam, July 1, 1967-June 30, 1968. Arizona Game and Fish Department.
- Stone, J.L. and N.L. Rathbun. 1969. Tailwater fisheries investigations: creel census and limnological study of the Colorado River below Glen Canyon Dam, July 1, 1968-June 30, 1969. Arizona Game and Fish Department.
- Sublette, J.E., L.E. Stevens, J.P. Shannon. 1998. Chironomidae (Diptera) of the Colorado River in Grand Canyon, Arizona, U.S.A., I: Taxonomy and ecology. *The Great Basin Naturalist* 58:97-146.
- Turner R.M. and M.M. Karpiscak. 1980. Recent vegetation changes along the Colorado River between Glen Canyon Dam and Lake Mead, Arizona. USGS Professional Paper 1132, Washington.
- U.S. Department of the Interior. 2006. Management policies: the guide to managing the National Park system. U.S. Department of the Interior, Washington (accessed 25 March 2009 at: <http://www.nps.gov/policy/mp/Index2006.htm>).
- U.S. Fish and Wildlife Service (USFWS). 2010. Reissuance of the 2009 Supplement to the 2008 Final Biological Opinion for the Operation of Glen Canyon Dam. U.S. Fish and Wildlife Service, Phoenix.
- U.S. National Park Service. 2006. Management Policies 2006, U.S. Department of the Interior National Park Service, Washington DC (ISBN 0-16-076874-8).
- Valdez, R.A. and R.J. Ryel. 1997. Life history and ecology of the Humpback Chub in the Colorado River in Grand Canyon. Pp. 3-32 in van Riper, C.III and e.T. Deschler, editors. Proceedings of the Third Biennial Conference of Research on the Colorado Plateau. National Park Service Transactions and Proceedings Series NPS/NRNAU/NRTP-97/12.
- Walters, C., J. Korman, L.E. Stevens, and B.D. Gold. 2000. Ecosystem modeling for evaluation of adaptive management policies in the Grand Canyon. *Conservation Ecology* 4:1[online] URL: <http://www.consecol.org/vol4/iss2/art1>.
- Willson, M. and A.Traveset. 1992. The ecology of seed dispersal. Pp. 85-110 in Fenner, M., editor. *Seeds: The Ecology of Regeneration in Plant Communities*, 2nd. Edition. CAB International, Wallingford.
- Wixom, H. 1997. Jacob Hamblin: His Own Story. Dixie College, St. George.

Yard, M.D., L.G. Coggins Jr., C.V. Baxter, G.E. Bennett, and J. Korman. 2011. Trout piscivory in the Colorado River, Grand Canyon: Effects of turbidity, temperature, and fish prey availability. *Transactions of the American Fisheries Society* 140:471-486.

DRAFT

**APPENDIX A:  
TAXA OF MANAGEMENT CONCERN IN THE CRE,  
SCORING SHEETS, ALTERNATIVE APPROACHES,  
AND COMMENTS BY EXPERTS**

**(Provided in electronic Excel format only)**



**AMP GOAL 3:  
ASSESSING RESTORATION POTENTIAL OF  
TAXA OF MANAGEMENT CONCERN  
IN THE COLORADO RIVER ECOSYSTEM  
DOWNSTREAM FROM GLEN CANYON DAM**

**Larry Stevens  
Grand Canyon Wildlands Council, Inc.**

- **Committee efforts**
- **Taxon list and results**
- **Prioritization**
- **Restoration issues**
- **Motion to AMWG**

## **AMP GOAL 3**

***Restore populations of extirpated species,  
as feasible and advisable.***

**The IN for Goal 3 is RIN 3.1.1:**

*What information (including technical, legal, economic, and policy issues) should be considered in determining the feasibility and advisability of restoring pikeminnow, bonytail, roundtail chub, river otter, or other extirpated species?*

### **AMWG April 2009:**

*In recognition of GCDAMP goals toward management of the Colorado River through an ecosystem approach, AMWG directs the TWG to establish the Species of Concern Ad Hoc Committee (SMCAHC) and requests the participation of GCMRC in that ad hoc committee, to produce a draft report to be presented to AMWG on or before by May 1, 2011, that contains the following with regard to species of management concern in the CRE: **A review of information about an assessment of the status of habitat needs and availability, and ecosystem roles of the species.***

(NOTE: Extension granted to August 2011 AMWG meeting)

## GOAL 3 AD HOC COMMITTEE MEMBERSHIP

bralston@usgs.gov; kdongoske@cableone.net; [Capron@wapa.gov](mailto:Capron@wapa.gov);  
wdavis@ecoplanaz.com; [Lori\\_Caramanian@ios.doi.gov](mailto:Lori_Caramanian@ios.doi.gov);  
lskrzynski@kaibabpaiute-nsn.gov; [michael.yeatts@nau.edu](mailto:michael.yeatts@nau.edu);  
richard.johnson@npgcable.com; Jan\_Balsom@nps.gov; [Martha\\_Hahn@nps.gov](mailto:Martha_Hahn@nps.gov);  
Brian\_Healy@nps.gov; Norm\_Henderson@nps.gov; [pam\\_sponholtz@fws.gov](mailto:pam_sponholtz@fws.gov);  
jasthiriot@crc.nv.gov; Sam\_Spiller@fws.gov; [tmelis@usgs.gov](mailto:tmelis@usgs.gov);  
bstewart@azgfd.gov; jcjordan1@cox.net; [todd\\_chaudhry@nps.gov](mailto:todd_chaudhry@nps.gov);  
rmoore@grandcanyontrust.org; JSorensen@azgfd.gov; [JSorensen@gf.state.az.us](mailto:JSorensen@gf.state.az.us);  
kelly@grandcanyonwildlands.org; mcrawford@usbr.gov; [RV\\_Ward@nps.gov](mailto:RV_Ward@nps.gov);  
Lori\_Makaric@nps.gov; Melissa\_Trammell@nps.gov; [wpersons@usgs.gov](mailto:wpersons@usgs.gov);  
dlward@usgs.gov; Shaula\_Hedwall@fws.gov; [svanderkooi@usgs.gov](mailto:svanderkooi@usgs.gov);  
farvana@aol.com; [mary@maryorton.com](mailto:mary@maryorton.com); numerous independent experts.

## AD HOC COMMITTEE MEETING DATES

2009: 30 Apr; 2010: 3 Jan; 2011: 27 Jan\*, 23 Feb\*, 12 May\*, 9 Jun\*  
Many emails, expert interviews, individual Committee discussions;  
TWG presentation 6/29/2011, approved motion forwarding the report.

\* Conference call

# **BIODIVERSITY AND ECOSYSTEM PROCESSES IN THE CRE**

- ~ 800 plant taxa
- ~ 10,000 macroinvertebrate taxa
- ~ 350 vertebrate taxa
- Present AMP focus on ES: HBC (KAS & SWFL)

## **EXTIRPATED OR DECLINING TMC**

- Which TMC lost in post-dam time, why, supporting info?
- Which TMC are declining, why, supporting info?
- How to prioritize restoration or rehabilitation?

# **METHODS AND ASSUMPTIONS**

## **Development of the List**

- Document rare, declining CRE taxa
  - Literature review
  - Expert opinions (not all experts agree)
- Exclude non-listed non-native species
- List repeatedly reviewed

## **AMP Issues, Possible Benefits**

- Advisory value to SOTI
- No AMP commitment on restoration funding
- Useful information for monitoring AMP concerns

# Prioritization Approaches

- Legal obligations (ESA)
- Multi-agency policy review
- Conservation status (NatureServe)
- Committee ranking - 17 variables, 0 to 4 scoring
  - Scored by experts
  - Unweighted summary score calculated/species
  - Arbitrary 50% score considered as TMC
- Trophic position – ecosystem approach
- Policy “reality check” and Committee discussion

# Prioritization Categories

Influence of Glen Canyon Dam on population

Conservation relevance

Quality of background information

Stock availability for restoration

Legal mandates, requirements

Habitat availability under present dam operations

Habitat availability under realistic future dam operations

Technical feasibility

**Agreement among stakeholders**

Inexpensiveness (low cost=high score)

Time-frame for success (short=high, long=low)

Logistical ease

Ease of compliance

Are genetics issues easily resolved?

Lack of disease transmission potential

Urgency in relation to climate change impacts

Beneficial interactions with other species

<b>Criterion</b>	<b>Number of CRE TMC</b>									
	<b>Plant</b>	<b>Trema- tode</b>	<b>In- sect</b>	<b>Snail</b>	<b>Fish</b>	<b>Amph ibians</b>	<b>Rep- tiles</b>	<b>Bird</b>	<b>Ma mm al</b>	<b>Total</b>
Extirpated	2?	0	2?	0	4	2	2	3 (4)	8	<b>14-23</b>
Federally ES or recent ES	0	0	0	1	5	2	0	9	1	<b>18</b>
Composite Score $\geq$ 50	0	1	3	1	8	3	3	9	19	<b>47</b>
3a. TMC about which little is known	1	0	9	0	3	2	5	6	26	<b>52</b>
3b. TMC of individual agency concern	6	0	3	1	8	4	3	16	21	<b>62</b>
3c. Rare or endemic TMC	1	1	3	1	0	1	0	5	2	<b>14</b>
3d-e. TMC affected by dam, (+ increasing or - decreasing)	6	1	4	1	8	4	4	13	11	<b>52</b>
Reality Check ("yes")	4	0	0	1	6	0	0	2	0	<b>13</b>
Reality Check ("mixed")	3	1	9	0	2	4	6	17	31	<b>73</b>
<b>Total number TMC *</b>	<b>7</b>	<b>1</b>	<b>9</b>	<b>1</b>	<b>8</b>	<b>9</b>	<b>6</b>	<b>19</b>	<b>31</b>	<b>91</b>

# PLANTS

<b>Taxon</b>	<b>Common Name</b>	<b>Total Score (percent of scores for non- blank categories; excluding stakeholder agreement scores)</b>
PLANT	Goodding's Willow (SAGO)	75.8
PLANT	Fremont Cottonwood (POFR)	60.9
PLANT	Western Honey Mesquite	Unscored but high – no OHWZ recruitment
PLANT	Netleaf Hackberry (CELA)	56.3

## Goodding's Willow (*Salix gooddingii*)



Once relatively common along wide reaches of the Colorado River in Grand Canyon; now gone from 7 of 18 post-dam sites, with most plants in poor health and no recruitment; losses due to increased beaver attack and recruitment impacts due to flow alteration.

# GCD Impacts on CRE Plant Diversity?

- Dam may be interrupting long-term colonization
  - particularly downstream and downslope movement & colonization
- Result: Loss of rare plant species over post-dam time

# INVERTEBRATES

<b>Taxon</b>	<b>Common Name</b>	<b>Total Score (percent of scores for non- blank categories; excluding stakeholder agreement scores)</b>
GAST	Kanab Ambersnail (KAS)	65.6
HEX	Vulcan's Well waterbug (BEFL)	62.5
HEX	Viceroy Butterfly (VBF)	56.25
HEX	Masked Clubskimmer	51.6

# KANAB AMBERSNAIL



# ENDEMIC, SMALL POPULATIONS, NO MONITORING, STATUS UNCERTAIN



“Masked Clubskimmer  
(LIBELLULIDAE: *Brechmorhoga pertinax*)”



Wetsalts Tiger Beetle  
(CICINDELIDAE:  
*Cicindela haemorrhagica arizonae*)



Vulcans Well Giant Waterbug  
(*Belostoma flumineum* new ssp?)



Viceroy Butterfly

*Limenitis archippus*

<b>FISH</b>		
<b>Taxon</b>	<b>Common Name</b>	<b>Total Score (percent of scores for non-blank categories; excluding stakeholder agreement scores)</b>
ICHTH	Humpback Chub (HBC)	71.9
ICHTH	Razorback Sucker (RBS)	66.7
ICHTH	Colorado Pikeminnow (CPM)	63.5
ICHTH	Bonytail Chub (BTC)	55.2
ICHTH	Roundtail Chub (RTC)	53.1
ICHTH	Flannelmouth Sucker (FMS)	89.1
ICHTH	Bluehead Sucker (BHS)	89.1
ICHTH	Speckled Dace (SPD)	87.0

# NATIVE FISH OF THE COLORADO RIVER IN GRAND CANYON



Bonytail Chub



Roundtail Chub



Colorado Squawfish



Razorback Sucker

Extirpated



Humpback Chub



Flannelmouth Sucker



Bluehead Sucker



Spotted Dace



Extant

# HERPETOFAUNA

<b>Order</b>	<b>Common Name</b>	<b>Total Score (percent of scores for non-blank categories; excluding stakeholder agreement scores)</b>
HERP	Zebra-tailed Lizard (ZTL)	79.7
HERP	Relict Leopard Frog (RLF)	69.5
HERP	Gila Monster	65.6
HERP	Northern Leopard Frog (NLF)	61.7
HERP	Rocky Mountain Toad	56.3

**Grand Canyon Pink Rattlesnake**



**Northern Leopard Frog  
-9 Mile Spring and Cardenas Marsh**



**Relict Leopard Frog at 248R**



# BIRDS

<b>Order</b>	<b>Common Name</b>	<b>Total Score (percent of scores for non-blank categories; excluding stakeholder agreement scores)</b>
AVES	SW Willow Flycatcher (SWWF)	63.8
AVES	Prairie Falcon (PRFA)	56.9
AVES	Bald Eagle (BAEA)	56.5
AVES	Gambel's Quail (GAQU)	55.4
AVES	Waterfowl	54.7
AVES	Golden Eagle	54.4
AVES	California Condor (CACO)	52.3
AVES	Osprey	50.0



## Southwestern Willow Flycatcher

- One of ~ 24 nesting neotropical migrant bird spp. in GC – an umbrella species?
- Extirpated from historic range, occasionally breeding on upper Lake Mead
- BH Cowbird brood parasitism
- Commonly nest in non-native Tamarisk – habitat changing

## Bald Eagle

- \* Winter along CR in GC upstream from LCR and at several tribs downstream
- \* Feed on trout
- \* Feeding aided by fluctuating flows
- \* Easily disturbed by passing boats
- \* Present status in GC?



## Peregrine Falcon

- \* Largest breeding population on any land management unit in the 48 states
- \* 90 % of prey biomass is waterbirds, mostly related to post-dam clearwater flow



**What about Prairie Falcon?**

# MAMMALS

Total Score (percent of scores for non-blank categories; excluding stakeholder agreement scores)

Order	Common Name	Total Score (percent of scores for non-blank categories; excluding stakeholder agreement scores)
MAMM	Yuma Myotis	65.8
MAMM	Badger (BDGR)	62.5
MAMM	Fringed Myotis	61.7
MAMM	Spotted Skunk (SpSk)	60.9
MAMM	Western Harvest Mouse (WHM)	60.2
MAMM	Allen's Big-eared Bat	60.0
MAMM	Long-eared Myotis	60.0
MAMM	Long-legged Myotis	60.0
MAMM	Western Pipistrelle	59.4
MAMM	Western Small-footed Myotis	58.3
MAMM	Arizona Myotis	56.7
MAMM	Gray Wolf (GW)	55.7
MAMM	Bobcat (Bobc)	55.5
MAMM	Southwestern River Otter (CRO)	54.8
MAMM	Spotted Bat	51.8
MAMM	Northern Grasshopper Mouse (NGM)	50.8
MAMM	California Mastiff Bat	50.0
MAMM	California Myotis	50.0

# MISSING MAMMALS



**Colorado River Otter  
Extinct? Role as mammalian  
fish predator**




**Badger**

**Muskrat: status in CRE?**



# SIMPLIFIED CONCEPTUAL MODEL OF THE COLORADO RIVER ECOSYSTEM, SHOWING THE TROPHIC POSITION OF EXTIRPATED AND LISTED FAUNA



# **ECOSYSTEM STEWARDSHIP IMPLICATIONS**

- **Restoration – Which, if any, TMC?**
  - **NPS, Cooperating Agencies, AMWG discussion and coordination**
- **Monitoring**
  - **Which taxa**
  - **Whether, how, when to include in future work planning**
- **Future budget and flow implications?**
- **Recommendations to AMWG**

# **PROPOSED MOTION**

**TWG recommends that AMWG accept the report of the Species of Concern Ad Hoc Committee entitled, “Assessment of Taxa of Management Concern in the Colorado River Ecosystem, Glen and Grand Canyons, Arizona, USA: Habitat Needs, Availability, and Ecosystem Roles” dated 15 June 2011, which meets the criteria set forth in the AMWG motion of April 2009, calling for “a review of information and assessment of the status of habitat needs and availability, and ecosystem roles” of native species of management concern.**