

on April 20, 1999 Attach 6 Bmaore

Vol. 9, No. 1 April 1999
GSA TODAY
A Publication of the Geological Society of America

INSIDE

- Call for Papers — Page D1
- Electronic Abstract Submission — Page D12

- Registration Issue — June GSA Today



Floods and Sandbars in the Grand Canyon

Ivo Lucchitta, 6969 Snowbowl View Circle, Flagstaff, AZ 86001
Luna B. Leopold, 400 Vermont Avenue, Berkeley, CA 94707


ABSTRACT

Erosion of sandbars and beaches in the Grand Canyon National Park downstream from Glen Canyon Dam has become a major problem that needs to be addressed. Geomorphic and geologic mapping provide a link between sandbar elevations and discharge measurements. This link allows an estimate of discharges that will deposit sand far enough above normal high water to prevent frequent depletions by erosion. The sand is needed to protect habitats and archaeological sites and to maintain beaches used by recreationists. It is proposed that when the Little Colorado is in flood, discharge at Glen Canyon Dam be increased to bring the total discharge to the desired high value. Analysis of the flow records show that such opportunities are presented on the average once in eight years, suggesting that the proposal has a reasonable chance of success.

INTRODUCTION

The Colorado River in the Grand Canyon section in Arizona once fluctuated greatly in its flow. Year-to-year and season-to-season variability was large. Peak discharges ranged from 300 000 cfs (cubic feet per second) to 19 200 cfs, a difference of 16 times. The amount of sediment transported as suspended load was very large. Measurements carried out at the Grand Canyon for the period December 1940 to June 1941 show that, at 50 000 cfs, about 2 000 000 tons were moved per day during the rising stage of the flood,¹ and 500 000 tons during the falling stage, whereas almost 5 000 000 tons per day were moved during the peak of the flood

Grand Canyon continued on p. 2



Glen Canyon Dam at high discharge during the June 1983 flood. The dam is 710 ft (216 m) high. The four jets of water issuing from near the lower right corner of the dam are from the outlet works. Releases from the right spillway are hidden by the cloud of mist and spray near the lower left corner of the dam. The left spillway, whose exit is visible a short distance downstream from the outlet-work jets, was inactive when the photo was taken. Discharge from the powerplant is below river level and not visible. Photo courtesy of David L. Wegner

¹Sediment transported during the rising stage of a flood is much greater than that transported during the falling stage (Leopold and Maddock, 1953).

GSA TODAY (ISSN 1052-5173) is published monthly by The Geological Society of America, Inc., with offices at 3300 Penrose Place, Boulder, Colorado. Mailing address: P.O. Box 9140, Boulder, CO 80301-9140, U.S.A. Periodicals postage paid at Boulder, Colorado, and at additional mailing offices. Postmaster: Send address changes to *GSA Today*, Membership Services, P.O. Box 9140, Boulder, CO 80301-9140.

Copyright © 1999, The Geological Society of America, Inc. (GSA). All rights reserved. Copyright not claimed on content prepared wholly by U.S. Government employees within the scope of their employment. Permission is granted to individuals to photocopy freely all items other than the science articles to further science and education. Individual scientists are hereby granted permission, without royalties or further requests, to make unlimited photocopies of the science articles for use in classrooms to further education and science, and to make up to five copies for distribution to associates in the furtherance of science; permission is granted to make more than five photocopies for other noncommercial, non-profit purposes furthering science and education upon payment of a fee (\$0.25 per page-copy) directly to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923 USA, phone (978) 750-8400, <http://www.copyright.com>; when paying, reference *GSA Today*, ISSN 1052-5173. Written permission is required from GSA for all other forms of capture, reproduction, and/or distribution of any item in this publication by any means, including posting on authors' or organizational Web sites, except that permission is granted to authors to post the abstracts only of their science articles on their own or their organization's Web site providing the posting includes this reference: "The full paper was published in the Geological Society of America's news-magazine, *GSA Today* [include year, month, and page number if known, where article appears or will appear]." GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

SUBSCRIPTIONS for 1999 calendar year: **Society members:** *GSA Today* is provided as part of membership dues. Contact Membership Services at (800) 472-1988; (303) 447-2020 or member@geosociety.org for membership information. **Nonmembers & Institutions:** Free with paid subscription to both *GSA Bulletin* and *Geology*, otherwise \$50 for U.S., Canada, and Mexico; \$60 elsewhere. Contact Subscription Services. **Single copies** may be requested from Publication Sales. Also available on an annual CD-ROM, (together with *GSA Bulletin*, *Geology*, *GSA Data Repository*, and an Electronic Retrospective Index to journal articles from 1972); \$89 to GSA Members, others call GSA Subscription Services for prices and details. **Claims:** For nonreceipt or for damaged copies, members contact Membership Services; all others contact Subscription Services. Claims are honored for one year; please allow sufficient delivery time for overseas copies, up to six months.

STAFF: Prepared from contributions from the GSA staff and membership.
Executive Director: Donald M. Davidson, Jr.
Science Editors: Suzanne M. Kay, Department of Geological Sciences, Cornell University, Ithaca, NY 14853; Molly F. Miller, Department of Geology, Box 117-B, Vanderbilt University, Nashville, TN 37235
Forum Editor: Bruce F. Molnia, U.S. Geological Survey, MS 917, National Center, Reston, VA 22092
Director of Publications: Peggy S. Lehr
Managing Editor: Faith Rogers
Assistant Editor: Vanessa Carney
Production Manager: Jon Olsen
Production Editor and Coordinator: Gaynor A. Bloom
Graphics Production: Gaynor A. Bloom, Joan E. Manly

ADVERTISING: Classifieds and display: contact Ann Crawford, (303) 447-2020; fax 303-447-1133; acrawford@geosociety.org.

Issues of this publication are available as electronic Acrobat files for free download from GSA's Web Site, <http://www.geosociety.org>. They can be viewed and printed on various personal computer operating systems: MSDOS, MSWindows, Macintosh, and Unix, using the appropriate Acrobat reader. Readers are available, free, from Adobe Corporation: <http://www.adobe.com/acrobat/readstep.html>.

This publication is included on GSA's annual CD-ROM, *GSA Journals on Compact Disc*. Call GSA Publication Sales for details.



50% Total
Recovered Fiber
10% Postconsumer

Printed in U.S.A. using pure soy inks.

IN THIS ISSUE

Floods and Sandbars in the Grand Canyon	1	1999 Annual Meeting Call for Papers ...	D1
In Memoriam	2	Letter	14
Memorial Preprints	2	<i>Environmental & Engineering Geoscience</i> Contents	16
Environment Matters	3	<i>Bulletin and Geology</i> Contents	18
SAGE Remarks	8	GSAF Update	19
Farouk El-Baz Award	10	Future GSA Meetings	20
New Service for <i>Bulletin</i> and <i>Geology</i> Authors	10	1999 GeoVentures	21
Washington Report	11	Calendar	22
Cordilleran Meeting Field Trips	12	Classifieds	23

In Memoriam

Richard J. Anderson
 Dublin, Ohio
 February 5, 1999

Peter J. Coney
 Tucson, Arizona
 February 20, 1999

Jesse L. Dally
 Houston, Texas
 February 7, 1999

Julian R. Goldsmith
 Chicago, Illinois
 January 24, 1999

Robert M. Hutchinson
 Golden, Colorado

Terry W. Offield
 Reston, Virginia
 February 5, 1999

Charles E. Sears
 Hampton, Virginia
 July 8, 1998

Memorial Preprints

The following memorial preprints are now available, free of charge, by writing to GSA, P.O. Box 9140, Boulder, CO 80301-9140.

Charles Banks Belt, Jr.
Edward S. Belt

Ralph J. Bernhagen
Michael C. Hansen, Thomas M. Berg

Wallace Woodrow Hagan
Martin C. Noger, James C. Cobb, Elizabeth Wade Hall

Wilson M. Laird
F. D. Holland, Jr.

Robert F. Walters
Daniel F. Merriam, Robert D. Cowdery

Quido Záruba
Jaroslav Pašek

Grand Canyon continued from p. 1

at ~130 000 cfs (Leopold and Maddock, 1953).

The closure of Glen Canyon Dam in 1963 wrought immense changes in the river. The high peaks were eliminated and the maximum controllable discharge—through power-plant and outlet works—is only ~45 000 cfs (1274 m³/s). The sediment, so characteristic of the river, is now deposited in the reservoir behind the dam, and the releases are clean and cold. All this materially affected the riparian zone² in the Grand Canyon, bringing about changes in erosion, vegetation, and the

biotic communities (Johnson, 1991; Stephens and Shoemaker, 1987; Stevens et al., 1995, 1997; Webb, 1996). The annual peak discharges now are usually in the range of 25 000 to 35 000 cfs. Only five times in the 35 years since dam closure has the peak flow in the Grand Canyon equaled or exceeded 45 000 cfs. Such discharges come about when water going over the dam spillways is added to the maximum discharge through power-plant and outlet works.

The curtailment of sediment passage in the canyon has resulted in little replenishment of the sandbars, on which the maintenance of the original riparian community depended. The release of water

²The zone along a river that is directly affected by the river and its fluctuations.

Grand Canyon continued on p. 3



Enhancing Integrated Science: The Leadership Imperative

Cathleen May, Director, Institute for Environmental Education
Mary Barber, Ecological Society of America
Linda Stanley, U.S. Geological Survey

What factors determine the success or failure of integrated scientific efforts? Are these factors scientific themselves? Social? Cultural? Are they fundamentally dollar-based and administrative? Will collaboration necessarily yield integrated approaches and results? Are there things that organizations can do to enable collaborative efforts among scientists?

The Geological Society of America, the Ecological Society of America (ESA), and the U.S. Geological Survey (USGS) organized a workshop, attended by 24 invited participants and 12 organizers, to investigate these questions. The workshop, "Enhancing Integrated Science" was held November 4-5, 1998, in Reston, Virginia. The participants represented life sciences, earth sciences, economics, social sciences, science administration, and institutional and agency leadership. The results of the workshop illustrate its emergent focus on leadership roles and include a set of "guiding principles" for integrated science, a set of recommendations for the USGS, and a set of recommendations for the larger community of science-enabling organizations and institutions. A summary report, including the results listed above as well as operating definitions, a list of participants, agenda and discussion topics, candid quotes from participants, and an action plan for disseminating the results of the workshop can be found on the USGS Website at: www.usgs.gov/integrated_science/index.html.

The process of organizing the workshop became a microcosm—a real-time illustration of the social and cultural progression by which individuals become a functional collective. A shared vision of the approach to the workshop evolved as individuals came to rely on one another's perspective of the problems we would tackle. Shared responsibility for the products of the workshop evolved as we came to trust each other's contribution during the workshop. These attributes—shared vision, interdependence, and trust—appear to be essential elements of genuine collaboration. Many of the features of the systems in which we work (academic, governmental, or otherwise) are inimical to interdependent, collaborative working relationships. This is not news. Science is a competitive endeavor.

How do we get beyond competition? One of the overarching understandings to emerge from the workshop was that leadership is imperative. If we are to expand the culture of science to include the collaboration so essential to integrated

scientific approaches, individuals and institutions alike must embrace the leadership imperative. Only leadership can modify funding structures to remove organizational barriers to collaboration. Only leadership, among the entire scientific community, can expand reward systems to include something other than individual achievement. Leadership is required to ensure that "interdisciplinary" does not come to mean "anti-disciplinary." Scientific societies, academic institutions, funding agencies, and federal agencies all share the leadership challenge. Leadership is required to safeguard the pursuit of deep understandings, rigorous methodologies, and technologies of individual disciplines without which we cannot achieve useful understandings of complex natural systems.

N. Metzger and R. N. Zare (*Science*, v. 283, p. 642-643) directly address the leadership imperative by focusing on science policy, "Federal structures . . . strongly militate against interdisciplinary programs cutting across jurisdictional lines." These authors recommend an ambitious interagency approach, funded by Congress through direct appropriations, to bring interdisciplinary research "from belief to reality." In essence, Metzger and Zare challenged Congress and the federal system to actively embrace the leadership imperative.

Even as the Metzger and Zare article reached the community, one federal agency moved to the fore with a bold action to enable interdisciplinary science. In a February 1, 1999, news release, USGS Director Chip Groat proclaimed his agency's "commitment to integrating USGS's scientific disciplines. . ." and announced budget restructuring that includes a line item for integrated science (see www.geosociety.org/science). This action signals, at least in the USGS, federal recognition of the exigencies of organizational leadership to interdisciplinary science.

GSA, ESA, and the USGS will continue their joint efforts to enhance integrated science by following through on the imperatives for organizational leadership we derived through our workshop last November. Some plans for implementation in the near future are contained in the summary report cited above. Long-range planning for collaborative leadership efforts is in the works. If your organization or institution is interested in active leadership to enable interdisciplinary science, please contact us through Cathleen May, (303) 447-2020 ext. 1956 at cmay@geosociety.org.

Grand Canyon *continued from p. 2*

from the dam has been governed by demand for electricity from the power plant. This demand fluctuates widely and quickly, so the discharge in the river has changed radically hour by hour and day by day. Not only has the riparian ecosystem been affected, but a decline in recreational use of the canyon has caused significant commercial loss.

These various effects have become so serious that scientific teams studying the

issue have recommended various remedies, including the trial release of water at high controllable discharges to test whether such discharges were sufficient to put in motion sand stored in the bed and in side draws and perhaps build back at least some of the sandbars. In 1996, a "test flood" consisted of the release of 45 000 cfs for a week; in 1997, a discharge of 31 000 cfs for 48 h was designed to transfer from channel bottom to channel margins at least some of the 2.2 million tons of sand and 2.7 million tons of silt and

clay that had been delivered to the Colorado River by a sequence of floods on the Paria River (Kaplinski et al., 1998). Both flows deposited sand, to the jubilation of all concerned, but an important part of the deposited volume was subsequently washed away (Kaplinski et al., 1998). Several floods on the Little Colorado River in January and February 1993 brought a large amount of new sand into the Colorado

Grand Canyon *continued on p. 4*




Figure 1. The only tributaries shown in this location map are the Paria and the Little Colorado Rivers, which are pertinent to this paper. Rivers are in blue; stream gages are in orange. The Lees Ferry gage measures the discharge from Glen Canyon Dam. The Cameron gage measures the discharge of the Little Colorado River, except for the relatively small flow from a spring near the confluence with the Colorado River. The Grand Canyon gage measures the combined discharges of the Colorado, Little Colorado, and Paria Rivers, plus the contribution of small tributary streams within the Grand Canyon, and minus seepage and evaporation losses. Oak Cave and Tsegi Wash, in green, are the location of Holocene deposits discussed in the text.

THE RECENT GEOLOGIC PAST

Extensive and reliably dated Quaternary deposits of the Colorado River in the Grand Canyon go back about 500 000 yr (Lucchitta et al., 1995, 1999), and perhaps about 750 000 yr (Machette and Rosholt, 1991). There have been nine discrete levels of deposition, datable by modern techniques, that are seen in the canyon varying from 10 to 205 m above the present river. Older and higher levels undoubtedly were once present, but have been eroded.

Most deposits consist of remnants of terraces produced by aggradation and underlain by far-traveled river gravel locally intermixed with coarser and more angular debris of local derivation from nearby tributary washes. Taken as a whole, the gravel terraces indicate overall downcutting through the Quaternary, interrupted periodically by aggradation, probably in glacial or late-glacial times. Cobbles greater than 10 cm are common in the gravel, indicating considerable energy and discharge. Tributary rivers and washes mimicked the activities of the main river.

The data show that the Colorado River has cut down an average rate of ~0.4 m/ka for this time interval, within the 0.4–1.09 m/ka range calculated (Lucchitta, 1988) for carving the Grand Canyon as a whole. In reality, this rate averages intervals of aggradation (negative downcutting) with intervals of downcutting at a faster, but unknown, rate. An estimate of the rate at which erosion occurs following a period of aggradation is given by the most recent deposits.

Near-Stream Terrace Levels

The more recent depositional units and sandbars of immediate concern are those that stand at elevations ~10 m or less above the present river. These deposits

Grand Canyon continued from p. 3

River and built conspicuous beaches throughout the Grand Canyon. A year later, these beaches were largely gone (I. Lucchitta, 1994, personal observation).

This paper offers an additional or alternative plan to rebuild the sandbars utilizing floods on the Little Colorado River. The plan is designed in such a way that important parts of the deposition are above the usual moderate peak discharges, so would be less likely to be eroded away.

THE PROPOSAL

A significant amount of sand is brought into the canyon from tributaries entering downstream of the dam, especially the Little Colorado River, which has a large drainage area, more than 26 000 mi² (67 340 km²). The Paria, with an area of 1570 mi² (4082 km²), is helpful (see Fig. 1).

It is proposed that when the Little Colorado approaches a significant flood discharge, the operators at Glen Canyon Dam open penstocks and outlet works so that a combination of high flow from the tributary added to a maximum release from the dam would produce a discharge in the Grand Canyon sufficiently high that its effects may provide on a small scale what the original

undammed river accomplished in sand movement and deposition.

This study uses information and data from several disciplines to examine the feasibility of the proposal. Such analysis involves more than the hydrology and hydraulics of the river basin but must perforce deal with the geomorphology of the canyon bottom, which gives an insight into the long-term functioning of the river. Then, in order to estimate what a flood of any size might deposit, it is necessary to analyze in simplified form the various strandlines and terraces, and determine their dates of deposition or renewal.




Figure 2. Strandlines and terraces in eastern Grand Canyon. Shown schematically are terrace treads and risers, strandlines marked by driftwood lines, and the vegetation types characteristic of each terrace level.

include the last aggradational terrace, as well as terraces produced during the current period of erosion, which includes the decades since Glen Canyon Dam was built. Geomorphic and geologic studies (Lucchitta and colleagues) provide a general profile of the units in this range, shown in Figure 2.

The archeological unit—the last aggradational terrace—is given this informal name (Lucchitta et al., 1995) because prehistoric Puebloan people lived and farmed on these deposits, leaving abundant evidence of their passage in the form of sites and artifacts of many kinds (Fairley et al., 1994). The unit corresponds in a general way to Hack's (1942, 1945) Tsegi Formation, present in washes tributary to the Colorado River in the southwestern Colorado Plateau. The relation of this formation to Puebloan artifacts and to the present channel is similar to those of the archeological unit, as is the fine grain size. Ages are comparable.

The archeological unit consists of a terrace whose base (strath) is below present river level, and the top (tread) about 10 m above it. The unit is composed predominantly of fine to very fine grained sand. The grain size, excellent rounding, and common frosting of the sand grains suggest derivation from Mesozoic eolian sandstone units abundant on the Colorado Plateau. Far-traveled material is scarce in most places. The sand grades laterally into coarser colluvium derived from local slopes and washes, giving the unit a striped appearance.

The archeological unit commonly is modified by wind action and overlain by resulting eolian deposits. In many places, these deposits bury Pueblo II sites built on top of fluvial material. These relations show that the unit stopped being deposited 800–750 yr B.P., and has been incised ~10 m since that time. The oldest age obtained so far from the unit is 5259–4985 calibrated ^{14}C yr B.P. (Lucchitta et al., USGS data) from a level ~3 m below the top of the unit, and at least 7 m above the base. This suggests that the beginning of deposition may well have been in early Holocene—or even late Pleistocene—time.

Pollen associated with the sediments in Oak Cave, southern Utah (Fig. 1), records the initiation of a regime of heavy summer rains, indicated by the expansion of ponderosa pine, which requires summer rain, and the stripping of interfluvial soil mantles, indicated by the waning of sagebrush, which requires deep soil (Lucchitta and others, USGS data). The late Pleistocene to early Holocene age of these sediments either slightly precedes that of the archeological unit, or corresponds to early deposition of this unit.

These depositional, age, and palynological data are interpreted to indicate regional stripping of soil mantles, with accompanying overloading and aggrada-




Figure 3. Discharge rating curve, Grand Canyon station. Many of the deviations from the curve are due to measurements being taken on the rising or falling stages of a flood.

tion of the regional drainage network resulting in deposition of the archeological unit and correlatives. We prefer the aggradational origin for the unit (the grade of the river increased with time) to the origin proposed by O'Connor et al. (1994), which invokes slackwater deposition from large floods (the grade remained constant), for the following reasons, among others: (1) Deposits correlative with the archeological unit are regionally widespread, and not restricted to the Colorado River and its floods. (2) Deposits of tributaries to the Colorado River are graded to the top of the unit, indicating general aggradation of the drainage network. (3) The unit is present in all types of reaches, not just back-eddy areas downstream from constrictions, where slackwater deposits would be typical.

The end of alluviation and onset of vigorous downcutting that incised the archeological unit started about 800 yr ago; downcutting has occurred at the high rate of 12.5 m/ka. The significance of these data is that the last period of major aggradation by the Colorado River is represented by the archeological unit, whereas the river has been operating in a regimen of strong downcutting for approximately the past 800 yr, continuing to this day. Glen Canyon dam was built during a strongly erosive regimen.

Among the levels below the archeological unit are two prominent terraces traceable along many reaches of the Grand Canyon. They have been named the Green Arrowweed and the Silver Arrowweed terraces, respectively (Lucchitta et

Grand Canyon continued on p. 6



Figure 4. Photo of beach on the Colorado River produced by the 1993 flood of the Little Colorado River. The beach is a short distance downstream from the confluence. The beach must have resulted from the flood on the Little Colorado because no such beaches were present upstream from the confluence, and because the sand contained abundant basaltic detritus. Basalt is common along the course of the Little Colorado River. One year later, the beaches visible in the photograph were virtually gone.

Grand Canyon continued from p. 5

al., 1995), from the local flora found there when the studies were carried out (Fig. 2). All levels below the archeological unit have been created and refurbished by floods during the historic period.³ Lines of flood debris and driftwood are found in consistent relations to treads and risers of the terraces, as shown in Figure 2. The younger (and lower) strandlines of large driftwood include abundant worked wood and artifacts such as cans. Plastic objects are common in the 1983 strandline, but very rare in the pre-dam 120 000–140 000 cfs strandline. In pre-dam time, beaches and strandlines produced by lesser floods tended to be modified or destroyed by the higher floods, so they cannot be recognized easily. Floods greater than 120 000–140 000 cfs were rare; the 1884 flood of 300 000 cfs (8490 m³/s)⁴ is marked locally by deposits and a strandline composed of large rotten logs, scarce worked wood, and a few artifacts that include no plastic items. The 97 300 cfs (2755 m³/s) flood of 1983, which marks the greatest post-dam discharge, is represented by a sand terrace and a drift line composed chiefly of fresh, relatively small wood and modern artifacts, including abundant plastic items. The various strandlines were surveyed by total station (Lucchitta et al., 1995, and USGS data).

The test flood of 1997 covered only the Green Arrowweed terrace. The 1996 test flood reportedly just reached the lower end of the Silver Arrowweed terrace (various sources, personal communications). Subsequent discharges have removed a considerable volume of the sand deposited by the test floods (personal observations by Lucchitta in 1998; Kaplinski et al., 1998; and Hazel et al., 1999).

Relation of Terraces to Floods

The geomorphic terraces must be placed within a hydrologic context. This is best done by relating the levels of the terraces to the discharge rating curve, which correlates gage height with discharge. The best gaging station to use for this purpose is that at Grand Canyon, near Bright Angel Creek. Measurements by current meter at this station began in 1921, but a reliable estimate of 300 000 cfs at Lees Ferry was also obtained for the flood of July 8, 1884 (see above).

A plot of the rating curve (Fig. 3) shows that there has been no progressive

change during the period of record. The rating curve for Grand Canyon is a reasonable approximation of the action of the river in the canyon and, combined with the general cross section provided by geomorphic studies, is used here to represent the relation of terrace heights to discharge.

From Figure 3, it can be seen that a discharge of about 55 000 cfs, corresponding to a gage height of 19 to 20 ft (6 m), is required for water to reach a significant part of the Silver Arrowweed terrace. However, an appreciable deposit of sand probably will require 65 000 to 70 000 cfs and a gage height of about 22 ft (7 m). Such a deposit would be relatively durable because it would be well away from the lower step of the terrace, and thus from the erosive action of the river.

In the 35 years since the dam began controlling the flow, only six events equaled or exceeded the 45 000 cfs of the 1996 "test flood" and of these, only two were significantly higher, 58 400 on June 15, 1965, and 96 200 cfs on June 29, 1983. All events exceeding 45 000 cfs required a contribution from the spillway of the dam.

The Little Colorado River can provide the 10 000 to 25 000 cfs needed to raise the maximum controllable outflow from the dam (45 000 cfs) to the required discharge. The floods of the Little Colorado in January and February 1993 indicate that not only is such a contribution possible, but that such floods also bring much sand into the Grand Canyon. The peak discharge of these floods was 18 200 cfs at Cameron, the sixth largest in the period of record. On March 3, a few weeks after the peak, the magnificent beaches (Fig. 4) produced by these floods lined the length of the river from the Little Colorado River confluence to at least as far as Diamond Creek, Mile 226 (Lucchitta and colleagues, personal observations).

ON THE POSSIBILITY OF RESTOCKING SAND ON THE BEACHES

In contrast to the test "floods," which essentially mined the sand in the channel of the Colorado River—a depletable resource—or redistributed sediment brought in by floods on the Paria—which are puny compared to the sediment transported by the Colorado River in pre-dam times—the Little Colorado flood introduced enough new sand to form beaches the length of the river below the confluence (Fig. 4), in addition to the sand that remained in the channel or made its way to Lake Mead. Such floods provide an opportunity for restocking beaches in the Grand Canyon downstream from the confluence.

First, one must achieve a combined Colorado–Little Colorado River discharge sufficient to keep sand entrained rather than allowing it to settle in the channel,

and this discharge must be maintained long enough to build adequate beaches. Second, the discharges must attain a high enough stage to park the sand where it will not be readily eroded by the river in its normal modern flows.

In the period of record, seven days of the highest peak discharges (> 18 000 cfs) of the Little Colorado at Cameron included four that increased the discharge at Grand Canyon enough to overflow the Silver Arrowweed terrace. These four occurrences in the 50 years of measurement means one occurrence about every 12 years.

Alternatively, one may estimate the possible contribution by counting the number of days when the daily flow near Cameron exceeded 10 000 cfs. The daily flow often is high on two to four successive days, which would mean little attenuation by channel storage. In the 50 years of record, there were six such occurrences, two of which were not identical with the seven analyzed above. Thus, there appears to be the possibility of six useful events in 50 years, or one in eight years on the average.

The concept is that the flow of the Little Colorado River be monitored at the most downstream gaging station at Cameron, Arizona. When the discharge reaches a high enough level, the operator of Glen Canyon Dam would, on short notice, increase the discharge through power-plant and outlet works to the maximum capacity of 45 000 cfs. A direct radio link from gage to dam would make the opportunity practical. Furthermore, advance notice of a possible peak at Cameron could be obtained from careful observation of the gages on the Little Colorado River upstream of Cameron.

Given the potential for useful events to occur on average once in eight years, it is quite within reason that the high flow of the Little Colorado River could be made to coincide with a release of 45 000 cfs from Glen Canyon Dam, resulting in the deposition of abundant sand at a substantial elevation above present river grade, within the Silver Arrowweed terrace. In this position, sand would have considerable durability, probably enough to last until the next restocking event.

The proposal discussed here pertains to a geological process—the restocking of beaches. However, it is well known that proper managing of the Grand Canyon involves a complex and often contradictory interplay of geological, biological, and other considerations (Schmidt et al., 1998). All of these must be considered when making management decisions. Nevertheless, the proposal attempts to recreate—to the extent possible now that the dam is built—the sediment transport and river stage characteristics of the pre-dam river. These are the conditions to which the Grand Canyon ecosystem was originally adjusted.

³Defined here as the time interval, starting in 1884, for which observational information is available.

⁴This discharge is based on the observed level reached by the flood at Lees Ferry—within the orchard of Lee's Lonely Dell Ranch. In the 1920s, a profile was surveyed from this level to the river, allowing determination of the channel cross section during the flood. This, in turn, led to an evaluation of the flood discharge.

ACKNOWLEDGMENTS

The U.S. Bureau of Reclamation's Glen Canyon Environmental Studies program supported Lucchitta's work in the Grand Canyon. A U.S. Geological Survey Gilbert fellowship to T. C. Hanks and Lucchitta supported the work on the Rainbow Plateau, where Oak Cave is. The Grand Canyon National Park staff kindly allowed research to be conducted within the park. Our colleagues at the U.S. Geological Survey's Water Resources section in Flagstaff, especially Frank Brewsaugh and Bob Hart, supplied us with river data and other assistance. Mark Caffee and Robert Finkel of the Center for Accelerator Mass Spectrometry at Lawrence Livermore National Laboratory provided the cosmogenic radio-nuclide exposure ages and radiocarbon ages. Sid Davis and Marie Davis of Davis² have concerned themselves with soils and paleoagriculture. James Hasbargen and Susan Smith of Northern Arizona University studied the palynology of the archaeological unit and equivalents. Our river-guide friends, for whom nothing is impossible, especially Steven Bledsoe, Kelly Burke, Brian Dierker, and Chris Geanious, made the study feasible by providing safe conveyance through the fastness and perils of the Grand Canyon. Finally, detailed reviews by L. L. Ely, Karl Karlstrom, J. L. Pederson, D. J. Topping, and two anonymous reviewers helped us to improve the manuscript. To all these individuals and organizations, thanks.

REFERENCES CITED

- Fairley, H. C., Bungart, P. W., Coder, C. M., Huffman, J., Samples, T. L., and Halsom, J. R., 1994, The Grand Canyon river corridor survey project: Archaeological survey along the Colorado River between Glen Canyon Dam and Separation Canyon: National Park Service Cooperative Agreement No. 9AA-40-07920, 276 p.
- Hack, J. T., 1942, The changing physical environment of the Hopi Indians of Arizona: Papers of the Peabody Museum of American Archaeology and Ethnology, Harvard University, v. 35, no. 1.
- Hack, J. T., 1945, Recent geology of the Tsegi Canyon: Appendix 1 of Archaeological studies in northeast Arizona, in Beals, R. L., et al, eds., Publications in American Archaeology and Ethnology, Volume 44: Berkeley and Los Angeles, University of California Press.
- Hazel, J. E., Kaplinski, M., Parnell, R. A., Manone, M. F., and Dale, A., 1999, Topographic and bathymetric changes at thirty-three long-term study sites: American Geophysical Union Monograph (in press).
- Johnson, R. R., 1991, Historic changes in vegetation along the Colorado River in the Grand Canyon: Washington, D.C., Water Science and Technology Board, Colorado River Ecology and Dam Management, p. 176-209.
- Kaplinski, M., Hazel, J. E., Manone, M. F., Parnell, R. A., Dale, A. R., and Topping, D., 1998, Sediment storage changes following short-duration, high-flow releases from Glen Canyon Dam, Colorado River in Grand Canyon: Geological Society of America Abstracts with Programs, v. 30, no. 6, p. 12.
- Leopold, L. B., and Maddock, T., Jr., 1953, The hydraulic geometry of stream channels and some physiographic implications: U.S. Geological Survey Professional Paper 252, 57 p.
- Lucchitta, L., 1988, Canyon maker: Plateau, v. 59, no. 2, 32 p.

PUBLICATIONS NEWS FROM THE GSA

BOOKSTORE

WATCH THIS COLUMN FOR NEWS ABOUT GSA PUBLICATIONS

MILITARY GEOLOGY IN WAR AND PEACE

edited by J. R. Underwood, Jr. and P. L. Guth, 1998

The classic dilemma for military geology has been whether support can best be provided by civilian technical-matter experts or by uniformed soldiers who routinely work with the combat units. In addition to the introductory paper this volume includes 24 papers, covering selected aspects of the history of military geology from the early 19th century through the recent Persian Gulf war, military education and operations, terrain analysis, engineering geology in the military, use of military geology in diplomacy and peace keeping, and the future of military geology. REG013, 256 p., ISBN 0-8137-4113-0, \$76.00, Member price \$60.80

GEOLOGIC EVOLUTION OF THE BARBERTON GREENSTONE BELT, SOUTH AFRICA

edited by Donald R. Lowe and Gary R. Byrty, 1999

The Barberton Greenstone Belt contains the oldest well-preserved volcanic and sedimentary rocks known on Earth and provides a unique window on surface environments, biological evolution, and crustal development 3,550 to 3,200 m.y. ago. This volume presents a systematic account of the geology of the Barberton Belt and its implications for interpreting surface conditions and processes on early Earth. The subjects include stratigraphy, structural divisions, sedimentary geology, and biological materials of the entire belt; mafic and ultramafic volcanic rocks and their petrogenesis; the sedimentology and volcanology of ultramafic volcanoclastic units; the depositional and tectonic setting of the oldest known foreland basin sediments of the overlying Fig Tree Group; and the petrology and tectonic implications of the upper syntectonic quartzose clastic sediments of the Moodies Group. An interpretative summary of the overall evolution of the Barberton Belt concludes the book. SPE329, 324 p., softcover, indexed, ISBN 0-8137-2329-9, \$60.00; Member price \$48.00

HIMALAYA AND TIBET: MOUNTAIN ROOTS TO MOUNTAIN TOPS

edited by Allison Macfarlane, Rasool Sorkhubi, Jay Quade, 1999

More than 30 international authors contributed to the first Himalaya-Karakoram-Tibet (HKT) Workshop held in America. The 11th HKT Workshop, held April 1996 in Flagstaff, Arizona, was a logical outcome of an increasing involvement of North American geologists in the studies of the Himalaya and Tibet over the past two decades. One of the prominent features of this 21 chapter volume is the integration of both "hard-rock" and "softrock" geology, paleotectonic and neotectonic history, and endogenic (tectonic) and exogenic (geomorphic)

processes in the Himalaya. Fully refereed, seven articles cover the Tibet, Trans-Himalaya, and Tethys unit. The High Himalayan Crystalline zone is the subject of another eight chapters; the final six chapters cover the Himalayan Foreland and its sediments, structures, and landforms. SPE328, 336 p., softcover, 1 large plate, indexed, ISBN 0-8137-2328-0, \$70.00, Member price \$56.00

LATE CENOZOIC XIANSUIHE-XIAOJIANG, RED RIVER, AND DALI FAULT SYSTEMS OF SOUTHWESTERN SICHUAN AND CENTRAL YUNNAN, CHINA

by E. Wang, B. C. Burchfiel, L. H. Royden, Chen Liangzhong, Chen Jishen, Li Wenxin, 1998

The Tibetan plateau and its surrounding mountain ranges and basins are a natural laboratory in which to study geological processes ranging from continental collision tectonics to effects of plateau development on climate. Three active fault systems, the Xianshuihe-Xiajiang, Red River, and Dali, offer clues about the extrusion of crustal fragments eastward from the Tibetan plateau, how far back into time the present pattern of deformation can be projected, and the relation between these fault systems and the intracrustal deformation of the India-Eurasia collision zone. The region of these fault systems is an example of rapid changes in partitioning of strain during 5 m.y. in a rotational tectonic regime. SPE327, 112 p., ISBN 0-8137-2327-2, \$41.00, Member price \$32.80

ENVIRONMENTAL EVOLUTION OF THE SOUTHWESTERN UNITED STATES: THE CLARENCE A. HALL, JR., VOLUME

edited by G.W. Ernst and C. A. Nelson, 1998

The serious professional will appreciate the breadth of this book devoted to the integrated study of earth and environmental sciences of the southwestern United States. IBS001, 500 pp., softcover, 7" x 10" format, ISBN 0-9665869-0-5, \$89.95, Member price \$72.00

SHOP GSA'S BOOKSTORE ON THE WEB!
www.geosociety.org

1-800-472-1988

GSA PUBLICATION SALES
P.O. Box 9140, Boulder, CO 80301
(303) 447-2020 or fax 303-447-1133
Prepayment required. Major credit cards accepted.
The Geological Society of America

- Lucchitta, L., Dehler, C. M., Davis, M. E., Burke, K. J., and Basdekas, P. O., 1995, Quaternary geologic map of the Palisades Creek-Comanche Creek area, eastern Grand Canyon, Arizona: U.S. Geological Survey Open-File Report 95-382, 39 p., maps.
- Lucchitta, L., Curtis, G. H., Davis, M. E., Davis, S. W., and Turrin, B., 1999, Cyclic aggradation and downcutting, fluvial response to volcanic activity, and calibration of soil carbonate stages in the western Grand Canyon, Arizona: Quaternary Research (in press).
- Machette, M. M., and Rosholt, J. N., 1991, Quaternary geology of the Grand Canyon, in Morrison, R. B., ed., Quaternary nonglacial geology: Conterminous U.S.: Boulder, Colorado, Geological Society of America, Geology of North America, v. K-2, p. 397-406.
- O'Connor, J. E., Ely, L. L., Wohl, E. E., Stevens, L. E., Melis, T. E., Kale, V. S., and Baker, V. E., 1994, A 4500-year record of large floods on the Colorado River in the Grand Canyon, Arizona: Journal of Geology, v. 102, p. 1-9.
- Schmidt, J. C., Webb, R. H., Valdez, R. A., Marzolf, R. G., and Stevens, L. E., 1998, Science and values in

- river restoration in the Grand Canyon: BioScience, v. 48, no. 9, p. 735-747.
- Stephens, H. G., and Shoemaker, E. M., 1987, In the footsteps of John Wesley Powell: Boulder, Colorado, Johnson Books, 286 p.
- Stevens, L. E., Schmidt, J. C., Ayers, T. J., and Brown, B. T., 1995, Flow regulation, geomorphology, and Colorado River marsh development in the Grand Canyon, Arizona: Ecological Applications, v. 5, no. 4, p. 1025-1039.
- Stevens, L. E., Buck, K. A., Brown, B. T., and Kline, N. C., 1997, Dam and geomorphological influences on Colorado River waterbird distribution, Grand Canyon, Arizona: Regulated Rivers, v. 13, p. 151-169.
- Webb, R. H., 1996, Grand Canyon: a century of change: rephotography of the 1889-1890 Stanton expedition: Tucson, University of Arizona Press, 290 p.
- Manuscript received August 12, 1998; accepted February 8, 1999.*