



United States Department of the Interior

BUREAU OF RECLAMATION
125 South State Street, Room 8100
Salt Lake City, UT 84138-1102



IN REPLY REFER TO:

UC- 410
2.2.4.21

VIA ELECTRONIC MAIL ONLY

Memorandum

To: Chief of Staff
Office of the Assistant Secretary for Water and Science

From: Wayne G. Pullan **WAYNE PULLAN**
Chair, Glen Canyon Leadership Team
Secretary's Designee to the Adaptive Management Work Group (acting)

Subject: Notification of Decision to Not Implement a Fall 2021 High Flow Experiment at Glen Canyon Dam

Digitally signed by
WAYNE PULLAN
Date: 2021.10.05
11:32:52 -06'00'

On September 28, 2021, the Glen Canyon Planning/Implementation Team (PI Team) finalized its recommendation regarding a potential fall High Flow Experiment (HFE) at Glen Canyon Dam in November 2021 (Attachment - *Final Recommendation Regarding a Fall 2021 High Flow Experiment at Glen Canyon Dam, November 2021*). While the PI Team reached consensus in its opposition to a 192-hour extended duration fall HFE, members were divided in the assessment of a shorter 60-hour fall HFE. The recommendation was developed and evaluated within the adaptive management framework and provisions of the 2016 *Record of Decision for the Glen Canyon Dam Long Term Experimental and Management Plan Final Environmental Impact Statement* (LTEMP ROD) concerning annual planning for flow-based experiments.

In accordance with the LTEMP, the Department may make the decision to conduct flow-based experiments (e.g., High Flow Experiments, Bug Flows, Trout Management Flows, and Low Summer Flows) at Glen Canyon Dam if it is determined that there are no unacceptable adverse impacts on other resource conditions. LTEMP states that "Prior to implementation of any experiment, the relative effects of the experiment on the following resource areas will be evaluated and considered: (1) water quality and water delivery, (2) humpback chub, (3) sediment, (4) riparian ecosystems, (5) historic properties and traditional cultural properties, (6) Tribal concerns, (7) hydropower production and WAPA's assessment of the status of the Basin Fund, (8) the rainbow trout fishery, (9) recreation, and (10) other resources." Water Year 2021 was the fourth full year of implementing the process for annual experimental planning under the LTEMP ROD. For future experimental planning, the Department welcomes input from each Leadership Team member as to whether the current process or another process should be used to satisfy the coordination and communication requirements under the LTEMP ROD.

Traditionally Associated Tribes shall be notified at least 30 days in advance of planned experimental flows. On September 20, 2021, notification of the possible fall HFE and offer for consultation was emailed to the Tribes and Parties to the LTEMP National Historic Preservation Act Section 106 Programmatic Agreement (LTEMP PA). No requests for consultation regarding the potential fall HFE were received.

The LTEMP ROD specifies the representation requirements for planning experiments at Glen Canyon Dam and is based on past successful planning and implementation of flow-based experiments. The PI Team includes technical representatives from the Bureau of Reclamation (Reclamation), the National Park Service (NPS), the U.S. Fish and Wildlife Service (FWS), the Bureau of Indian Affairs (BIA), the U.S. Geological Survey's (USGS) Grand Canyon Monitoring and Research Center (GCMRC), Western Area Power Administration (WAPA), the Arizona Game and Fish Department (AZGFD), the seven Colorado River Basin States (States), and the Upper Colorado River Commission (UCRC). The Glen Canyon Leadership Team (Leadership Team) is made up of decision makers from these same agencies and states along with the UCRC.

The Leadership Team has reviewed and considered the PI Team's recommendation, including the assessment of key resources that may be impacted or affected by a 60-hour fall HFE. The Leadership Team met via webinar on September 29, 2021 and, consistent with the PI Team's recommendation, members were divided in their assessment of a 60-hour fall HFE.

The 2021 water year was an historically difficult year for the Western states, including tribes, fisheries, wildlife, farmers, ranchers, and communities. The Colorado River Basin is experiencing its 22nd year of drought and many years of low-runoff conditions. Following two extremely dry periods of runoff in 2020 (54% of average annual inflow) and 2021 (33% of average annual inflow), the two reservoirs (Lake Mead and Lake Powell) are at their lowest levels since they were originally filled. The Colorado River Drought Contingency Plan Authorization Act (P.L. 116-14) directs Interior to "operate Colorado River reservoirs" according to Drought Contingency Plan agreements, including the Drought Response Operations Agreement (DROA), which establishes an objective for Glen Canyon Dam operations to minimize the risk of falling below a "Target Elevation" of 3525 feet (ft). The potential effect of a HFE on the Target Elevation is an appropriate consideration when determining whether to conduct a HFE.

As a response to the hydrologic conditions and the need to protect critical levels at Lake Powell as part of the DROA, Reclamation began releasing water from upper reservoirs to Lake Powell in July under the Emergency Action provisions of DROA. These releases from upper reservoirs are to help reduce the risk of Lake Powell going below the DROA-identified Target Elevation of 3525 ft. These DROA actions will continue through the calendar year. While these actions have helped to reduce the risk for next year, it is likely that additional actions may or will be needed to protect critical levels.

The sediment triggers outlined by LTEMP ROD have been met to consider a 2021 fall HFE. Advocates for the 2021 fall HFE are concerned that the ability to implement HFEs in the future may be limited, should drought conditions continue. They also cite the much-needed sediment

resource benefits from a fall HFE given that it has been three years since the most recent HFE (fall 2018) was triggered and implemented. Implementation of a fall HFE would not impact the annual release volume from Lake Powell, which is set at 7.48 maf in WY 2022. Due to relatively low winter flows, sandbar increases would be retained through winter such that the 60-hour fall HFE would result in a net 20% increase to sandbar size in April 2022 relative to October 2021.

The majority of Leadership Team members indicated that one or more of the following concerns amount to unacceptable adverse impacts on resource areas as identified for consideration by the LTEMP ROD.

Implementation of a 60-hour fall HFE would temporarily reduce Lake Powell reservoir elevation by approximately 2 feet, which reduction is contrary to the objective of an approximately 3 feet of elevation increase made possible due to additional releases from upper basin initial units (Flaming Gorge, Navajo, and Aspinall) currently being implemented under the DROA. As a result, Leadership Team members have indicated that implementation of a fall HFE in November 2021 may result in reduced support for future drought response operations—which releases will likely be necessary to avoid the water elevation dropping below 3,525 at Lake Powell. Further, under most probable hydrology a 60-hour fall HFE is projected to result in up to 15 additional days below 3,525 ft and an annual minimum elevation 0.7 ft lower than if no HFE were to be implemented. While these impacts to reservoir elevation could be mitigated by reallocation (reducing) releases in December, January, and February rather than March, April, and May, doing so would result in a greater direct cost to the Basin Fund for purchase power. The current condition and projections for the Basin Fund have elicited unprecedented actions by WAPA and Reclamation including deferred maintenance, a proposed rate increase, elimination of certain firm supply provisions in contracts, and a one-time pursuit of appropriations for environmental program base funding. The proposed HFE would worsen the condition of the Basin fund; direct costs to the Basin Fund are estimated at \$1.30M for March, April, May reallocation and at \$3.04M for December, January, February reallocation. The majority of Leadership Team members indicated that one or more of these concerns amount to unacceptable adverse impacts.

Based on the concerns expressed by both the Leadership Team and the PI Team, the determination of unacceptable adverse impacts, and the lack of consensus to implement a 60-hour fall HFE, I concur with the determination of unacceptable adverse impacts and I have decided not to conduct a fall HFE in November 2021. We will continue to work with our partners in future HFEs and in the protection of the Grand Canyon and our most important resources.

I am grateful to both the Leadership and the Planning/Implementation Teams for their dedication and commitment to the process for annual experimental planning and for your continued support of the Glen Canyon Dam Adaptive Management Program.

For the hearing impaired please call the Federal Relay System at (800) 877-8339 (TTY).

Attachment - Final Recommendation Regarding a Fall 2021 High Flow Experiment (HFE) at
Glen Canyon Dam, November 2021

cc: Camille Calimlim Touton, Deputy Commissioner

September 28, 2021

To: Glen Canyon Leadership Team for Implementation of Experiments under the Long Term Experimental and Management Plan (LTEMP)

From: Glen Canyon Dam Technical Implementation / Planning team

Re: Final Recommendation Regarding a Fall 2021 High Flow Experiment (HFE) at Glen Canyon Dam, November 2021

I. Recommendation Summary

Based on the LTEMP Record of Decision (ROD), the Glen Canyon Dam technical implementation/planning team (PI Team) has worked over the past several weeks to evaluate existing information and data in determining this recommendation regarding a HFE at Glen Canyon Dam. The PI Team's recommendation regarding implementation of a fall 2021 HFE is based on a careful assessment of resources and best available science.

By consensus, the PI Team is opposed to recommending that the Department implement a 192-hour extended duration HFE in fall 2021.

Technical representatives were divided in the assessment of a shorter duration 60-hour HFE, having evaluated the alternative for its ability to reduce negative resource impacts while still providing sufficient benefit to the sediment resource in terms of learning and effectiveness. The majority of representatives on the PI Team are opposed to recommending that the Department implement a 60-hour HFE this fall, several members support recommending that the Department implement a 60-hour duration HFE this fall, and two members have abstained from making a recommendation.

Purpose of the Recommendation Memo

The purpose of this memorandum is to transmit a recommendation to the Glen Canyon Leadership Team and to the Department in accordance with the LTEMP ROD. The PI Team includes technical representatives from the National Park Service (NPS), U.S. Fish and Wildlife Service (FWS), Bureau of Indian Affairs (BIA), U.S. Geological Survey (USGS) Grand Canyon Monitoring and Research Center (GCMRC), Bureau of Reclamation, Western Area Power Administration (WAPA), Arizona Game and Fish Department (AGFD), seven Colorado River Basin States (States), and the Upper Colorado River Commission (UCRC).

The PI Team has worked over the past several weeks to evaluate existing data and coordinate the potential implementation of an HFE. The PI Team evaluated the latest data from agency experts, and considered multiple issues in making its recommendation, as summarized below. The PI Team arrived at this recommendation after several weeks of PI Team conference calls, and after receiving feedback from Adaptive Management Program stakeholders. The Secretary of the Interior and/or her Designee will consider the recommendation of the PI Team, but retains sole discretion to decide how best to accomplish operations and experiments in any given year

pursuant to the 2016 LTEMP ROD and other binding obligations for Glen Canyon Dam. The PI Team incorporated the most current science and data and considered multiple issues with agency experts, as summarized below, in making this final recommendation.

192-hour Extended Duration Fall HFE

The PI Team arrived at this recommendation after several weeks of PI Team conference calls, and after receiving feedback from Adaptive Management Program stakeholders. **By consensus, the PI Team is opposed to recommending that the Department implement a 192-hour extended duration HFE in fall 2021.** The PI Team identified the following resource concerns associated with a 192-hour extended duration fall HFE: challenges and risks associated with reallocating a large volume of water (415 kaf) while meeting hourly, daily and monthly operating objectives identified in the LTEMP ROD and consistent with a 7.48 maf annual release pattern; uncertainties and risks associated with exacerbating low reservoir elevations (e.g. additional days below 3,525 ft, lower annual minimum); the strain on the already low status for the Basin Fund as projected by WAPA; and potential impacts to non-native fish, including increased brown trout immigration to Lees Ferry and increased risk of fish entrainment at lower reservoir elevations. PI Team members highlighted uncertainties regarding Glen Canyon Dam's ability to reliably and effectively deliver water and generate power when pool elevation drops below 3,525 ft. Further, a highly anticipated spring HFE, if triggered, would be prohibited from occurring in the same water year as an extended duration HFE.

60-hour Fall HFE

Technical representatives were divided in the assessment of a shorter duration 60-hour HFE, having evaluated the alternative for its ability to reduce negative resource impacts while still providing sufficient benefit to the sediment resource in terms of learning and effectiveness. The majority of representatives on the PI Team are opposed to recommending that the Department implement a 60-hour HFE this fall, several members support recommending that the Department implement a 60-hour duration HFE this fall, and two members (WAPA and AGFD) have abstained from making a recommendation.

PI Team representatives opposed to a 60-hour HFE include all seven Basin States (AZDWR, CA, CO, NM, UT, WY, NV), UCRC, and Reclamation. Those opposed to implementing a shorter duration HFE this fall indicated that the reduced duration did not sufficiently address all unacceptable adverse impacts to resources identified above (low pool elevation, Basin Fund status). They also expressed concern that potential benefits to the sediment resource would not outweigh potential adverse impacts. PI Team members again highlighted uncertainties regarding Glen Canyon Dam's ability to reliably and effectively deliver water and generate power when pool elevation drops below 3,525 ft.

Those that support or that do not oppose implementation of a 60-hour HFE include BIA, NPS, USGS, and USFWS. PI Team members in favor of an HFE cited the much-needed sediment resource benefits from a fall HFE and included this option as a potential compromise for the Leadership Team to consider that addresses both concerns expressed while still providing benefits to sediment (predicted +30% sandbar size in November 2021 relative to October 2021). They also noted that, in LTEMP, fall HFEs were predicted to be conducted frequently (~3 out of

every 4 years) and it has been three years since the most recent HFE (fall 2018) was triggered and implemented, such that this HFE would rebuild some of the beaches that were lost during this time. GMCRC noted that sandbar benefits from a 60-hour HFE would likely be greater in Marble Canyon, but would result in sandbar deposition throughout Marble Canyon and Grand Canyon. It was noted that maintaining camping beaches is important for the LTEMP sediment resource goal. Also noted is the potential for improved legacy of sandbar increases resulting from a fall HFE (predicted +20% sandbar size in April 2022 relative to October 2021) due to relatively low winter flows consistent with the 7.48 maf annual release pattern planned for Lake Powell in Water Year (WY) 2022. The GCMRC highlighted the learning value of implementing an HFE in November 2021 in that it would contribute to addressing, 1) the long-term question about the cumulative effect of multiple HFEs over the 20-year period of the LTEMP, and 2) questions about the potential combined effects of HFEs and riparian vegetation management for sand storage in aeolian dunefields covering archaeological sites. Although the best available science indicates a fall HFE may result in small benefits to undesired non-native species, these effects were highly uncertain (see rainbow trout section below) or the marginal impact was expected to be small (see brown trout section). Thus, expert opinion among fishery biologists was that a 60 hour fall HFE would not substantively increase risk to endangered and native fishes in Grand Canyon above the existing level of risk.

Scope of Assessment

Flow Experiments at Glen Canyon Dam are an important feature of the LTEMP ROD that will help determine specific dam operations that could be implemented to improve conditions, continue to meet the Grand Canyon Protection Act's (GCPA) requirements, and to minimize adverse impacts on the downstream natural, recreational, and cultural resources. Per the LTEMP ROD, the PI Team is charged with seeking a consensus technical recommendation regarding the assessment of resources at and below Glen Canyon Dam (see Section III for list of resources). Water management activities occurring elsewhere within the Colorado River Basin may be relevant to the Department's decision of whether to implement flow experiments, but are being addressed separately outside the scope of this technical assessment. For example, water managers, including many PI Team members, are concurrently working to evaluate the potential effects of drought on water resources throughout the Basin, including evaluations of options to minimize the risk of Lake Powell falling below the "Target Elevation" of 3,525 ft. under the Drought Response Operations Agreement. The LTEMP process for evaluating potential flow experiments requires assessment of the effects on resources, and provides an off-ramp in the event of unacceptable adverse impacts. If implemented, flow experiments cannot change the annual release volume from Glen Canyon Dam. The PI Team appreciates the opportunity to provide this report and anticipates that the Leadership Team may wish to address considerations that were outside of the scope of this technical assessment.

II. Introduction

The purpose of HFEs conducted in the context of the LTEMP ROD is to determine if sandbar building during HFEs exceeds sandbar erosion during periods between HFEs, such that sandbar size can be increased or maintained over the 20-year period of the LTEMP. This study supports the LTEMP Resource Goal for the sediment resource to "Increase and retain fine sediment

volume, area, and distribution in the Glen, Marble, and Grand Canyon reaches above the elevation of the average base flow for ecological, cultural, and recreational purposes” (DOI 2016b). Table 4 in Appendix B of the LTEMP ROD summarizes implementation criteria for LTEMP experiments, and an excerpt of the criteria for fall HFEs is provided below (**Table 1**).

Table 1. LTEMP Implementation Criteria for Fall High Flow Experiments

Experimental Treatment	Trigger ^a and Primary Objective	Replicates	Duration	Annual Implementation Considerations ^b	Long-Term Off-Ramp Conditions ^c	Action if Successful
<i>Sediment-Related Experiments (Cont.)</i>						
Fall HFE ≤96 hr up to 45,000 cfs in Oct. or Nov.	Trigger: Sufficient Paria River sediment input in fall accounting period (Jul.–Nov.) to achieve a positive sand mass balance in Marble Canyon with implementation of an HFE Objective: Rebuild sandbars	Implement in each year triggered, dependent on resource condition and response	≤96 hr	Potential short-term unacceptable impacts on resources listed in Section 1.3; unacceptable cumulative effects of sequential HFEs	This type of fall HFE is not effective in building sandbars; or long-term unacceptable adverse impacts on the resources listed in Section 1.3 are observed	Implement as adaptive treatment when triggered and existing resource conditions allow
Fall HFEs longer than 96-hr duration up to 45,000 cfs in Oct. or Nov.	Trigger: Sufficient Paria River sediment input in fall accounting period (Jul.–Nov.) to achieve a positive sand mass balance in Marble Canyon with implementation of an HFE longer than a 96-hr, up to 45,000-cfs flow Objective: Rebuild sandbars	Implement in each year triggered; limited to total of four tests in LTEMP period	Up to 250 hr depending on availability of sand duration of first test not to exceed 192 hr	Potential short-term unacceptable impacts on resources listed in Section 1.3; unacceptable cumulative effects of sequential HFEs	Extended-duration fall HFEs are not effective in building sandbars; resulting sandbars are no bigger than those created by shorter-duration HFEs; or long-term unacceptable adverse impacts on the resources listed in Section 1.3 are observed	Implement as adaptive treatment when triggered and existing resource conditions allow

- a Triggers will be modified as needed during the 20-year LTEMP period in an adaptive manner through processes including ESA consultation and based on the best available science utilizing the experimental framework for each alternative.
- b Annual determination by the DOI. Any implementation will consider resource condition assessments and resource concerns using the annual processes described in Sections 1.3 and 1.4.
- c Suspension of experiment if the DOI determines effects cannot be mitigated.

Source: 2016 LTEMP ROD, Appendix B, Table 4 – Implementation Criteria for Experimental Treatments of Alternative D

To date, five HFEs have been conducted under sand-enriched conditions since the HFE Protocol was initiated in 2012. Those HFEs occurred in November each year in 2012, 2013, 2014, 2016, and 2018. In each case, sandbar building results were consistent with the results from previous HFEs (Grams, 2019). All HFEs resulted in substantial deposition at all sandbar types (see Mueller and others, 2018 for description of sandbar types), with some evidence of cumulative gains in sandbar volume over multiple HFEs. While observations demonstrate that HFEs benefit campsites and cause temporary increases in campsite area, vegetation encroachment continues to cause progressive declines in campsite areas at some locations.

III. LTEMP Process for Implementing Experiments

The 2016 LTEMP ROD provides the framework for implementing flow-based experiments at Glen Canyon Dam when resource conditions warrant. The purpose of LTEMP experiments is to learn, through adaptive management, how to better protect, mitigate adverse effects to, and improve resources downstream of Glen Canyon Dam, while complying with relevant laws. Ongoing research and monitoring through the Glen Canyon Dam Adaptive Management

Program ensures the best science and data are available for making decisions related to flow experiments.

Under the LTEMP, the Department may conduct flow-based experiments (HFEs, Bug Flows, Trout Management Flows, and Low Summer Flows) at Glen Canyon Dam when resource conditions warrant and if it is determined that there will not be unacceptable adverse impacts on other resources.

Prior to implementation of any experiment, the relative effects of the experiment on the following resource areas will be evaluated and considered: (1) water quality and water delivery, (2) humpback chub, (3) sediment, (4) riparian ecosystems, (5) historic properties and traditional cultural properties, (6) Tribal concerns, (7) hydropower production and WAPA's assessment of the status of the Basin Fund, (8) the rainbow trout fishery, (9) recreation, and (10) other resources.

--P. B-8, Implementation Process for Experiments Under Alternative D, 2016 LTEMP ROD

The process for recommending experiments under the LTEMP, which has been used for past experiments and has been followed here, involves outreach to GCDAMP partners through regular meetings and additional notification to Tribes inviting consultation. The process also involves coordination with the PI Team to plan for the possible experiment, evaluate the status of resources, and make the technical recommendation of whether to conduct an experiment. The PI Team presents its recommendation to the Leadership Team, which makes a recommendation to The Department. The Secretary's Designee to the Adaptive Management Work Group (AMWG) is the chair of the Leadership Team and may make the decision for the Department regarding the experiment, as delegated by the Secretary of the Interior.

LTEMP HFE Protocol

As described in the LTEMP ROD, HFEs are experimental in nature and are designed to achieve a better understanding of whether, how, and when to incorporate high releases into future dam operations in a manner that maintains or improves beaches, sandbars, and associated habitat. The LTEMP HFE Protocol establishes a decision-making framework consisting of three components: (1) planning and budgeting, (2) modeling, and (3) decision and implementation.

Under the LTEMP, HFE releases are restricted to limited periods of the year when the highest volumes of sediment are most likely available for building sandbars. Sediment-triggered HFEs may be implemented in spring (March or April) or fall (October or November; **Figure 1**). Fall extended-duration HFEs range from greater than 96 hr to 250 hr. Spring and fall HFEs that are not extended-duration range from less than 1 hr to 96 hr. Proactive HFEs may be implemented in spring or early summer (April, May or June), and have a duration range up to 24 hr. HFE magnitudes range from 31,500 cfs to 45,000 cfs. The frequency of HFEs is determined by tributary sediment inputs, annual release volumes, resource conditions, and decision of the Department. Extended-duration fall HFEs are limited to a frequency of 4 times total in the 20-year LTEMP period.




Figure 1. Average monthly sand load from the Paria River (1923-2010) and Little Colorado River (1987-2010) showing the fall and spring HFE accounting periods and implementation windows (DOI 2016a).

HFE Sand Budget Model

The LTEMP HFE Protocol uses predictive models to make recommendations for the magnitude and duration of potential HFEs using real-time measurements and models of sand inflow from the Paria River and forecasted hydrologic data to determine whether suitable sediment and hydrology conditions exist for a high-flow experimental release.

A sand transport/budget model (Wright et al. 2010) was used to predict the mass of sand that would be transported by an HFE and to estimate if a potential HFE would transport more or less sand than had been delivered from the Paria River to the Colorado River during the fall accounting period (July 1 to November 30). Only HFE durations that resulted in a “positive sand balance” were considered. Output of the modeling runs provides the initial recommendation for the magnitude and duration of the HFE. However, because modeling only considers a simple range of possible HFE peak magnitudes and durations, the Protocol includes a review of the model output that may modify the recommended HFE to benefit relevant resources.

In addition to reviewing the sand budget model output, the PI Team assessed the status of the LTEMP resources and the potential effect of an HFE on these resources in making the recommendation described here.

IV. Hydrograph Alternatives

HFE Sand Budget Model Results

Throughout the summer and fall, and in accordance with the Protocol, Reclamation regularly updated its modeling estimates based on cumulative sediment inputs to determine the largest HFE that resulted in a positive sand balance in Marble Canyon. The modeled HFE shape was based on past years’ input from scientists at GCMRC and designed to meet the twin objectives of providing the greatest resource benefit and developing scientific information that will help better

inform future decision making. Hydrology inputs were provided as hourly Glen Canyon Dam releases (historic and future) for the accounting period (July 1 – November 30).

The final September 27, 2021 model results predicted there was sufficient sediment for implementation of a 192-hr extended duration HFE—the longest duration allowed for the first implementation of an extended duration HFE—under the LTEMP HFE Protocol. The final cumulative sand load estimates for the lower and upper bounds were, respectively, 982,000 and 1,471,000 metric tons. This model run used the conservative lower bound estimate for Paria River sand input and estimated 136,000 metric tons would remain on November 30, 2021 following a 192-hour extended duration HFE and at the end of the accounting period.

Hydrograph Alternatives

In response to resource impact concerns, the PI Team proposed two hydrograph alternatives that had potential to reduce negative impacts while still meeting objectives for sediment benefits and learning (**Figure 3**). These alternatives reduced the duration of the experiment (see **Table 2**, **Table 3**).




Figure 2. Sand Budget Model Results for Fall 2021 Accounting Window, 192-hour Extended Duration HFE




Figure 3. November 2021 Schedule of Releases for Alternative Fall HFE Hydrographs

Table 2. Summary of Water Releases, Pool Elevation, and Basin Fund Impacts for Alternative Fall HFE Hydrographs

Fall HFE Hydrograph Alternatives	Water Release		Pool Elevation ²			Relative Impact to Basin Fund
	November Total Release Volume (kaf)	Bypass Volume (kaf)	End of November Elevation (ft)	Number of Additional Days below 3525 ft	WY 2022 Minimum (ft) Difference from No Action	
No HFE	500	0	3540.70	0	3518.58 --	\$0
60-hr HFE (MAM) ³	634	82	3538.85	15	3517.89 (0.69)	\$(1.30M)
60-hr HFE (DJF) ³	634	82	3538.85	0	--	\$(3.04M)
96-hr HFE (MAM)	710	126	3537.79	20	3517.50 (1.08)	\$(1.94M)
96-hr HFE (DJF)	710	126	3537.79	0	3518.59 0.01	\$(4.66M)
192-hr HFE (MAM)	915	245	3534.92	34	3515.42 (3.16)	\$(3.70M)

1 – Peak capacity for all alternatives is 33,900 cfs.

2 – Assumes most probable hydrology as indicated in the September 24 Month Study.

3 – Refers to the months in which less water would be released in order to balance a November HFE: March, April, May (MAM) or December, January, February (DJF).

Table 3. Summary of Learning and Effectiveness for Alternative Fall HFE Hydrographs

Fall HFE Hydrograph Alternatives	Learning Relative to Hypotheses (ranked most to least)	Effectiveness Relative to Objective (ranked most to least)	Sediment Benefit ¹
No HFE	4 – Very Low ²	4 – None	0
60-hr HFE (MAM)	3 – Moderate (short duration is not consistent w/ LTEMP experimental design; contributes to Western Grand Canyon sediment study) ²	3 - Moderate (reduced peak/low duration will build sandbars, but will likely be limited in downstream reaches)	60%
96-hr HFE (MAM)	2 – Moderate/High (regular HFEs are a critical part of the LTEMP experimental design; contributes to western Grand Canyon sediment study) ²	2 - Moderate/high (reduced peak/long duration expected to build sandbars throughout most of Marble and Grand Canyon)	70%
96-hr HFE (DJF)	See above ²	See above	See above
192-hr HFE (MAM)	1 - High (regular HFEs are a critical part of the LTEMP experimental design; learning associated with the extended duration is high despite low peak discharge, because this represents first extended duration HFE)	1 - High (extended duration hypothesized to result in greater sandbar building in downstream reaches)	105%

1 – Sediment benefit is described as "bar building relative to a base HFE of 96-hours at 44,000 cfs"

2 – The unique conditions of high sediment input and low winter flows will be valuable to inform sediment accounting window discussions under all scenarios. Low winter flows will allow substantial sand retention in Marble Canyon through at least June 30, 2022.

Experimental Design and Description - 60-Hour Duration

Potential 60-hour HFE Hydrograph:

- Ramp-up from base releases at 4,000 cfs/hr at approximately 7:00 am on Monday, November 1, 2021 (all times Mountain Standard Time) until reaching powerplant capacity (~18,900 cfs)
- Open first bypass tube at 10:00 am on November 1
- Ramp-up from powerplant capacity to full bypass (~33,900 cfs) at one full bypass tube (~3,750 cfs) per hour in 4 hrs
- Stay at peak release (~33,900 cfs) for 60 hrs
- Ramp-down from peak release to base releases at beginning at half bypass of 1,875 cfs/hr until reaching powerplant capacity and then decreasing at 2,500 cfs/hr

These recommendations result in the following release schedule at Glen Canyon Dam (also, **Figure 4**):

- Begin ramp-up from 6,500 cfs at 7:00 am on November 1 (Monday)
- Reach powerplant capacity at approximately 10:00 am on November 1
- Open bypass tubes at approximately 10:00 am on November 1
- Reach full bypass at 2:00 pm on November 1
- Begin ramp-down from bypass at 2:00 am on November 4 (Thursday)
- Complete HFE (back to 9,000 cfs) at 1:00 pm on November 4 (Thursday)




Figure 4. November 2021 Glen Canyon Dam hourly releases for Potential 60-hr HFE

V. Monitoring Plan

GCMRC developed a science plan for the LTEMP that describes a program of monitoring and research activities that support ongoing information needs associated with implementation of the LTEMP and associated experiments like HFEs (VanderKooi et al. 2017). This approach relies on water quality, sediment, aquatic biology, and other resource monitoring and research projects funded in the GCDAMP Fiscal Year (FY) 2021-2023 TWP (Reclamation and GCMRC TWP, U.S. Department of the Interior, 2020). These projects will inform the effect of future HFEs on the downstream resources of Glen, Marble, and Grand Canyons. Projects from the TWP specific to monitoring HFEs are further discussed below.

Project A, *Streamflow, Water Quality, and Sediment Transport and Budgeting* in the Colorado River Ecosystem, and Project B, *Sandbar and Sediment Storage Monitoring and Research*, are essential components to implementation of the HFE Protocol under LTEMP because the protocol calls for high flow releases from Glen Canyon Dam whenever a specified minimum amount of fine sediment delivered from the Paria River is exceeded. Under Project A, the measurements needed to trigger HFEs are collected. Project B supports the direct measurements of the volume of fine sediment, especially sand, that is stored on the bed of the Colorado River, in its eddies, or at higher elevation along the river's banks; these measurements allow assessment of the effectiveness of the HFE Protocol. A substantial accomplishment of these programs in previous work plans was the development of web-based interfaces to serve sediment transport and water quality data, calculate fine sediment mass balances (see https://www.gcmrc.gov/discharge_qw_sediment/), and to serve photographs of approximately 50 sandbars located from Lees Ferry to Diamond Creek (see <https://www.usgs.gov/apps/sandbar/>). The latter data allow stakeholders to evaluate the effects of controlled floods implemented under the HFE Protocol.

As described in the HFE Protocol EA (U.S. Department of the Interior 2011) and the LTEMP FEIS (U.S. Department of the Interior 2016a), the potential HFE planned for fall 2021 would not be an isolated event, but a component of a longer-term experiment to restore and maintain sandbars with multiple high flows over a period of several years. The monitoring data that are needed to assess the outcome of this multi-year experiment include annual sandbar monitoring at selected long-term monitoring sites, periodic monitoring of changes in sand storage in the river channel, and measurements of sandbar size at more than 1,000 sites based on aerial photographs that are collected periodically. These activities are described in detail in the Reclamation and GCMRC TWP (U.S. Department of the Interior 2020). It is also important, however, to evaluate the sandbar building response of each high flow to assess whether the sandbar building objectives are being achieved incrementally. This evaluation will be based on sites that are monitored by remotely deployed digital cameras and repeat topographic surveys of sites that will occur in fall 2022.

GCMRC scientists have installed digital cameras that capture 5 images every day at 43 sandbar monitoring sites throughout Marble and Grand Canyons between Lees Ferry and Diamond Creek. The images acquired by these cameras will be used to evaluate both the magnitude and spatial distribution of sandbar building caused by the HFE (Grams et al. 2018). They will also be used to assess the rate of post-HFE sandbar erosion. Because the remote cameras are monitoring

the same sites that are monitored by the annual surveys and the same sites that were monitored during the previous high flows, it will be possible to evaluate sandbar-building effectiveness of the planned HFE relative to the previous HFEs.

Project D will continue monitoring for changes in sediment storage in dunefields covering archaeological sites. An HFE has not occurred since the NPS LTEMP vegetation management project was implemented to open up sandbar campsites and increase aeolian transport to archaeological sites. Thus, we have data for scenarios 1-3 but not #4 below:

- o 1) Year(s) without vegetation removal and without HFE,
- o 2) Year(s) without vegetation removal but with HFE,
- o 3) Year(s) with vegetation removal but without HFE,
- o 4) Year with vegetation removal and HFE.

An HFE implemented in 2021 would provide GCMRC and NPS the opportunity under their collaborative LTEMP vegetation management project to address a critical uncertainty (i.e., scenario #4 above) about the potential combined effects of HFEs and riparian vegetation management.

GCMRC will also collect data on water quality (including nutrients), native and nonnative fishes, aquatic invertebrates, riparian plants, and other resources, as described in the Reclamation and GCMRC TWP (DOI 2020). Project N, *Hydropower Monitoring and Research*, is focused on conducting monitoring and research of proposed experiments in the LTEMP ROD and consider impacts on hydropower and energy as part of the experimental design. Project G, *Humpback Chub Population Dynamics throughout the Colorado River ecosystem*, continues long-term monitoring in Grand Canyon and will provide information about potential response of Humpback chub to a fall HFE.

GCMRC will also collect additional data in support of project O.2, which is a study of channel response to flow pulses in the western Grand Canyon. The HFE will provide the opportunity to measure bed response to the larger 33,900 cfs HFE than was measured in the 2021 spring disturbance flow. The required data are repeat measurements of channel bathymetry and bank topography for 3-km long study reach. The objectives and budget are described in Project B.6.5 in the GCDAMP FY 2021-2023 TWP.

VI. Assessment of Resources

In coordination with the GCMRC, the PI Team completed an assessment of key resources that may be impacted or affected by an HFE. This assessment is based on the current condition of resources and on findings and observations from fall HFEs conducted in 2012, 2013, 2014, 2016, and 2018. Key resources were evaluated relative to the timing, duration, and magnitude of the potential fall 2021 HFE using the best available science: November 1, 2021 start date, 60-hour duration, and 33,900 cfs peak magnitude. This section summarizes the assessment of resources and expected effects of the potential HFE.

Archaeological and Cultural Resources

Reclamation and NPS (DOI 2016b) determined that HFEs, as identified in the LTEMP, could, through multiple experiments, potentially affect historic properties and the effect would be

adverse per 36 CFR 800.5(2)(iv). The agencies also found that adverse effects to sacred sites could result from the HFEs, primarily from limitation of access of tribes to sacred sites during the period of HFE releases. Reclamation, as lead federal agency for National Historic Preservation Act section 106 compliance, completed the LTEMP Programmatic Agreement (PA, Reclamation 2017) with affected tribes and other parties to address these effects. Effects of HFEs to cultural resources are primarily from erosion and redistribution of sediment. Inundation can adversely affect sites through erosion, but deposition may help protect sites by providing sources of sand that can bury historic properties via aeolian transport (Reclamation 2011, DOI 2016b, East et al. 2016). HFEs also may affect access of tribes to historic properties and alter visitation patterns to historic properties (Reclamation 2011, DOI 2016b).

The PA incorporates, by reference and specified in Appendix D, a commitment to the stipulations identified in previous compliance agreements, most notably the Memorandum of Agreement for the 2012 High Flow Experiment. Reclamation, as lead federal agency, notified Tribes and consulting parties on September 20, 2021 identifying the potential for a fall HFE in 2021, in conformance with the stipulations in the PA (and previous MOA).

GCMRC monitoring has shown that recent HFEs have eroded terraces that contain archaeological sites in Glen Canyon National Recreation Area (GLCA; East et al. 2016). HFEs also rebuild or maintain sandbars that provide sand to resupply aeolian dunefields containing archaeological sites throughout Marble and Grand Canyons (Sankey et al. 2018). Aeolian dunefields were resupplied with sand from HFE deposits in half of the instances monitored after the 2012, 2013, 2014, and 2016 HFEs (Sankey et al. 2018). There is also evidence for cumulative effects of sediment resupply of dunefields when annual HFEs are conducted consistently in consecutive years (Sankey et al. 2018). No adverse effects to historic properties were identified from the 2012, 2013, or 2014 HFEs. Results from monitoring following the 2016 and 2018 HFE showed that several archaeological sites have transitioned from net-erosion to net-deposition dominated topographic changes in association with the higher frequency of HFEs during the time period of the current HFE protocol. Additionally, the NPS LTEMP vegetation management project was implemented beginning in 2019 to open up sandbar campsites and increase aeolian transport to archaeological sites. An HFE has not occurred since the vegetation management experiments were implemented. Thus, questions about the potential combined effects of HFEs and riparian vegetation management for sand storage in aeolian dunefields covering archaeological sites have not yet been addressed. **A fall HFE in 2021 would not pose risks to archaeological and cultural resources that were not previously analyzed in the LTEMP.**

Natural Processes (Aquatic Food Base)

Flood disturbance is a critical natural process in streams and rivers (Poff and others 1997). By disrupting ecosystem structure and altering the availability of substrates and resources, flood disturbance can help maintain native biological diversity (Carlisle and others 2017). Disturbance magnitude, for example the extent of drying at low flow or the proportion of the bed that is mobilized at high flows, can influence ecosystem outcomes by determining the extent of biomass loss and the quantity of newly scoured habitat patches available for recolonization by fast-growing algae and aquatic insects. Disturbance frequency and timing (e.g., spring vs. fall) can also influence the rate and trajectory of ecosystem recovery from disturbance. The life cycles of

many species of native algae, insects and fish are directly tied to flood disturbances, and alterations to river flood regimes can adversely affect ecosystem health. In fact, a national synthesis of flow and biological data from over 700 streams and rivers in the lower 48 states found that intact and healthy communities of native aquatic invertebrates and fish were most often present where flood disturbance still occurred, and where flood timing was seasonally appropriate (i.e., similar to the natural condition; Carlisle and others 2017). Although the Colorado River in Grand Canyon could not be included in this 2017 synthesis owing to the absence of pre-dam ecological data, the mechanisms linking periodic flow disturbance to stream ecosystem health were evaluated in a wide variety of streams and regions. It is therefore reasonable to predict that similar mechanisms linking appropriately timed flow disturbance to ecosystem health also operate in the Colorado River.

The pre-dam Colorado River was characterized by spring snow-melt floods that often exceeded 100,000 ft³/s and typically peaked in late June, followed by flash flood flows during the summer monsoon season, and extensive low base flows from the fall through early spring (Topping and others 2003). This seasonally variable flow regime was an important driver of natural processes in the Colorado River, and the unique fish species that evolved here were adapted to frequent flow disturbances. For example, the small eyes and tiny embedded scales that are common to several native fish are thought to be adaptations to the sediment laden floods that scoured the Grand Canyon annually. In the pre-dam river, turbidity was always high (Voichick and others 2016), suggesting that algae growth was likely restricted to the river's edge or shallow cobble habitats; detritus and leaf litter are thought to have been primary sources of energy fueling food webs (Blinn and Cole 1991).

Regulation of the Colorado River by Glen Canyon Dam in 1963 eliminated the annual spring high flow disturbances, it substantially increased base flow thereby eliminating periods of low flows, and hourly variation in discharge increased substantially owing to hydropower production (Topping and others 2003, Kennedy and others 2016). In addition to changing the river's flow regime, Glen Canyon Dam also changed other aspects of the physical template, particularly temperature, sediment, and nutrient regimes (Dibble and others 2021, Topping and others 2000, Yackulic and others 2018). These changes to the physical template led to dramatic changes in the natural processes that sustain river food webs. For example, Colorado River food webs are now primarily built upon algae production owing to clear water (Stevens and others 1997, Cross and others 2013). Many types of aquatic insects have been extirpated from Grand Canyon owing to multiple stressors including cold water temperatures and hourly fluctuations in discharge that leads to acute mortality of aquatic insect eggs laid along constantly changing shorelines (Kennedy and others 2016). Three species of native fish have also been extirpated from Grand Canyon owing to cold water, predation by non-native fish, and dams that block migration routes (Minckley 1991).

Meanwhile, many species of non-native invertebrates (e.g., *Gammarus lacustris*, New Zealand mud snails, quagga mussels) and fish (e.g., rainbow trout, brown trout) have become established throughout the Grand Canyon segment because they are well-suited to this new physical template and altered flow regimes. Predation by non-native fish species is a constant and potentially growing threat to many native fish species that remain in Grand Canyon. Additionally, simplified food webs that are inherently unstable and overall low food base

production arising from nutrient limitation have been shown to limit native and desired non-native fish populations in some reaches and years (Cross and others 2013, Korman and others 2021). In spite of these changes to the ecosystem and the natural processes that support food webs, native fish populations in Grand Canyon have been relatively stable over the past two decades compared to other segments of the Colorado River (Yackulic and others 2014, Healy and others 2020, Dibble and others 2021).

The LTEMP seeks to enhance key resources through experimentation and management of both flow and non-flow actions. High Flow Experiments are the principal type of flow disturbance evaluated as part of LTEMP. Because the annual snowmelt flood of the Colorado River was in spring and early summer, it has long been hypothesized that spring HFEs would be more likely to benefit natural processes of the river compared to fall HFEs (Kennedy and Ralston, 2011). Consistent with these predictions, monitoring of fall HFEs in 2012, 2013, 2014, 2016, and 2018 indicate these flow disturbances have neutral-to-negative effects on algae production, aquatic insect abundance and diversity, and other natural processes (Kennedy and Muehlbauer 2019). Unfortunately, these tools for monitoring natural processes were not in place in 2008 the last time a spring HFE was tested, so the role of spring HFEs in enhancing natural processes remains unclear. Nonetheless, monitoring of natural processes occurred in March 2021 during the spring disturbance flow test at Glen Canyon Dam, and future analysis of these monitoring data may help shed light on whether flow disturbance in spring benefits natural processes. **A 2021 fall HFE would not pose unacceptable risks to natural processes that were not previously considered or analyzed in the LTEMP.**

Humpback chub

The adult humpback chub population in the Little Colorado River aggregation is currently above the Tier-1 threshold of 9,000 adults identified in the Biological Opinion for the LTEMP (GCMRC unpublished data, USFWS unpublished data). Past HFEs have had no measurable direct effects, positive or negative, on humpback chub or other native fish. Their populations in the Little Colorado River aggregation remained relatively stable over the decade from 2009-2019, a period that included HFEs in 2012, 2013, 2014, and 2016 and increased water temperatures (Kennedy and Ralston 2011, GCMRC unpublished data). However, populations are now declining in the Little Colorado River aggregation and are expected to continue to decline over the next few years. The three-year average (2018-2020) of large sub-adult humpback chub in the Colorado River mainstem from river mile 63.45 to 65.2 (juvenile chub monitoring reach) was estimated at 600 fish, which is below the 810 fish required to prevent initiation of a Tier 1 fish management action trigger. Fish biologists are working to identify factors that may have led to poor recruitment, assess the current outlook for humpback chub populations in the Grand Canyon, and evaluate early intervention actions that can be taken to reverse the decline.

HFEs may indirectly affect humpback chub through increases in rainbow trout or brown trout populations, which are discussed below in those respective sections. Based on provisional unpublished data, humpback chub were not directly affected by the 2012, 2013, 2014, 2016, or 2018 HFEs, with adult populations appearing stable over the period of these HFEs and juvenile populations fluctuating in response to variable recruitment in the Little Colorado River. For example, there was poor production of juveniles this year in the Little Colorado River in agreement with past findings that in years lacking winter/spring runoff humpback chub

recruitment is depressed (Van Haverbeke et al. 2013). **Based on the current population status of humpback chub and the unclear relationship of fall HFEs to humpback chub, a fall 2021 HFE does not pose unacceptable risks to humpback chub that were not previously considered in the LTEMP.**

Hydropower and Energy

WAPA has firm electric power contracts and will continue to meet these contract obligations either with generation from Colorado River Storage Project powerplants or from purchases from the wholesale electrical market. Low-volume releases from Glen Canyon Dam both pre- and post-HFE requires extra electrical purchases to meet WAPA’s contract obligations. Conversely, during the HFE, high volume releases from Glen Canyon Dam requires the energy be sold to the open market. WAPA estimated the financial cost of implementing a 2021 fall HFE is between \$1.30 million and \$4.66 million, depending on duration and which months water is moved from to provide volume for the HFE (see **Table 4**). For comparison with past HFEs, WAPA determined the costs at \$1.1 million for 2012, \$2.6 million for 2013, \$2.1 million for 2014, \$1.2 million for 2016, and \$1.3 million for 2018.

Table 4. WAPA’s Estimate of the Impact of Fall HFE Alternatives on the Basin Fund

Fall HFE Hydrograph Alternatives	November Total Release Volume (kaf)	Bypass Volume (kaf)	Months water moved from for HFE	Volume moved (1000’s AF)	Relative Impact to Basin Fund
No HFE	500	0	--	--	\$0
60-hr HFE	634	82	Mar	46	\$(1.30M)
			Apr	44	
			May	44	
60-hr HFE	634	82	Dec	44	\$(3.04M)
			Jan	46	
			Feb	44	
96-hr HFE	710	126	Mar	72	\$(1.94M)
			Apr	69	
			May	69	
96-hr HFE	710	126	Dec	69	\$(4.66M)
			Jan	72	
			Feb	69	
192-hr HFE	915	245	Mar	141	\$(3.70M)
			Apr	137	
			May	137	

Water releases from Glen Canyon Dam during the HFE may be affected by disturbances of the electrical system. Responses to these disturbances are required by Reclamation and WAPA under law, contracts, and other agreements. Regulation and contingency reserves are the two types of assistance provided by the CRSP system for electrical disturbances, and both are managed by WAPA’s Western Area Colorado-Missouri (WACM) Balancing Authority. Regulation is used to respond to frequency deviations on the electrical system. Glen Canyon Dam is the only CRSP powerplant capable of the immediate responses required for regulation and can increase or decrease releases by as much as ±1,280 cfs (40 mw) for up to 1 hour and 59

minutes. WAPA would move reserves to an alternate CRSP unit for a fall 2021 HFE after determining this will not likely result in additional cost.

WAPA estimates that the balance of the Upper Colorado River Basin Fund (Basin Fund) will be about \$87 million at the end of fiscal year 2021. This is considerably below WAPA's target for an end of year balance. WAPA further projects significant reductions in the Basin Fund balance over the next two years primarily due to reduced reservoir levels (hydraulic head), low annual releases from Glen Canyon Dam (7.48 maf in water years 2022 and 2023) and increases in purchase power prices and amount. Without significant mitigation efforts, the Basin Fund would not be able to sustain the project. Mitigation actions in progress to reduce impacts to the Basin Fund include deferrals of O&M expenses and capital projects, and a new public process for a potential rate adjustment aimed at avoiding a cost recovery charge and avoiding significant projected purchase power costs in the near future. Regardless of a possible new rate, WAPA intends to determine the cost of HFEs and other experiments at Glen Canyon Dam as has been done in the past and will account for them as a constructive return. Given the extraordinary measures taking place to maintain the Basin Fund by deferring millions of dollars in expenditures, and a proposed new two-year rate, WAPA is very concerned with actions that would further exacerbate the Basin Fund balance. **Therefore, the current status of the Basin Fund is not at a level sufficient to support implementation of a Fall 2021 HFE without considerable risk of adverse impacts.**

Other Native Fishes

A small reproducing population of endangered razorback sucker occurs downstream in Lake Mead, and recent monitoring data indicate that razorback sucker occupy and spawn in western Grand Canyon (Kegerries et al. 2017). In 2012, a single adult was captured near Spencer Canyon (Bunch et al. 2012) and several other sonic-tagged individuals were detected in the same relative area (Kegerries et al. 2017). Razorback suckers have been captured in small numbers in this same area in subsequent years including two adults, one untagged and one sonic tagged, in 2013, one sonic-tagged adult in 2016, and one untagged adult in 2018 (AGFD unpublished data). Razorback sucker larvae were captured as far upstream as river mile 127.3 in 2019; however, numbers of larvae found during standardized monitoring have steadily declined since 2014 from 462 to 0 in 2020 and 2021 (Kegerries et al. 2021, S. Platania, ASIR, Inc., personal communication).

Regardless of declines in observed larvae, changes in flows due to a fall HFE are unlikely to have a substantial effect to razorback suckers, since life stages that might be sensitive to higher flows (e.g., spawning adults, larval fish) are not present in the fall months. In recent years, native fish have increased in abundance (or remained stable) and distribution in western Grand Canyon, with large numbers of juvenile humpback chub and flannelmouth sucker present (Kegerries et al. 2021, Van Haverbeke et al. 2017, Rogowski et al. 2018).

Invasive salmonids in Marble Canyon may benefit from a fall HFE through reduced turbidity after the HFE and enhanced growth and survival (Korman and others 2021), which could lead to indirect effects on native fishes through predation. Nonetheless, this potential indirect effect depends on factors driving turbidity that are less certain, such as additional tributary sediment

inputs. **Thus, no unacceptable negative response would be expected among native fishes to a fall HFE this year, based on current monitoring results and previous HFEs.**

Recreational Experience

The majority of recreational users along the river in both GLCA and GRCA access the river by boat. Upriver trips from Lees Ferry are primarily related to day use activities such as angling, motorized and non-motorized watercraft, and camping. GLCA has seen a significant increase in non-motorized watercraft (kayaks, canoes, paddleboards) since 2019 and the commercial outfitters that use motorized watercraft to transport these boaters upstream (i.e., backhaul services) have also increased significantly during the fall months. Recreational use is now throughout the week with higher numbers on the weekends as opposed to previously. If a HFE is implemented, press, website, and social media posts as well as direct contacts with the commercial operators will need to occur to inform the potential recreationists. Day use visitors also use Paria Beach for picnicking and shoreline recreation but most use in this area is during the summer and early fall, with limited use in the colder, late fall into winter months. Consequently, little to no impact to shoreline users would be anticipated from the HFE.

Both commercial and private angling trips for rainbow trout and now for brown trout under the Incentivized Harvest program also occur on all days of the week and may be increasing in early November as brown trout start appearing in and around their fall spawning beds. During the peak flow of the HFE, there would be a direct impact to fishing as it would produce flows large enough to impede fishing activity and may also affect foraging behavior of trout immediately following the HFE, reducing catch rates.

White-water boating in Grand Canyon is a year-round recreational experience, and all Grand Canyon river users with permits for use of the river during the HFE could be affected by changing flow patterns. Effects would primarily be related to safety considerations, covered in Section VII of this report. Day raft trips from Glen Canyon Dam to Lees Ferry, conducted under contract by Wilderness River Adventures (WRA), cannot operate during HFEs because flow into the Colorado River uses the bypass tubes at Glen Canyon Dam near the launch point for these trips. This commercial operator would be notified of the HFE dates and may be closing their limited 2021 operations around the time of the HFE.

Impacts to recreational experiences associated with the HFE would be both short- and long term in GRCA. Although the HFE is proposed to take place on the first day when only non-commercial launches are permitted, there would be many commercial rowing trips on the water from launches during the two weeks prior to the HFE (403 people on a total of 35 river trips). Some private trips may choose to cancel rather than be on the water during the high flows. Both commercial and private trips may alter their itineraries to avoid rafting if they are near rapids, which become more dangerous at high flows. The stage elevation changes (5 – 10 feet depending on the camp, per Magirl's models) at the start and end of the flows may create problems in camp and make some campable areas unusable. Medium- to long term impacts to recreational experience would be expected to be minimal as flows of this magnitude are unlikely to alter the rapids. The HFE should benefit recreational users through improvements to campable area and vegetation in the near shore habitat.

Sediment

During the period between the last HFE in November 2018 and July 1, 2021, roughly ½ million metric tons of sand have been eroded from Marble and Grand canyons. Large magnitudes of sand erosion occurred in Upper Marble Canyon, Eastern Grand Canyon, and East-Central Grand Canyon, with sand deposition occurring in Lower Marble Canyon and West-Central Grand Canyon. The changes in sand mass were as follows: Upper Marble Canyon (RM 0–30) - 580,000 ±130,000 metric tons; Lower Marble Canyon (RM 30–61) +120,000 ±110,000 metric tons; Eastern Grand Canyon (RM 61–87) -310,000 ±260,000 metric tons; East-Central Grand Canyon (RM 87–166) -200,000 ±240,000 metric tons; West-Central Grand Canyon (RM 166–225) +450,000 ±230,000 metric tons. Importantly, whereas the sand eroded from Upper Marble Canyon, Eastern Grand Canyon, and East-Central Grand Canyon likely came, in part, from high-elevation sandbars, the sand that was deposited in Lower Marble Canyon and West-Central Grand Canyon was deposited at lower elevations in the channel and eddies but not in high elevation sandbars. The most recent sandbar monitoring data show erosion of sandbar since the fall 2018 HFE and photographic evidence from summer 2021 shows extensive gully erosion at many sandbars in Marble Canyon and Grand Canyon caused by summer 2021 thunderstorms.

See discussion in Section IV for current sediment conditions relative to the HFE Protocol. Five HFEs have been conducted under sand-enriched conditions since the HFE Protocol was initiated in 2012. Those HFEs occurred in November of 2012, 2013, 2014, 2016, and 2018. In each case, sandbar building results were generally consistent with the results from previous sand-enriched HFEs as described by Schmidt and Grams (2011). All prior HFEs resulted in substantial deposition at all sandbar types (see Mueller et al. 2018 for description of sandbar types) followed by erosion of about half the new deposits within six months. Response immediately after the 2018 HFE based on digital camera images of sandbars from Lees Ferry to Diamond Creek indicated that there was a substantial gain (deposition) for 28 sandbars (66% of sites), no substantial change for nine sandbars (22% of sites), and substantial loss (erosion) for five sandbars (12% of sites) (Grams 2019). The HFE deposits typically begin eroding immediately following each HFE and the bulk of the newly deposited sand persists for approximately six to 12 months. Annual topographic surveys of sandbars were conducted between September 29 and October 16, 2020. Data from these surveys indicate there has been some net increase in the size of reattachment sandbars since the beginning of the HFE protocol in 2012. The size of other types (Mueller et al. 2018) of sandbars has fluctuated, with no significant net increase or decrease. Thus, repeated HFEs under the protocol have caused some cumulative increases in the size of some sandbars. Deposition of sand during HFEs has caused temporary increases in campsite area; however, there has been a net long-term decline in campsite area caused mostly by vegetation encroachment (Hadley et al. 2018a, Hadley et al. 2018b). Anecdotal reports from river guides indicate that hillslope runoff from summer rainstorms has caused substantial erosion at many sandbars during summer 2021. These eroded sandbars will not rebuild without an HFE.

Sandbars provide sand to resupply dunefields via aeolian transport throughout Marble and Grand Canyons (Sankey et al. 2018). Aeolian dunefields were resupplied with sand from HFE deposits in half of the instances monitored after the 2012, 2013, 2014, and 2016 HFEs (Sankey et al. 2018). There is also evidence for cumulative effects of sediment resupply of dunefields when annual HFEs are conducted consistently in consecutive years (Sankey et al. 2018). In 2019, under the LTEMP vegetation management project, NPS began experimentally removing riparian

vegetation that creates a barrier for aeolian transport of sand from sandbar campsites to dunefield archaeological sites. An HFE has not occurred since the vegetation management experiments were implemented. Thus, questions about the potential combined effects of HFEs and riparian vegetation management for sand storage in aeolian dunefields covering archaeological sites have not yet been addressed.

The aggregate sand mass-balance conditions since inception of the HFE Protocol (i.e., for the period between July 1, 2012, and September 1, 2021) for the different segments of the Colorado River in Marble Canyon (from https://www.gcmrc.gov/discharge_qw_sediment/reaches/GCDAMP) are:

- Upper Marble Canyon: +1.90 million metric tons (the range of this measurement is between -0.88 and +4.60 million metric tons)
- Lower Marble Canyon: +1.20 million metric tons (the range of this measurement is between +0.30 and +2.20 million metric tons)

Thus, there was likely more sand in the Colorado River corridor in Marble Canyon on September 1, 2021, than on July 1, 2012, when the HFE Protocol was first implemented. **The potential fall HFE does not pose risks to the sediment conditions that were not previously analyzed in the LTEMP and will benefit most sediment resources.**

Tribal Resources

All resources in the canyon are of importance to Tribes, thus all resources are tribal resources. As such, careful consideration of the potential effects of an HFE on all resources has been considered. In addition, the taking of life in the canyon is a serious concern expressed by Tribal partners. The potential HFE would not be expected to directly or indirectly result in increased taking of life in the canyon, either during the experiment or in the future as a result of the experiment. **The potential experiment would not be expected to cause unacceptable adverse impacts to tribal resources.**

Rainbow Trout Fishery

The NPS Comprehensive Fisheries Management Plan completed in cooperation and coordination with AGFD (NPS 2013), and the AGFD Management Plan establish objectives for the rainbow trout fishery at Lees Ferry. Two of those objectives are to maintain angler catch rates of ≥ 1 rainbow trout per hour, and fish condition ≥ 1 in the summer. After a population crash of rainbow trout in the winter of 2015-2016, the population started rebounding with increased young of the year production in 2017-2018, however that did not result in an increase of catchable fish, and overall abundance has been decreasing since 2018.

Overall, angler catch rates for rainbow trout have decreased from 2018 to 2020. Results from 2020 creel surveys for boat anglers was 0.639 fish/hr (CI 0.588, 0.689), and preliminary data from 2021 (January - August) revealed a similar result of 0.584 fish/hr [0.520, 0.648]. In 2020, walk-in anglers had a catch per unit effort of 0.251 fish/hr [0.191, 0.310], and preliminary data

for 2021 was estimated as 0.297 fish/hr [0.0615, 0.419]. These values are below the targeted minimum catch rate of one fish/hr identified within AGFD's Management Plan.

Fish condition of rainbow trout has been shown to be negatively correlated with abundance. There is a limited food base in Glen Canyon for sub-adult and adult fish, and as fish abundance increases, food becomes limiting and fish condition begins to decline. In 2020 and 2021, the relative condition of rainbow trout accessible to anglers (> 6 inches) was greater than one indicating healthy fish (AGFD unpublished data).

The Natal Origins and Trout Recruitment and Growth Dynamics projects led by GCMRC to more closely examine rainbow trout have provided data to assess the effect of fall HFEs on rainbow trout abundance and vital rates in Lees Ferry and at the Little Colorado River (LCR). Analysis of data from fall HFEs conducted in 2004, 2012, 2013, 2014, 2016, and 2018 indicate there are no large, consistent effects of fall HFEs on the rainbow trout fishery in Lees Ferry. Rainbow trout growth has been high in years both with and without HFEs. Additionally, modeling indicates other factors besides HFEs, including fish density/competition and reservoir effects on nutrient dynamics and food base production are more important determinants of the health and abundance of the rainbow trout population in Lees Ferry (Korman et al. 2020). Further, there is no direct link between fall HFEs and emigration of rainbow trout out of Lees Ferry. **As such, a fall 2021 HFE would not be expected to have negative impacts on the rainbow trout fishery aside from the temporary and direct effect of the HFE on angling access that are described in the recreation section.**

Downstream near the LCR, fall HFEs may temporarily lead to higher growth, survival, and reproduction of rainbow trout owing to a period of relatively low turbidity after the HFE that benefits these visual sight-feeding fish (Korman et al. 2020). Conducting an HFE in fall 2021 may lead to slightly lower turbidity at the LCR over the winter compared to no-HFE, however the marginal effect of conducting an HFE on overwinter turbidity is somewhat uncertain owing to the absence of a quantitative model of turbidity and the unknown amount of moisture this winter. Analysis of historic turbidity data demonstrates that tributary flooding has an overriding influence on turbidity over the winter (Voichick and Topping 2014). But in the absence of any tributary flooding over the winter, conducting an HFE in fall 2021 could lead to small increases in water clarity (less turbid water) compared to if no-HFE were conducted. Analysis of historic turbidity data indicate marginal increase in water clarity without conducting an HFE would be very small. However, rainbow trout growth and predation rates decline exponentially as a function of turbidity. Therefore, because the relation between turbidity and rainbow trout growth is exponential, even small increases in water clarity associated with no-HFE, when compounded over several months, could have a significant effect on rainbow trout predation at the LCR. Maintaining elevated turbidity in Marble Canyon has been discussed as a potentially useful strategy for limiting rainbow trout populations and associated predation on native fish since at least 2011 (Runge and others 2011).

If conducting an HFE in fall 2021 does in fact lead to clearer water conditions around the LCR confluence area, then prior research demonstrates this would lead to improved feeding efficiency and predation by rainbow trout, and lower juvenile humpback chub survival rates until the next major turbidity flooding event resets and increases turbidity in Marble Canyon (Korman et al.

2020, Yackulic et al. 2018). In 2020 rainbow trout abundance around the LCR was ~700 (Yackulic personal communication). Although abundance estimates for 2021 are not yet available, catch rates have been slightly higher than 2020, suggesting rainbow trout abundance is now >700. For comparison, rainbow trout abundance at the LCR ranged from around 1500-2000 during 2012-2014, followed by near-zero abundance by 2017 owing to systemwide declines in rainbow trout. Thus, abundance of rainbow trout at the LCR confluence has been increasing since around 2017 but they remain below the abundance estimates seen during 2012-2014.

Nonnative Invasive Species - Green Sunfish

New cohorts of young nonnative green sunfish were observed by NPS staff in the Upper Slough within Glen Canyon in August of 2021. Due to this successful reproduction, NPS believes there is a need to remove green sunfish in the slough prior to an HFE. This removal effort is likely to eliminate most individuals and reduce the risk of downstream dispersal to critical endangered fish habitat downstream. Similar to previous removal efforts, most green sunfish removed and the few individuals that may remain will be new cohorts less than 40 mm TL. As done during the past two Octobers, NPS will pump out the 1/3-acre Upper Slough on October 6th-8th until only approximately 1,000 gallons of water remains. NPS will then electrofish and collect as many of the green sunfish as possible. These will be removed alive, quarantined for up to 14 days in one or more holding tanks, and then released into Lake Powell near the Wahweap Marina which is approximately 4 miles upstream of the dam. The release will be completed under an AGFD permit and will follow a Hazard Analysis and Critical Control Point (HAACP) plan to ensure no New Zealand mudsnails or other river invasives are released into Lake Powell. No adult or juvenile green sunfish were captured in the much larger Lower Slough in March or July of 2021 indicating that treatment of the main slough is unnecessary this year. Based on these planned actions and findings, green sunfish do not represent a resource concern within the Glen Canyon Reach that would prevent a fall 2021 HFE.

Within Grand Canyon, individual green sunfish were detected in several locations in 2021. Green sunfish identified in Kanab Creek in June were not detected again during monitoring in September in Kanab Creek which may have been due to a large tributary flood in July. Five were found while seining in a large backwater near RM 243 during July, but none were found in August in the same backwater. Four green sunfish were captured in the mainstem between RM 157-175 in 2021. No changes to the status of green sunfish in GRCA are expected related to the fall HFE. **No unacceptable risk to native fishes in Grand Canyon is expected, assuming green sunfish are removed from the Upper Slough in Glen Canyon prior to a fall HFE.**

Nonnative Invasive Species- Brown Trout

Brown trout are a highly piscivorous species known to prey on humpback chub and other native species (Yard et al. 2011). Monitoring of juvenile humpback chub suggests that increased rainbow trout abundances (and perhaps brown trout abundances) are associated with lower juvenile chub survival rates (Yackulic et al. 2018); however, this effect is uncertain and may be weak relative to other drivers of humpback chub dynamics (e.g., temperature, juvenile recruitment, food availability). Nonetheless, brown trout can have a population-level impact on native fish in Grand Canyon (Healy and others 2020). Coinciding with suppression of brown trout in a primary area for reproduction (Bright Angel Creek; Healy and others 2020), brown trout catches at the Little Colorado River confluence have generally been low since

implementation of the HFE protocol in 2012 (GCMRC unpublished data). In contrast to observations near the Little Colorado River, brown trout abundance has increased in Glen Canyon in recent years and is a cause for concern. This species is known to be a fall-spawner that has successfully spawned and recruited in Glen Canyon since implementation of the HFE protocol in 2012 (Runge et al. 2018).

The model of Lees Ferry brown trout populations developed for the 2018 Brown Trout report (Runge and others 2018) and subsequent updates suggest that high immigration of adult brown trout into the Lees Ferry reach has occurred concurrent with some, but not all, fall HFEs (i.e., 2014 and 2018 fall HFEs are associated with immigration of ~1000 medium-large brown trout into Lees Ferry, while HFEs in 2012, 2013, and 2016 are not associated with high immigration; Yackulic, 2021). Further, high immigration of adult brown trout into Lees Ferry has never occurred in the absence of a fall HFE. A separate model of brown trout movement developed using individually tagged brown trout throughout Marble and Grand Canyon also suggests that movement probability of brown trout to Lees Ferry is greater during intervals that include fall HFEs (Schelly and others 2021). Of 39 brown trout implanted with sonic tags in Glen Canyon, three fish were found to make temporary downstream movements – one of which was detected in the vicinity of the LCR, and then returned to Glen Canyon within ~5 days (NPS unpublished data). In addition, brown trout favor food items (Gammarus and mud snails) that have shown increases during the period where regular testing of fall HFEs has occurred (Kennedy and others 2019). Although fall HFEs have been linked to increased immigration of adult brown trout into Lees Ferry and potential food base changes that may also favor brown trout, there is not strong evidence linking fall HFEs to increased brown trout recruitment in Lees Ferry; specifically, years with the largest recruitment of new juvenile brown trout in Lees Ferry are not associated with fall HFEs (i.e., two of the highest years of brown trout recruitment are in fall of 2016 and fall of 2020, but no fall HFE occurred in the year prior). Increases in adult numbers due to immigration may lead to increased brown trout production. Nonetheless, a leading hypothesis suggests the number of rainbow trout spawners may influence brown trout recruitment through interference spawning. Since rainbow trout catch rates in 2021 are among the lowest observed in Glen Canyon since monitoring began, brown trout population growth may continue with or without a fall HFE in 2021.

Given increases in the estimated brown trout population, GLCA implemented an Incentivized Harvest program targeting brown trout in November of 2019. Nearly 480 larger sized (6” to 25”) brown trout have been harvested under this program to-date. GLCA has increased incentives this fall starting in August 2021, and is evaluating increases again in December and running through January 2022 to try and increase fishing pressure and harvest rates during the fall/winter spawning period when brown trout are most susceptible to angling. GLCA increased incentives last November of 2020 as well. The HFE may impact fishing in early November during the event, but fishing would return to normal after the HFE.

Although a November 2021 HFE may trigger additional immigration of adult brown trout into Lees Ferry, brown trout populations in Lees Ferry are large enough at this point that additional immigration arising from a 60-hour fall HFE is not anticipated to substantively increase the risk to endangered and native fishes that are downstream in Grand Canyon above the existing level of risk. Nonetheless, GLCA will increase incentives to anglers to

increase harvest of brown trout with the intent of offsetting the arrival of new adult brown trout that might immigrate into the reach during the fall.

Nonnative Invasive Species - Other Fishes

Other nonnative fish species observed in recent years in Glen Canyon that could threaten humpback chub and other native fishes if they became more abundant and widespread in the Colorado River downstream of Glen Canyon Dam include smallmouth bass, striped bass, and walleye. From 2017 to 2019 one smallmouth bass was captured each year below Glen Canyon Dam, while in recent years (2020-2021) no smallmouth bass were captured. Two smallmouth bass were caught in 2013 and one each in 2011 and 2010 (AGFD unpublished data). Striped bass have been periodically captured in Glen Canyon since the early 1980s. More recent captures of striped bass by GCMRC in Glen Canyon include one caught in each 2020 and 2016; two in 2019; and three in 2015. In August of 2018, five striped bass were observed in the slough (GCMRC unpublished data). Small numbers of walleye have been captured annually in Glen Canyon, from 2006 to 2019 AGFD has captured an average of three (range 0 to 8) each year (AGFD unpublished data, GCMRC unpublished data), primarily downstream of Glen Canyon Dam, with none captured in 2020 or in 2021. There is no evidence that detection of these non-native species in recent years is related to past HFEs.

If the current drought continues and reservoir levels continue to decline, there will be an increased risk of fish entrainment through Glen Canyon Dam and increased river temperatures conducive to warm water invasive fishes. At reservoir elevations below 3,516 ft, surface levels are less than 46 ft away from the penstock centerline and fish entrainment risk increases, as past monitoring has revealed that the majority of pelagic fish are located in the top 46 ft of the water column (U.S. Department of the Interior, 2012). In spring 2022, non-native fish will be in closer proximity to the penstocks than previous years when reservoir elevations were typically at least 50 feet higher. The elevated level of entrainment risk associated with low lake levels exists whether an HFE is conducted or not, but an HFE with a March, April, and May reallocation could somewhat exacerbate the risk by increasing the number of days spent near elevations where non-native fish entrainment is predicted to increase non-linearly (i.e., days around 3,516 ft). Reallocating the additional water released during a fall HFE to December, January, and February is not projected to increase the number of days below 3,525 ft under the most probable hydrology (see Table 2).

Riparian Vegetation

There is no evidence that the potential HFE would significantly impact riparian vegetation resources. The primary impact would be to extend the active channel upslope, which is the zone of daily inundation, for the duration of the HFE. This may slightly extend the suitable habitat for obligate wetland herbaceous species that respond positively to inundation, though longer-lived perennial species are unlikely to respond significantly to this short-term increase in inundation (Butterfield et al. 2018). Nonetheless, possible impacts of HFEs will be assessed through statistical modeling of changes in riparian vegetation composition based on 2018 vegetation surveys and hydrological variables, specifically inundation duration and elevation above base flows, calculated from the hydrograph and sandbar exceedance equations. These results will be available for discussion at the upcoming Annual Reporting Meeting. Future work to assess physiological responses of plants to HFEs will be conducted in the GCMRC FY21-23 TWP.

Water Delivery - Monthly, Daily, and Hourly Releases

The recommended HFE would result in changes to the weekly release prior to and after the HFE and the monthly volume distribution during Water Year (WY) 2022. Neither the tier determination nor the annual release volume as outlined in the 2007 Interim Guidelines for Lower Basin Shortages and Coordinated Operations for Lakes Powell and Mead (Reclamation 2007) will be impacted by a potential HFE. Reclamation determined the annual release volume for water year 2022 will be 7.48 million acre feet (maf) under all scenarios in accordance with the 2007 Interim Guidelines process and the August 2021 24-Month Study January 1, 2022 elevation projection.

The best estimate for total release from Glen Canyon Dam for an HFE in November 2021 is 33,900 cfs (18,900 cfs through the powerplant and 15,000 cfs of bypass). This estimate is based on the most recent unit testing completed in August 2021, a maintenance assumption that six of the eight units at Glen Canyon Powerplant would be available in November 2021, and an approximately 100% gate opening on the available six units. In addition, this estimate assumes that 40 megawatts (MW) (approximately 1,280 cfs) of system regulation will be maintained at Glen Canyon.

The release volume required in November for the potential 60-hour HFE is 634,000 acre-feet. The September 24-Month Study projected 500,000 acre feet release volume in November, therefore it is necessary to reallocate 134,000 acre-feet from March, April and May later in WY2022 into November. The November total volume for a 60-hour HFE is 634,000 acre-feet and bypass volume is 82,000 acre-feet. WAPA and Reclamation would coordinate on the scheduled reallocation of monthly release volumes with the goal of protecting minimum monthly thresholds for daily release fluctuations whenever practicable as described in the LTEMP as well as maximizing the economic value of hydropower. The September 24-Month Study most probable annual release for water year 2022 under Interim Guidelines is projected to be 7.48 maf, with all months projected to be above the minimum monthly thresholds regardless of the HFE release.

The LTEMP maximum ramp rates (4,000 cfs per hour when increasing and 2,500 cfs per hour when ramping down) would be adhered to throughout the experiment, as would the maximum daily fluctuations (9 times the monthly release volume in May and June; and 10 times the monthly release volume in July and August). Hourly releases for the days prior to and after the potential HFE in November are anticipated to fluctuate between 6,500 to 9,000 cfs and complies with the daily fluctuating range not to exceed 8,000 cfs outlined in the LTEMP. In addition, minimum releases of 5,000 cfs during the nighttime and 8,000 cfs during the daytime would be maintained.

Water Quality

A fall 2021 HFE does not pose unacceptable risks to water quality in the Colorado River and an HFE may help mitigate poor water quality (high temperature, low dissolved oxygen) depending on when lake turnover occurs. The bypass tubes withdraw water from deeper in the reservoir than the penstocks. As such, releasing water from the bypass tubes during an HFE temporarily cools water temperatures, increases dissolved oxygen concentrations, and

increases concentrations of dissolved nutrients. During an HFE, water temperatures in the Glen Canyon reach can be around 1-2°C colder compared to normal operations when the bypass tubes are not in operation. However, turnover in Lake Powell reservoir can occur from mid-October to November. Thus, depending on when lake turnover occurs, an HFE could mitigate undesired effects of warm water temperatures and low dissolved oxygen conditions. Notably, during an HFE, dissolved oxygen concentrations in the Glen Canyon reach are 100% saturated (no oxygen deficit), because the bypass tubes effectively aerate the water. Biologically available phosphorus concentrations have been increasing in the last year and are currently ~0.003 to 0.008 mg/L; this represents a moderate value relative to historical values. By releasing water from lower in the reservoir using the bypass tubes, an HFE would temporarily increase phosphorus concentrations; phosphorus concentrations are higher near the bypass tube elevation compared to the penstock elevation. For example, the 2018 HFE increased Glen Canyon's biologically available phosphorus concentration by ~ 0.001 mg/L from 0.003 to 0.004 mg/L. However, this effect of elevated phosphorus would be short-duration, and an HFE is unlikely to cause any long-term changes in phosphorus concentrations or loading to the Colorado River. Effects to downstream resources of an HFE arising from changes in phosphorus would therefore be negligible.

VII. Safety Considerations

Recreational Safety

As identified in the LTEMP HFE Protocol, potential effects on public health and safety could occur in conjunction with an HFE, primarily impacting recreational anglers and boaters. All daily fluctuations, minimum flows and maximum flows associated with any potential HFE are within the range experienced by recreational users in the past. Reclamation and NPS continue to work together to ensure that safety measures are implemented, including restricting access to the river immediately below the dam during potential HFEs, and, as noted below, providing public notice about the timing of the HFE implementation. NPS Boating Safety Rules always apply to all boaters using the river.

Reclamation and NPS coordinate to address safety and security issues related to HFEs. Additionally, the three park service units affected, GLCA, Grand Canyon National Park (GRCA) and Lake Mead National Recreation Area (LAKE) will work together to collaboratively plan necessary actions for a potential HFE. NPS units work to maximize continuity of efforts and resources, particularly in those areas where responsibilities are shared, specifically Lees Ferry and Pearce Ferry. Each park has clearly designated responsible parties and staffing needs and actions that need to occur prior to and during an HFE. The parks have coordinated communications plans, medical plans and resource capabilities for search and rescue responses. The three park units maintain frequent communication and information sharing leading up to and during any implemented HFE.

In preparation for an HFE, GRCA, GLCA, and LAKE identify and communicate with all commercial operations on the river, as well as permitted Colorado River trip permit holders that have the potential to be impacted by the HFE while rafting the Colorado River within each respective park unit. Planning is implemented to provide alternative trip dates for trips potentially affected by an HFE. All permit holders are directed to access up-to-date information provided by Reclamation, NPS, and the USGS/GCMRC websites. Additionally, all backcountry

hikers who access the Colorado River as part of their backcountry hike are alerted to potential campsite inundation areas.

Prior to an HFE, GLCA communicates with the holders of commercial use authorizations for commercial services (raft trips, back-haul services, fishing guides, etc.) on the Colorado River within GLCA to provide information on the time and duration of the HFE. During past HFEs, relatively few recreational boaters traveled upstream from Lees Ferry as the event was occurring. Information about a pending HFE and safety considerations are provided to recreational users at Lees Ferry in coordination with the PI Team Communications group. Information is also provided via public media, the GLCA website and on-site NPS staff. A fact sheet explaining potential impacts to park visitors is distributed to potentially affected visitors. Notifications are provided at Lees Ferry and Phantom Ranch and the fact sheet is available at these locations, as well as the GRCA Backcountry Information Center and primary visitor center.

Research and Monitoring Safety

In addition, safety considerations regarding sampling efforts by GCMRC have been incorporated into planning to ensure that safety of field staff is an overarching priority. GCMRC crews are deployed to locations in the days before the high flow release and will be supported by motorized rafts, and boats and cableways. They take critical measurements of discharge, suspended sediment transport, and organic drift. There is a lag between the time that water is released from the dam and the time that water arrives at a particular downstream location (often referred to as “travel time”). GCMRC crews deployed during the experimental flows would be made aware of the timing of the experimental flows. The range of flows for the potential HFE are within the range GCMRC and contracted boat operators have experienced in the past.

At sites downstream from the Paria River (river mile [RM] 1), work can only be safely conducted during daylight hours. This is especially the case on the first day of an HFE when the water surface typically is covered with woody debris that potentially can clog props of outboard engines or snag equipment suspended from cableways. Likewise, large logs that float just below the water surface can pose a threat to the safety of sampling staff. To address these issues, all field measurements by GCMRC personnel would be done during daylight hours in order to maximize the safety of field personnel.

VIII. Communications Plan

The potential HFE presents an opportunity to share with the public the purpose of the LTEMP flow-based experiments and anticipated benefits to resources downstream of Glen Canyon Dam. The communications plan for a fall 2021 HFE would consist of communications product development and media coordination; no public or media events at Glen Canyon Dam would be planned.

Prior to HFE implementation, Reclamation’s Upper Colorado Basin Region Public Affairs Office, in coordination with NPS, USGS, WAPA, and the Department, would develop a communications plan. In the instance of an event, an initial media advisory is sent to alert media representatives and the public of the HFE, including its purpose and expected start and finish dates. A more detailed news release for publication on or near the HFE dates may be prepared

for distribution by the Department. Social media outlets are also used to communicate with the public leading up to and during the experiment, including to share imagery of the HFE. If determined necessary, newsworthy and prudent by the communication team, the same methodology may be used if a determination not to conduct an HFE is made.

IX. Monitoring and Coordination During Experiment Implementation

If a fall 2021 HFE is implemented, members of the PI Team would continue to meet regularly prior to, during, and following implementation of the experiment. Ongoing communication occurs through the regularly scheduled monthly Glen Canyon Dam operations coordination calls and through additional coordination calls, as needed. Scientists conducting field surveys during the experiment and agency technical experts report on data collected and preliminary results to the Department and the GCDAMP at regularly scheduled meetings. Glen Canyon Dam operations may be adjusted in the event of unexpected impacts from the HFE and/or in the event of an emergency.

X. Post Experiment-Reporting and Planning for Future Experiments

The PI Team would coordinate to report initial findings at the 2021 GCDAMP Annual Reporting Meeting, scheduled for January 2022 in Phoenix, AZ.

- Monitoring of sediment transport and sandbar responses to an HFE would include measuring sediment transport at several sites in Marble and Grand Canyons as well as the volume of fine sediment, especially sand, that is stored on the bed of the Colorado River, in its eddies, or at higher elevation along the river's banks as described in Section V. GCMRC will also collect data on water quality (including nutrients), native and nonnative fishes, aquatic invertebrates, riparian plants, and other resources as described in the Reclamation and GCMRC TWP (DOI 2020). GCMRC will use the information from these studies to evaluate the effects of HFEs on downstream resources in Glen, Marble, and Grand Canyons and to help in the design of future experiments.
- The 7.48 maf releases will lead to lower sand export compared to typical 8.23 maf release patterns. Based on a recent *Journal of Geophysical Research* article by Topping and others (2021) and analyses from the sand routing model used to trigger HFEs, if an HFE does not occur this November, it is likely that roughly 1.2 million metric tons (~90%) of this year's Paria River sand inputs would be retained somewhere in Marble Canyon until at least June 30, 2022.
- The PI Team would meet in early 2022 to review the implementation and results of all 2021 experimental activities, and to begin coordination on the evaluation of resources and potential experiments that may be conducted in 2022.
- In addition, the PI Team would report ongoing findings at meetings of the GCDAMP Technical Work Group (TWG) and Adaptive Management Work Group (AMWG). Reclamation has a commitment to provide an annual monitoring report to the FWS Arizona Ecological Services Office (AESO) in compliance with the 2016 Biological Opinion; this report will also include a summary of the effects of any flow experiments conducted under the LTEMP ROD. Reclamation would use the monitoring information

and feedback from AESO and GCDAMP stakeholders to inform monitoring for future experiments, and to design and implement any measures necessary to address any adverse effects that may occur due to these flows.

- In accordance with the LTEMP, the Department may make the decision to conduct future flow-based experiments (High Flow Experiments, Bug Flows, Trout Management Flows, and Low Summer Flows) at Glen Canyon Dam if it is determined that there are no unacceptable adverse impacts on other resource conditions. Information and data from this or other experiments will be considered in future recommendations and decisions.

XI. Consultation

Reclamation and GCMRC provided much of the information in this report that was available at the time in the GCMRC 2020 Annual Report. Newer not yet published information was reported to the Adaptive Management Program Partners at the GCDAMP Annual Reporting Meeting on January 20-22, 2021 as well as to the AMWG at its August 18-19, 2021 meeting. Notification of the potential for a 2021 fall HFE was communicated to GCDAMP stakeholders on August 19, 2021, shortly after the sediment trigger was met. A follow-up informational webinar was held on September 21, 2021 with GCDAMP stakeholders as an opportunity to ask questions and provide feedback. Representatives from the Basin States participated in the development of this recommendation. Based on feedback and discussion at the GCDAMP webinar, the recommendation and report were finalized without major changes. Reclamation and GCMRC will present the findings and recommendation of this report to the TWG on October 13, 2021.

On September 20, 2021, the required 30-day advance notification and offer for consultation were sent electronically to the Tribes and Parties to the LTEMP cultural Programmatic Agreement regarding the potential for a High Flow Experiment beginning November 1, 2021. As of September 24, 2021, Reclamation has not received any requests for consultation on the potential experiment. A follow-up notification will be sent electronically to the Programmatic Agreement signatories, including Tribes, following the Department's decision regarding the potential High Flow Experiment.

XII. Conclusion

Determining whether to recommend an HFE required coordination and effective communication among the technical staff of multiple agencies. The PI Team relied heavily on and would like to acknowledge the contributions of staff from GCMRC, Reclamation, and WAPA in making this recommendation.

The PI Team has thoroughly evaluated the issues discussed above and has taken into consideration the information and analyses included in the LTEMP FEIS and ROD. The PI Team's recommendation regarding implementation of a fall 2021 HFE is based on a careful assessment of resources and best available science. **By consensus, the PI Team is opposed to recommending that the Department implement a 192-hour extended duration HFE in fall 2021. Technical representatives were divided in the assessment of a shorter duration 60-hour HFE.**

References Cited

- Blinn, D.W., and Cole, G.A., 1991. Algal and invertebrate biota in the Colorado River: comparison of pre-and post-dam conditions. *Colorado River ecology and dam management*. National Academy Press, Washington, DC, p102-123.
- Bunch, A.J., Osterhoudt, R.C., Anderson, M.C., and Stewart, W.T. 2012. Colorado River Fish Monitoring in Grand Canyon, Arizona— 2012 Annual Report. Final report prepared by Arizona Game and Fish Department for Grand Canyon Monitoring and Research Center, Flagstaff, Arizona.
- Burnham, K.P., and Anderson, D.R. 2002 Model selection and multimodel inference: a practical information-theoretic approach. New York, New York, Springer-Verlag.
- Butterfield, B.J., Palmquist, E.C., and Ralston, B.E.. 2018, Hydrological regime and climate interactively shape riparian vegetation composition along the Colorado River, Grand Canyon: *Journal of Applied Vegetation Science*: 2018:1-12. Dietze, M.C., 2017, *Ecological forecasting*: Princeton, New Jersey, USA, Princeton University Press, 288 p.
- Carlisle, D.M., Grantham, T.E., Eng, K. and Wolock, D.M., 2017. Biological relevance of streamflow metrics: regional and national perspectives. *Freshwater Science*, 36(4), pp.927-940.
- Cross, W.F., Baxter, C.V., Rosi-Marshall, E.J., Hall Jr, R.O., Kennedy, T.A., Donner, K.C., ... & Yard, M.D., 2013. Food-web dynamics in a large river discontinuum. *Ecological Monographs*, 83(3), pp.311-337.
- Dibble, K.L., C.B. Yackulic, T.A. Kennedy, K.R. Bestgen, and J.C. Schmidt., 2021, Water storage decisions will determine the distribution and persistence of imperiled river fishes. *Ecological Applications* 31, no. 2 (2021): e02279.
- East, A.E., Collins, B.D., Sankey, J.B., Corbett, S.C., Fairley, H.C., and Caster, J., 2016, Conditions and processes affecting sand resources at archeological sites in the Colorado River corridor below Glen Canyon Dam, Arizona: U.S. Geological Survey Professional Paper 1825, 104 p., <http://dx.doi.org/10.3133/pp1825>.
- Grams, P. E., Tusso, R. B., & Buscombe, D. (2018). Automated Remote Cameras for Monitoring Alluvial Sandbars on the Colorado River in Grand Canyon, Arizona. U.S. Geological Survey Open-File Report 2018-1019, 61. <https://doi.org/10.3133/ofr20181019>
- Grams, P.E., 2019, Sandbar deposition caused by high-flow experiments on the Colorado River downstream from Glen Canyon Dam—November 2012–November 2018, in High-flow experiments assessment extended abstracts—Glen Canyon Dam Adaptive Management Program Annual Reporting Meeting presentations, March 12-13, 2019, Phoenix, Ariz.: U.S. Geological Survey, Grand Canyon Monitoring and Research Center, p. 12-22, https://www.usbr.gov/uc/progact/amp/amwg/2019-03-06-amwg-meeting/20190301-HFE_Extended_Abstracts-Combined_FINAL.pdf.

Hadley, D. R., Grams, P. E., & Kaplinski, M. A. 2018a. Quantifying geomorphic and vegetation change at sandbar campsites in response to flow regulation and controlled floods, Grand Canyon National Park, Arizona. *River Research and Applications*, (June), 1–11.

<https://doi.org/10.1002/rra.3349>

Hadley, D. R., Grams, P. E., Kaplinski, M. A., Hazel, J.E., J., & Parnell, R. A. 2018b. Geomorphology and vegetation change at Colorado River campsites, Marble and Grand Canyons, Arizona. U.S. Geological Survey Scientific Investigations Report 2017–5096, 64.

<https://doi.org/10.3133/sir20175096>

Healy, B. D., R. C. Schelly, C. B. Yackulic, E. C. O. Smith, and P. Budy. 2020. Remarkable response of native fishes to invasive trout suppression varies with trout density, temperature, and annual hydrology. *Canadian Journal of Fisheries and Aquatic Sciences* 77, pp.1446–1462.

Kegerries, R. B., B. C. Albrecht, E.I. Gilbert, W. H. Brandenburg, A.L. Barkalow, M.C. McKinstry, H.E. Mohn, B.D. Healy, J.R. Stolberg, E.C. Omana Smith, C.B. Nelson, and R.J. Rogers. 2017a. Occurrence and Reproduction by Razorback Sucker (*Xyrauchen texanus*) in the Grand Canyon, Arizona. *The Southwestern Naturalist* 62(3): 227–232.

Kegerries, R.B., R.J. Rogers, B. Albrecht, S.L. Wood, M.J. Chavez, S.P. Platania, M. McKinstry, K. Pedersen, B. Healy, J. Stolberg, and E. Omana Smith. 2021. Razorback Sucker *Xyrauchen texanus* research and monitoring in the Colorado River inflow area of Lake Mead and the lower Grand Canyon, Arizona and Nevada. Report prepared by BIO-WEST, Inc., for the US Bureau of Reclamation, Upper Colorado Region, Salt Lake City.

Kennedy, T. A., and B. E. Ralston. 2011. Biological responses to high-flow experiments at Glen Canyon Dam. Effects of three high-flow experiments on the Colorado River ecosystem downstream from Glen Canyon Dam, Arizona. US Geological Survey Circular, 1366(147), pp. 93-125.

Kennedy, T.A., Muehlbauer, J.D., Yackulic, C.B., Lytle, D.A., Miller, S.W., Dibble, K.L., Kortenhoeven, E.W., Metcalfe, A.N., and Baxter, C.V. 2016, Flow management for hydropower extirpates aquatic insects, undermining river food webs: *BioScience*, v. 66, no. 7, p. 561-575,

<http://dx.doi.org/10.1093/biosci/biw059>.

Kennedy, T.A., and J.D. Muehlbauer, 2019, Big flood, small flood, spring flood, fall flood: HFE timing affects food base response. Presentation at 2019 Annual Reporting Meeting and HFE workshop, Phoenix, AZ. [https://www.usbr.gov/uc/progact/amp/twg/2019-03-14-twg-meeting/20190314-](https://www.usbr.gov/uc/progact/amp/twg/2019-03-14-twg-meeting/20190314-BigFloodSmallFloodSpringFloodFallFloodHFETimingAffectsFoodBaseResponse-Presentation-508-UCRO.pdf)

[BigFloodSmallFloodSpringFloodFallFloodHFETimingAffectsFoodBaseResponse-Presentation-508-UCRO.pdf](https://www.usbr.gov/uc/progact/amp/twg/2019-03-14-twg-meeting/20190314-BigFloodSmallFloodSpringFloodFallFloodHFETimingAffectsFoodBaseResponse-Presentation-508-UCRO.pdf)

Korman, J., and Campana, S.E. 2009. Effects of hydropeaking on nearshore habitat use and growth of age-0 rainbow trout in a large regulated river. *Transactions of the American Fisheries Society*, v. 138, no. 1, p. 76-87, <http://dx.doi.org/10.1577/T08-026.1>.

Korman, J., Kaplinski, M., and Buszowski, J. 2006. Effects of air and mainstem water temperatures, hydraulic isolation, and fluctuating flows from the Glen Canyon Dam on the water temperatures in shoreline environments of the Colorado River in Grand Canyon: Flagstaff, Arizona, Ecometric Research Inc., Northern Arizona University, and Namtek Inc., submitted to U.S. Geological Survey, Grand Canyon Monitoring and Research Center, cooperative agreement no. 04WRAG00006, modification 1, 52 p.

Korman, J., Walters, C., Martell, S.J.D., Pine, W.E., III, and Dutterer, A. 2011. Effects of flow fluctuations on habitat use and survival of age-0 rainbow trout (*Oncorhynchus mykiss*) in a large, regulated river: Canadian Journal of Fisheries and Aquatic Sciences, v. 68, no. 6, p. 1097-1109, <http://dx.doi.org/10.1139/f2011-045>.

Korman, J., Yard, M., Walters, C.J., and Coggins, L.G. 2009. Effects of fish size, habitat, flow, and density on capture probabilities of age-0 rainbow trout estimated from electrofishing at discrete sites in a large river. Transactions of the American Fisheries Society, v. 138, no. 1, p. 58-75, <http://dx.doi.org/10.1577/T08-025.1>.

Mueller, E. R., Grams, P. E., Hazel, J. E., & Schmidt, J. C. 2018. Variability in eddy sandbar dynamics during two decades of controlled flooding of the Colorado River in the Grand Canyon. Sedimentary Geology, 363, 181–199. <https://doi.org/10.1016/j.sedgeo.2017.11.007>

Robinson, A.T., Clarkson, R.W., and Forrest, R.E. 1998. Dispersal of larval fishes in a regulated river tributary. Transactions of the American Fisheries Society, v. 127, no. 5, p. 772-786, [https://doi.org/10.1577/1548-8659\(1998\)127<0772:DOLFIA>2.0.CO;2](https://doi.org/10.1577/1548-8659(1998)127<0772:DOLFIA>2.0.CO;2).

Poff, N.L., Allan, J.D., Bain, M.B., Karr, J.R., Prestegard, K.L., Richter, B.D., Sparks, R.E. and Stromberg, J.C., 1997. The natural flow regime. *BioScience*, 47(11), pp.769-784.

Rogowski, D.L, R.J. Osterhoudt, H.E. Mohn, and J.K. Boyer. 2018. Humpback chub (*Gila cypha*) range expansion in the Western Grand Canyon. Western North American Naturalist 78: 26-38.

Robinson, A.T., Clarkson, R.W., and Forrest, R.E. 1998. Dispersal of larval fishes in a regulated river tributary. Transactions of the American Fisheries Society, v. 127, no. 5, p. 772-786, [https://doi.org/10.1577/1548-8659\(1998\)127<0772:DOLFIA>2.0.CO;2](https://doi.org/10.1577/1548-8659(1998)127<0772:DOLFIA>2.0.CO;2).

Ross, R., and Grams, P.E. 2013. Nearshore thermal gradients of the Colorado River near the Little Colorado River confluence, Grand Canyon National Park, Arizona. U.S. Geological Survey Open-File Report 2013-1013, 65 p., <http://pubs.usgs.gov/of/2013/1013/>.

Runge, M.C., Bean, Ellen, Smith, D.R., and Kokos, Sonja, 2011, Non-native fish control below Glen Canyon Dam—Report from a structured decision-making project: U.S. Geological Survey Open-File Report 2011–1012, 74 p., at <https://pubs.usgs.gov/of/2011/1012/>.

Runge, M.C., Yackulic, C.B., Bair, L.S., Kennedy, T.A., Valdez, R.A., Ellsworth, C., Kershner J.L., Rogers, R.S., Trammell, M.A., and Young, K.L. 2018. Brown trout in the Lees Ferry reach

of the Colorado River—Evaluation of causal hypotheses and potential interventions. U.S. Geological Survey Open-File Report 2018–1069, 83 p., <https://doi.org/10.3133/ofr20181069>. Sabo, J.L., and Post, D.M. 2008. Quantifying periodic, stochastic, and catastrophic environmental variation. *Ecological Monographs*, v. 78, no. 1, p. 19–40.

Sankey, J.B., Caster, J., Kasprak, A. and East, A.E. 2018. The response of source-bordering aeolian dunefields to sediment-supply changes 2. Controlled floods of the Colorado River in Grand Canyon, Arizona, USA. *Aeolian research*, 32, pp.154-169. <https://doi.org/10.1016/j.aeolia.2018.02.004>

Schmidt, J. C. and Grams, P. E. 2011. The high flows--physical science results. In T. S. Melis (Ed.), *Effects of three high-flow experiments on the Colorado River ecosystem downstream from Glen Canyon Dam, Arizona, U.S. Geological Survey Circular 1366* (pp. 53–91). Retrieved from <https://pubs.usgs.gov/circ/1366/>

Stevens, L.E., Shannon, J.P., and Blinn, D.W., 1997. Colorado River benthic ecology in Grand Canyon, Arizona, USA: dam, tributary and geomorphological influences. *Regulated Rivers: Research & Management* 13(2), pp.129-149.

Topping, D.J., Rubin, D.M., Nelson, J.M., Kinzel III, P.J. and Corson, I.C., 2000. Colorado River sediment transport: 2. Systematic Bed-elevation and grain-size effects of sand supply limitation. *Water Resources Research*, 36(2), pp.543-570.

Topping, D. J., Schmidt, J. C., & Vierra, L. E., 2003. Computation and analysis of the instantaneous-discharge record for the Colorado River at Lees Ferry, Arizona: May 8, 1921, through September 30, 2000. USGS Professional Paper 1677.

Topping, D. J., Grams, P. E., Griffiths, R. E., Dean, D. J., Wright, S. A., & Unema, J. A., 2021. Self-limitation of sand storage in a bedrock-canyon river arising from the interaction of flow and grain size. *Journal of Geophysical Research: Earth Surface*, 126(5), e2020JF005565.

U.S. Department of the Interior. 2011. Environmental assessment—Development and implementation of a protocol for high-flow experimental releases from Glen Canyon Dam, Arizona, 2011 through 2020: Salt Lake City, Utah, Bureau of Reclamation, Upper Colorado Region, 176 p. plus appendices, <http://www.usbr.gov/uc/envdocs/ea/gc/HFEProtocol/HFEEA.pdf>.

U.S. Department of Interior. 2016a. Glen Canyon Dam Long-term Experimental and Management Plan final Environmental Impact Statement (LTEMP FEIS): U.S. Department of the Interior, Bureau of Reclamation, Upper Colorado Region, National Park Service, Intermountain Region, 8 chapters plus 17 appendices, <http://ltempeis.anl.gov/documents/final-eis/>.

U.S. Department of Interior. 2016b. Record of Decision for the Glen Canyon Dam Long-term Experimental and Management Plan final Environmental Impact Statement (LTEMP ROD): Salt Lake City, Utah, U.S. Department of the Interior, Bureau of Reclamation, Upper

Colorado Region, National Park Service, Intermountain Region, 196 p., http://itempeis.anl.gov/documents/docs/LTEMP_ROD.pdf.

U.S. Department of the Interior. 2020. Glen Canyon Dam Adaptive Management Program Triennial Budget and Work Plan—Fiscal Years 2021-2023—December 22, 2020 final signed by the Secretary of the Dept. of Interior: Flagstaff, Ariz., U.S. Geological Survey, Grand Canyon Monitoring and Research Center and Salt Lake City, Utah, Bureau of Reclamation, Upper Colorado Region, 384 p.

VanderKooi, S.P., Kennedy, T.A., Topping, D.J., Grams, P.E., Ward, D.L., Fairley, H.C., Bair, L.S., Yackulic, C.B., Schmidt, J.C., and Sankey, J.B. 2017. Scientific monitoring plan in support of the selected alternative of the Glen Canyon Dam Long-Term Experimental and Management Plan: U.S. Geological Survey, Grand Canyon Monitoring and Research Center, U.S. Geological Survey Open-File Report 2017-1006, 18 p., <https://doi.org/10.3133/ofr2017100>

Van Haverbeke, D.R., D.M. Stone, M.J. Dodrill, K.L. Young, and M.J. Pillow. 2017. Population expansion of humpback chub in western Grand Canyon and hypothesized mechanisms. *The Southwestern Naturalist* 62: 285-292.

Voichick, N., & Topping, D. J. 2014. Extending the turbidity record—making additional use of continuous data from turbidity, acoustic-Doppler, and laser diffraction instruments and suspended-sediment samples in the Colorado River in Grand Canyon. US Geological Survey Scientific Investigations Report, 5097, 31.

Voichick, N., Kennedy, T.A., Topping, D.J., Griffiths, R.E., and Fry, K.L., 2016, Water clarity of the Colorado River—Implications for food webs and fish communities: U.S. Geological Survey Fact Sheet 2016–5053, 4 p., <http://dx.doi.org/10.3133/fs20163053>

Ward, D.L. 2015. Green Sunfish *Lepomis cyanellus*; Risk Assessment for the Colorado River ecosystem. Memorandum submitted to Katrina Grantz, Bureau of Reclamation, Upper Colorado Region, Salt Lake City.

Yackulic, C. B., Yard, M.D., Korman, J., & Van Haverbeke, D.R., 2014. A quantitative life history of endangered humpback chub that spawn in the Little Colorado River: variation in movement, growth, and survival. *Ecology and Evolution*, 4(7), pp.1006-1018.

Yackulic, C.B., Dibble, K., and Deemer, B., 2018. Temperature and nutrients as aquatic ecosystem drivers in the CRe. Glen Canyon Dam Adaptive Management Work Group, Annual Reporting Meeting <https://www.usbr.gov/uc/progact/amp/twg/2018-01-25-twg-meeting/AR11.pdf>